

## Original Research Article

# Role of Magnetic Resonance Imaging in Evaluation of Hip and Groin Sports- Related Injuries

### Abstract

**Background:** Magnetic Resonance Imaging (MRI) proved to be highly accurate imaging modality allows clear differentiation of the normal anatomic features of the hip. In addition to that it allows clear differentiation of the individual component of the normal joint from one another. The purpose of this work is was to diagnose sports-related injuries of hip and groin using MR imaging with different sequences.

**Methods:** This prospective study was carried out on 59 athletic subjects presented clinically with different hip or groin joint complaints, both sexes within adult age. All patients were subjected to clinical examination, imaging (conventional radiography of both hips were performed mostly in antero-posterior and lateral views to detect fractures, dislocations, bony lesions or soft tissue swelling), MRI examinations and image analysis.

**Results:** There was significant difference between MRI type of injuries and different age groups of studied sportive patient. MRI show lesion in 59 cases in comparison to X-ray which has less sensitivity as 46 case has negative results show lesions in MRI.

**Conclusions:** MRI of the hip has established to be the gold standard for the diagnosis of sports-related hip and groin injuries in the setting of negative radiographs, also it can determine early signs of stress fracture before it appear in X-ray. so, more details about different lesions can be obtained from MRI and This help in accurate diagnosis, rapid treatment of different athlete lesions and determine time plane of returning to the sport by follow up imaging.

**Keywords:** Magnetic Resonance Imaging, Evaluation of Hip and Groin, Sports- Related Injuries

## **Introduction:**

Groin is the region between the abdomen and the legs, groin and hip pain in Athletes is one of the most difficult clinical entities to diagnose and treat in sports medicine. This is due to the amount of differential diagnoses and the complexity of the groin and hip region's anatomy, where many structures interact.<sup>[1]</sup>

The understanding, assessment, and management of hip and groin pain in athletes have improved significantly over the last decade. In the past, the assessment of hip pain and groin injuries were limited to obvious disorders. The progress have improved awareness of the complexities of groin and hip athletic injuries, as well as the biomechanical basis for recent hip abnormalities.<sup>[2]</sup>

Magnetic Resonance Imaging (MRI) proved to be highly accurate imaging modality allows clear differentiation of the normal anatomic features of the hip. In addition to that it allows clear differentiation of the individual component of the normal joint from one another. As MRI allows identification of the muscles, bone marrow, cortex, fascia, vessels and nerves with high contrast between them. Although MRI has no ionizing radiation and thus no biological hazards have been identified<sup>[3]</sup>.

Local injections can be beneficial in some cases, allowing high level athletes to continue competing throughout the season. To prevent aggravating hip pain in athletes, avoid violent stretching to increase range of motion. Also circumvent deep hip flexion and low repetition and heavyweight strength training. Once the athlete is no longer experiencing pain, a gradual return to play can be started<sup>[2]</sup>.

Radiology is play a critical role in guiding the monitoring of athletic injuries. It is essential to understand the benefits and drawbacks of imaging modalities as it helps in accurate and timely diagnosis.

MRI is the best evaluation technique for the different injuries, Despite the fact that the initial evaluation with conventional radiography and the use of other imaging techniques as ultrasonography (US), computed tomography, and bone scintigraphy may be helpful or preferred in certain cases <sup>[4]</sup>.

Because the hips are major weight-bearing joints, athletes depend on them to perform well in their respective sports activities. Hip pain is a frequent complaint among athletes as a result of acute and chronic injuries affecting the hip.

Recent progress in imaging approaches aid radiologists in their understanding of hip anatomy, biomechanics, and pathologic conditions, thereby assisting our orthopaedic surgeon colleagues in the early management of these cases <sup>[4]</sup>.

The purpose of this study was to diagnose sports-related injuries of hip and groin using MR imaging with different sequences.

### **Patients and Methods:**

This was a prospective study that carried out on 59 athletic cases presented clinically with different hip or groin joint complaints, both sexes within adult age. The study was done after approval from the Ethical Committee Tanta University Hospitals. All patients provided informed consent.

Exclusion criteria: Participants who hadn't sport injury, history of hip trauma, hip joint malignancy, contraindicated to MRI, with metal implants, such as hip replacements, vascular clips, or foreign bodies and those who suffer from severe claustrophobia may be unable to

tolerate an MRI scan, although more open scanners are now available, and medical sedation (which is not needed in our study) is available of making the test more bearable.

**All patients were subjected to:** Full history taking , clinical examination, imaging (conventional radiography of both hips were performed mostly in antero-posterior and lateral views to detect fractures, dislocations, bony lesions or soft tissue swelling), MRI examinations and image analysis.

**MRI protocols: -**

Both hips were examined for suspected bilateral deformities using the body coil with a wide field of view to determine the extent of the lesions and to enable for comparison between the normal and abnormal sides. The thickness of the slices was adjusted to be between 4 and 8mm, and thinner slices were used when additional detail was needed. Scout observe of the hip: In all cases, a coronal T1 weighted image was taken to aid in the localization of subsequent slices of various planes and to delineate the anatomy. A repetition time (TR) < 800 msec, an echo time (TE) < 30 msec, a slice thickness of 4-8 mm, FOV= 30-40 cm. Coronal and Axial T1 weighted spin echo images: A repetition time (TR) < 800 msec, an echo time (TE) < 30 msec, a slice thickness of 4-8 mm, FOV= 30-40 cm. Sagittal, axial and coronal T2 weighted spin echo images: repetition time (TR) >2000 msec, an echo time (TE) > 60 msec, a slice thickness of 4-8 mm, FOV= 30-40 cm. Coronal and axial STIR (short time inversion recovery) weighted images: A repetition time (TR) > 2000 msec, an echo time (TE) > 30 msec, a slice thickness of 4-8 mm, FOV= 30-40 cm.

**Image analysis:**

Interpretation was done regarding assessment of the followings: hip joint alignment on both sides to determine femoral head acetabular articulation and exclude femoral head dislocation. Femoral heads regarding size, shape and the signal intensity. Femoral head marrow signal intensity was estimated in both T1 & T2WIS and graded as normal, increased or decreased

judged relative to the contralateral femoral head. The joint space for presence or absence of joint effusion and is graded into mild, moderate and severe as in labrum tear. The pelvic bones and the upper femur, for exclusion of any bone fractures, edema and abnormal shape and assess the signal intensity of the bone marrow and evaluate any abnormal signal in all pulse sequences. The periarticular soft tissues for any abnormal signal or any abnormal swelling. The surrounding muscles and intra pelvic soft tissue structures for the presence of any abnormal signal for exclusion of muscle tear, contusion or strain

### Statistical method

The data were arranged, tabulated and statistically analyzed by SPSS software (Statistical Package for the Social Sciences, version 21, SPSS Inc. Chicago, IL, USA). The range, mean, and standard deviation were calculated for quantitative data. For qualitative data, which characterize a categorical set of data by the frequency, percentage, or proportion of each category, the Chi-square test was used to compare two groups and more. Significance was accepted at  $p < 0.05$  for explanation of results of tests of significance.

### Results:

**Error! Not a valid bookmark self-reference.** showed patient characteristics and complaint of the studied patients.

**Table 1: Age, sex, affected side and complaint of the studied patients who play sports and presented with persistent or recurrent hip or groin pain (n=59).**

Variables		The studied patients (n=59)
Sex	Female	25(42.4%)
	Male	34(57.6%)
Age years	14-18	9(15.3%)
	>18-25	16(27.1%)
	>25-45	22(37.3%)
	>45-65	12(20.3%)
	Range	14-65
	Mean±SD	31.37±13.40
	Median	30.00
Side affected	Right	32(54.2%)
	Left	21(35.6%)
	Right & left	6(10.2%)

Complaint	
Pain	14(23.7%)
Pain-catching & locking	2(3.4%)
Pain tenderness	17(28.8%)
Pain & snap with flexion	2(3.4%)
Pain & swelling	2(3.4%)
Pain & limitation of movement	20(33.9%)
Pain & limitation of movement & swelling	2(3.4%)

Data are represented by mean± SD or number (%)

Table 2 showed MRI types of injuries found in 59 studied sportive patients

**Table 2: MRI types of injuries found in 59 studied sportive patients**

MRI types of injuries	The studied patients (n=59)
Bone injuries	14(23.7%)
Joint injuries	6(10.2%)
Muscle injuries	36(61.0%)
Bone and muscle injuries	1(1.7%)
Joint and muscle injuries	2(3.4%)

Data are represented by number (%)

There was significant difference between MRI type of injuries and different age groups of studied sportive patient. Table 3

**Table 3: Relation between MRI type of injuries and different age groups of studied sportive patients (n=59)**

MRI types of injuries found in studied patients	Age years of the studied patients (n=59)				$\chi^2$	P
	14-18 (n=9)	>18-25 (n=16)	>25-45 (n=22)	>45-65 (n=12)		
Bone injuries	6(66.7%)	6(37.5%)	0(0%)	2(16.7%)	35.984	0.0001*
Bone and muscle injuries	1(11.1%)	0(0%)	0(0%)	0(0%)		
Joint injuries	0(0%)	0(0%)	6(27.3%)	0(0%)		
Muscle injuries	2(22.2%)	10(62.5%)	14(63.6%)	10(83.3%)		
Joint and muscle injuries	0(0%)	0(0%)	2(9.1%)	0(0%)		

\*: Statistically significant at  $p \leq 0.05$ , Data are represented by number (%)

Table 4 showed final detailed diagnosis of lesions by MRI of the studied patients who play sports and presented with persistent or recurrent hip or groin pain and relation between final MRI finding and X-ray finding in 59 sportive patients with painful hip (n=59).

**Table 4: Final detailed diagnosis of lesions by MRI of the studied patients who play sports and presented with persistent or recurrent hip or groin pain (n=59) and Relation between final MRI finding and X-ray finding in 59 sportive patients with painful hip**

Final MRI finding in investigated patients	The incidence (n=62)
--	----------------------

<b>bone injuries:</b>	<b>(n=15)</b>	
Anterior inferior iliac spine avulsion	6(40.0%)	
Femoral neck stress fracture	4(26.7%)	
Greater trochantric fracture	2(13.3%)	
Anterior superior iliac spine avulsion	1(6.7%)	
Thigh splint (Adductor insertion Avulsion)	2(13.3%)	
<b>Joint injuries:</b>	<b>(n=8)</b>	
Femero - acetabular impingment	2(25.0%)	
Labrum tear	6(75.0%)	
<b>Muscle injuries:</b>	<b>(n=39)</b>	
Greater trochanter syndrome	12(30.8%)	
Muscle contusion	2(5.1%)	
Snapping hip syndrome	2(5.1%)	
athletic publalgia	12(30.8%)	
Strain	11(28.2%)	
Final MRI finding in investigated patients	<b>Patient do X –ray n=59</b>	
	<b>Positive X- ray n=13</b>	<b>Negative X-ray n=46</b>
Anterior inferior iliac spine avulsion	5	1
Femoral neck stress fracture	1	3
Greater trochantric fracture	2	0
Anterior superior iliac spine avulsion with muscle strain	1	0
Thigh splint (Adductor insertion Avulsion)	2	0
Femero - acetabular impingment	1	1
Labrum tear	0	4
Labrum tear with muscle strain	0	2
Greater trochanter syndrome	0	12
Muscle contusion	0	2
Snapping hip syndrome	0	2
Athletic publalgia	0	12
Muscle Strain	1	7

Data are represented by number (%), MRI: Magnetic Resonance Imaging

MRI show lesion in 59 cases in comparison to X-ray which has less sensitivity as 46 case has negative results show lesions in MRI. Table 5

**Table 5: MRI signs in different sequences in investigated sportive patient with painful hip**

MRI	T1 weighted image	T2weighted image	STIR image	No. of cases
<b>Bone marrow edema</b>	Hypo intense	Hyper intense	Hyper intense	25
<b>Fracture line</b>	Hypo intense	Hypo intense Surrounded by high signal	Hypo intense Surrounded by high signal	11
<b>Muscle strain</b>	Hypo intense	Hyper intense	Hyper intense	23
<b>Grade 1: edema</b>	Hypo intense	Hyper intense with partial retracted	Hyper intense with partial retracted	3
<b>Grade 2: partial</b>				16

<b>tear</b>	Hypo intense	fiber	fiber	
<b>Grade3: complete tear</b>		Hyper intense with complete retracted muscle	Hyper intense with complete retracted muscle	4
<b>Tendnosis</b>	Hypo intense	Hyper intense	Hyper intense	14
<b>Intermuscular hemorrhage</b>	Hypo intense	Hyper intense	Hyper intense	7
<b>Labrum tear</b>	Non- specific labrum hypo intense in T1	Hypo intense area within hyper intense labrum	Hypo intense area within hyper intense labrum	6
<b>Joint effusion</b>	Hypo intense	Hyper intense	Hyper intense	11

## Discussion

Hip and groin pain is an usual complaint among all ages of athletes and has been estimated to account for 5% to 6% of all types of sports injuries. Hip and groin pain are frequently caused by a combination of factors, including bony and soft tissue abnormalities that can occur intra- or extra-articularly in both acute and chronic situations of overuse<sup>[5]</sup>.

In this study according to MRI types of injuries there are 62 lesions in 59 patients, 14 patients showed bone injuries (23.7 %), 6 patients with joint injuries (10.2%) and 36 patients with muscle injuries (61%), Some of patient had multiple injuries 1 patient had bone and muscle injuries (1.7%) and 2 patient had joint and muscle injuries (3.4%). This result showed that muscular injuries which represent extra-articular injuries is the most common lesion in 39 patients (66.1%) .

This agree with Chan et al.<sup>[6]</sup> who had study showed that the majority of the injuries were mild to moderate and the most frequently occurring ones were abrasion (37%), contusion (21%), cramp (20%), sprains (9%), and strains (7%). All represent 94% and mostly include muscles and only 8% had severe injuries include other types like fractures.

Comparison between MRI types of injuries and different age groups showed that out of 47 patients within age group 14 - 45 ys (n=12 patients,25.5 %) had bone injuries , (n=26 patients, 55.3 %) had muscle injuries, (n=6 patients, 12.8%) had joint injuries, (n=2 patients ,

4.3%) of them had joint and muscle injuries and (n=1 patient, 2.1%) of them had bone and muscle injuries the most prevalent lesions were muscle injuries.

Out of 12 patients within age group > 45-65 ys (n=2 patients, 16.7 %) had bone injuries and (n=10 patients , 83.3 %) had muscle injury and also this showed that the most prevalent lesions were muscle injuries.

In our study, we noticed that: Muscle injuries is the most common lesion represent 39 of 59 patients (66.1%) with this lesions and this showed that strain represent 59% of muscular injuries and 39% of all injuries so it was the commonest lesion within athlete group.

This agree with Kumaravel et al. <sup>[7]</sup> who had MRI study on American football player at the high school level and at the collegiate level and found that the large proportion of all American football injuries are muscular. Strains of muscle represent 12–24% of all high school injuries [and approximately 20% of all collegiate practice injuries. Approximately half of all muscle strains occur in the thigh. The hamstrings, hip flexors, and quadriceps were the most frequently injured muscles during the New York Giants training camp, in decreasing order of incidence.

Bone injuries had higher incidence in age group 14-18 with total cases 7cases out of 9 patients in this group (77.8%) and in Final detailed diagnosis of lesions by MRI the most common lesion in bone category was AIIS .

This agree with Yeager et al. <sup>[8]</sup> who had study on Pelvic Avulsion Injuries in the Adolescent Athlete that showed high rate of this type of injuries within old adolescent boys. in review of 228 pediatric pelvic avulsion fractures, It was established that 49% were AIIS avulsions and that the most frequently occurring mechanism was running (39 %).

8 patients had joint injuries found in 59 studied sportive patients most lesions were labrum tear 6 cases represent 10.2% of total cases. This not agree with Narvani et al. <sup>[9]</sup> who had published a report on the prevalence of acetabular labrum tears in athletes suffering from

groin pain and showed that in four out of eighteen athletes (22%) had acetabular labrum tear and this difference in result due to use higher modality MR arthrography that more sensitive than MRI in labrum tear detection.

In our study, MRI finding in 59 investigated patients ; 15 patient with bone lesions show AIIS avulsion in 6 cases 40%, stress neck fracture in 4 case 26.7 %, greater trochanteric fracture in 2 case 13.3 %, ASIS avulsion in 1 case 6.7 % and Thigh splint (Adductor insertion Avulsion) in 2 case 13.3%. as most bone lesion include fracture or avulsion we find that conventional radiograph positive in all cases except 4 cases : 1case showed non- healed AIIS avulsion fracture show non-healed fracture line in T1 weighted image and 3 cases of early signs of stress fracture showed grade 1 to 2 edoema of the bone marrow with hyper intense signal in T2 weighted image and STIR image.

8 patient had joint lesions showed labrum tear in 6 case 75% and FAI in 2 case 25 % all patient has MRI signs and negative conventional radiograph except 1case of FAI which had positive X-ray signs

39 patient had muscular lesions showed Greater trochanter syndrome in 12 cases 30.8%, Muscle contusion in 2 case 5.1%, Snapping hip syndrome in 2case 5.1%, Strain of athletic pubalgia in 12 cases 30.8% and Strain in 11 cases 28.2%. all had negative X-ray except 1case showed soft tissue shadow by X-ray and complete muscular tear with hematoma in MRI.

In our study, out of 59 patient with MRI lesions only 13 patients (22%) had positive sign in conventional radiograph include; bone lesion 11 of 13 (84.6%) case with fractures,1 of 13 (7.7%) case in joint lesions had FAI and 1 of 13 (7.7%) case in muscle lesion had complete tear and hematoma in MRI and only showed soft tissue shadow in X-ray this result confirm low sensitivity of X-ray in different soft tissue injuries.

This agree with Mourad et al. <sup>[10]</sup> who showed that X-ray is rapid technique but It is insensitive to soft tissue changes and has limited sensitivity to bony changes.

The limitations of our study were The limited numbers of athletes from each sport separately as every type is known for different types of injury, also wide range of age groups due to random choice of sport type making each group number small and not very accurate results, in addition to that high cost of MRI.

### **Conclusions:**

In the absence of negative radiographs, MRI of the hip has established itself as the gold standard for diagnosing sports-related hip and groin injuries, also it can determine early signs of stress fracture before it appear in X-ray. so, more details about different lesions can be obtained from MRI and This help in accurate diagnosis, rapid treatment of different athlete lesions and determine time plane of returning to the sport by follow up imaging.

### **COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

### **References:**

1. Suarez JC, Ely EE, Mutnal AB, Figueroa NM, Klika AK, Patel PD, et al. Comprehensive approach to the evaluation of groin pain. *J Am Acad Orthop Surg.* 2013;21:558-70.
2. Kelly BT, Maak TG, Larson CM, Bedi A, Zaltz I. Sports hip injuries: assessment and management. *Instr Course Lect.* 2013;62:515-31.

3. Verbeeten KM, Hermann KL, Hasselqvist M, Lausten GS, Joergensen P, Jensen CM, et al. The advantages of MRI in the detection of occult hip fractures. *Eur Radiol.* 2005;15:165-9.
4. Hegazi TM, Belair JA, McCarthy EJ, Roedl JB, Morrison WB. Sports Injuries about the Hip: What the Radiologist Should Know. *Radiographics.* 2016;36:1717-45.
5. Kerbel YE, Smith CM, Prodromo JP, Nzeogu MI, Mulcahey MK. Epidemiology of Hip and Groin Injuries in Collegiate Athletes in the United States. *Orthop J Sports Med.* 2018;6:2325967118771676.
6. Chan WP, Liu YJ, Huang GS, Lin MF, Huang S, Chang YC, et al. Relationship of idiopathic osteonecrosis of the femoral head to perfusion changes in the proximal femur by dynamic contrast-enhanced MRI. *AJR Am J Roentgenol.* 2011;196:637-43.
7. Kumaravel M, Bawa P, Murai N. Magnetic resonance imaging of muscle injury in elite American football players: Predictors for return to play and performance. *Eur J Radiol.* 2018;108:155-64.
8. Yeager KC, Silva SR, Richter DL. Pelvic Avulsion Injuries in the Adolescent Athlete. *Clin Sports Med.* 2021;40:375-84.
9. Narvani AA, Tsiridis E, Kendall S, Chaudhuri R, Thomas P. A preliminary report on prevalence of acetabular labrum tears in sports patients with groin pain. *Knee Surg Sports Traumatol Arthrosc.* 2003;11:403-8.
10. Mourad C, Omoumi P, Malghem J, Vande Berg BC. Conventional Radiography of the Hip Revisited: Correlation with Advanced Imaging. *Magn Reson Imaging Clin N Am.* 2019;27:661-83.