

PREVALENCE AND ASSOCIATED RISK FACTORS OF ORAL MUCOSAL LESIONS AND OROFACIAL DYSKINESIA IN PATIENTS WITH PSYCHOTIC DISORDERS

ABSTRACT

Aim: To determine the prevalence of oral mucosa lesions, facial movement disorders and the associated risk factors among patients with psychotic disorders and controls.

Study design: A cross sectional study

Place and duration of study: The department of mental health and the General outpatients' clinic of the Obafemi Awolowo University Teaching Hospitals Complex, Ile- Ife, Nigeria. between October 2016 to November 2017.

Methodology: One hundred and eighty consenting patients (90 cases and 90 controls) were recruited. Oral mucosa assessment was based on W.H.O guide to epidemiology and diagnosis of oral mucosa diseases. Involuntary facial movement was assessed using the Abnormal Involuntary Movement Scale. Xerostomia assessment was done by volumetric sialometry.

Results: Prevalence of oral mucosa lesions in patients with psychotic disorders was 83.33% while it was 71.11% for the controls ($P=0.05$). The likelihood of developing oral mucosa lesions in patients with psychotic disorders increased with age (Odds ratio 1.02, 95% C.I (1.00-1.03), $P=0.04$). Orofacial dyskinesia was observed in 17 (18.89%) of the cases on antipsychotic medications and none in the controls ($P<0.001$). The risk of developing orofacial dyskinesia also increased (Odds Ratio 1.096, 95% C.I (1.03-1.16), $P=0.003$) with increase in age in the cases.

Conclusion: Oral mucosa lesions and orofacial dyskinesia were more prevalent in patients with psychotic disorders than the controls. Increasing age was a risk factor for developing oral mucosa lesions and orofacial dyskinesia among patients with psychotic disorders.

Keywords: oral lesions, orofacial dyskinesia, psychotic disorders

INTRODUCTION

Oral health affects the personal, social, and psychological aspects of life[1]. Consequently, a bidirectional relationship exists between mental health and oral health[2]. Individuals with mental disorders are prone to oral health problems. This may be as a result of general self-neglect associated with mental illness, fear of treatment, inability to access dental services and the side effects of medications[3].

Many patients on long term antipsychotic medications are predisposed to xerostomia, leading to increased risk of caries, oral fungal infections, mucosal, lip and tongue lesions, gingivitis and periodontitis[4]. Likewise, some oral health conditions have been associated with mental health challenges. For instance, burning mouth syndrome is a chronic oral dysaesthesia characterized by a burning sensation of the oral cavity with clinically normal mucosa and in the absence of any detectable organic cause[5]. Aetiological factors for burning mouth syndrome include psychological abnormalities, menopause, xerostomia among others[6]. Self-injurious tendencies characterize some mental health disorders. It is a deliberate harm to one's body without suicidal intent. It may occur with a series of psychiatric disorders or in patients with mental retardation, autism, epilepsy[7]. Oral and perioral structures are the tissues on which most self-inflicted lesions are produced[8]. Oral structures are also affected by bruxism: an oral habit consisting of involuntary rhythmic or spasmodic non-functional gnashing, grinding, or clenching of teeth, which may lead to occlusal trauma. The most dangerous form of this pathology is night bruxism, which has a psychoemotional origin[9]. If not treated, it leads to damage of the teeth, periodontium and oral mucosa, pathology of the muscles constituting the masticatory system, headache, cervical pain, and temporomandibular disorders[10]. Psychological and behavioral traits are important in the diagnosis and management of chronic orofacial pain. Furthermore, high levels of depression and somatization have been observed in patients with chronic orofacial pain[11].

Orofacial dyskinesia is an involuntary repetitive movement of the mouth and face[12]. Both orofacial dyskinesia and appendicular dyskinesia are considered sub syndrome of tardive dyskinesia. In most cases, orofacial dyskinesia occurs as a result of long-term exposure to antipsychotic drugs of the phenothiazine and butyrophenone groups. Antipsychotics are classified into typical/first generation and atypical/second generation[13]. The mode of action of the first generation is predominantly to act as antagonists at the brain dopamine D₂ receptors[13]. Some associated adverse drug reaction includes extrapyramidal motor effects (acute dystonia, akathisia, parkinsonism, tardive dyskinesia). The mode of action of second generation is predominant antagonism of 5-HT_{2A} receptors with a lesser degree antagonism of dopamine D₂ receptors. Second generation can be differentiated from the first by their low or negligible side effects, effectiveness and in general supposed increased safety[13].

Long term effects of orofacial dyskinesia include myalgia of the muscles of mastication and temporomandibular disorders as a result of the relentless movement typical of this disorder[14].

Despite the 20% prevalence of mental disorders in the Nigerian population[15], there is paucity of data relating to the risk factors for oral lesions and orofacial dyskinesia in patients living with mental health challenges. Likewise, the prevalence of oral lesions and orofacial dyskinesia in patients receiving psychiatric care in Nigeria is not fully known. Several oral

conditions affect the quality of life of patients with mental disorders. Hence a need for a study describing oral mucosal lesions in a subset this group of patients. The attributes and exposure that increases the likelihood of developing such lesions are also worth exploring. This is with a view to ensuring a holistic patient management. Furthermore, there is need to bring these findings to the fore in order to encourage a more robust multidisciplinary approach in the management of patients with mental disorders.

Accordingly, this study was designed to investigate the prevalence of oral mucosa lesions, facial movement disorders in our environment and the associated risk factors among patients diagnosed with psychotic disorders and controls.

MATERIAL AND METHODS

Ethics: The procedures for this study were in accordance with the ethical standards of the Helsinki Declaration of 1975, as revised in 2000. Ethical approval for this study was obtained from the ethics and research committee, Obafemi Awolowo University Teaching Hospitals Complex, Ile-Ife with registration Number: NHREC/27/02/2009a.

Patients

A cross sectional study investigating oral lesions, orofacial dyskinesia and associated risk factors in patients with psychotic disorders attending the mental health clinic of the Obafemi Awolowo University Teaching Hospitals Complex, Ile-Ife, Nigeria and controls from the General outpatients' clinic between October 2016 to November 2017.

The study population comprised of outpatients diagnosed with psychotic disorders. They were randomly selected from the patients being managed in the mental health clinic. Patients were 18 years and above with no co-morbid medical conditions like primary hypertension and diabetes mellitus. The control group was recruited to compare with patients with psychotic disorders. They comprised healthy individuals who visited the general outpatients' department of the hospital on account of pre-employment, pre-admission and routine medical checkup. Other inclusion criteria for control subjects were negative history of diagnosed mental disorder, use of antipsychotic medications and co morbid medical conditions.

Sample size determination

The formula for sample size estimation of two proportions, which gave the total number of sample size required for the two groups, was used[16]. With a statistical significance (Alpha) of 5% and power of 90%, a known prevalence of dyskinesia within the general population was 1.3%[17] and was allotted to population 1 (P1). Also the known prevalence of dyskinesia in patients with mental disorders was 15%[18] and was allotted to population 2 (The comparison group, P2). The effect size (D) of 13.7% (P2-P1) difference was adopted for this study. Attrition rate of 10%

(rounded off to 16 patients) was added into the calculated sample size giving a total of 180. Since there are two groups, this translated to 90 patients per group.

Data collection

Patients who met the inclusion criteria were informed about the study after which a signed consent was obtained. Biodata such as name, age, sex, address, ethnicity, marital status and occupation were recorded. Information on the patients' past medical history and drug history of note as well as habits that may result in similar oral conditions such as tobacco and alcohol consumption were also recorded in addition to blood pressure and weight. Other relevant information such as established diagnosis of mental disorders and facial movement disorder and the present medication for each participant were obtained from the hospital record.

Detailed extra oral and intraoral examination was done. Oral mucosa assessment was based on the W.H.O guide to epidemiology and diagnosis of oral mucosal diseases[19].

Involuntary facial movement

Involuntary facial movement was assessed using the Abnormal Involuntary Movement Scale (AIMS)[20]. The scale aids in early detection of dyskinesia and also provides a method for ongoing surveillance. Patients were first observed unobtrusively at rest in the waiting area. Patients were seated on a firm chair without arms for questioning and examination following standard operations as outlined in the scale. Observed abnormal movements of the face and mouth, extremities, trunk and global judgements were rated according to severity on the AIMS scale (0=none, 1=minimal, 2=mild, 3=moderate, 4=severe) by one examiner.

Oral hygiene: Oral hygiene was assessed using Greene and Vermilion's Oral Hygiene Index Simplified [21]. Six teeth were examined one from each sextant (upper anterior right central incisor, upper right first molar, upper left first molar, lower left central incisor, lower right first molar and lower left first molar) of the jaw and the corresponding scores were allotted.

The addition of the total score was divided by total number of teeth examined (Six). This was done for each of debris and calculus index respectively. Summation of both debris and calculus index gave the oral hygiene index and was graded as follows: 0-1.2 = good; 1.3-3.0 = fair; 3.1-6.0= poor.

Xerostomia: Assessment of hyposalivation was done by sialometry, measuring resting unstimulated whole saliva using standardized spitting method[22]. Patients were asked to abstain from eating and drinking thirty minutes before saliva collection. Patients were asked to spit into a sputum jar for a period of five minutes. Measurement of the volume of collected saliva was done using a graduated Pasteur pipette. The volume was divided by five minutes to get the flow rate in milliliters/minute (mls/min). Flow rate of 0.3-0.5mls/min was adjudged normal while rate less than or equal to 0.1mls/min was adjudged hyposalivation[23].

Examiner calibration

An initial calibration of the researcher was done by both the supervisor (a consultant mental health physician) and the researcher. The inter-examiner consistency was determined by calculating the Kappa (k) statistics [24]. The Kappa coefficient calculated for this study was 0.91 for oral mucosal lesions and 0.87 for orofacial dyskinesia.

Mental health status of the comparison group

Mental health status of the control group was assessed using the General Health Questionnaire 12, GHQ-12 [25]. It has been validated for use in Nigeria [26]. Prospective patients with scores greater than 15, which is indicative of distress, were not included in the study.

Statistical analysis

Data analysis was done using Stata 13 statistical software (Statacorp, College Station, Texas). Descriptive statistics were used to characterize socio-demographic variables such as age, sex, marital status, and occupation. Bivariate analysis such as Chi-square or Fisher's exact test was used as appropriate to compare categorical variables.

The association between the predictors and the outcome was determined using odds ratios for patients with mental disorders with and without the outcome and for the controls and patients with mental disorders. All the risk factors were fitted into a logistic regression model to see how each factor contributed to each of oral lesions and orofacial dyskinesia and the interactions between the factors. Logistic regression was used to determine the role of the predictor and confounders on the primary outcome.

Statistical significance was set at $p < 0.05$ and confidence interval was set at 95% for all the analysis.

RESULTS

Distribution of patients

One hundred and eighty patients participated in the study, 93 (51.67%) females and 87 (48.33%) males. The controls comprised 49 (54.44%) males and 41 (45.56%) females with mean age of 33.30 (± 12.61) years. Overall mean age of patients was 36.68 \pm 12.45 years (Table 1).

Table 1. Demographic and Clinical characteristics of patients

CHARACTERISTIC	TOTAL (N=180)	CASES (N=90)	CONTROLS (N=90)	P VALUE
Age (mean \pm SD, years)	36.68(\pm 12.45)	40.06(\pm 11.40)	33.30(\pm 12.61)	
Sex (n%)				
Male	87(48.33)	38 (42.44)	49 (54.44)	
Female	93(51.67)	52(57.78)	41(45.56)	
Oral hygiene index simplified (n%)				
Good	69(38.33)	12(13.33)	57(63.33)	
Fair	46(25.56)	33(36.67)	13(14.44)	
Poor	65(36.11)	45(50.00)	20(22.22)	<0.001*
Substance Abuse (n%)				
No	177(98.33)	87(96.67)	90(100.00)	
Yes	3(1.67)	3(3.33)	0(0.00)	0.25
Alcohol Intake (n%)				
No	175(97.22)	84(93.33)	75(83.33)	
Yes	21(11.67)	6(6.67)	15(16.67)	0.04*
Cigarette Smoking (n%)				
No	175(97.22)	85(94.44)	90(100.00)	
Yes	5(2.78)	5(5.56)	0(0.00)	0.06
Oral lesions (n%)				
Absent	41(22.78)	15(16.67)	26(28.89)	
Present	139(77.22)	75(83.33)	64(71.11)	0.05
Orofacial Dyskinesia (n%)				
Absent	163(90.56)	73(81.11)	90(100.00)	

Present	17(9.44)	17(18.89)	0(0.00)	<0.001*
Hyposalivation(n%)				
Absent	170(94.44)	81(90.00)	89(98.89)	
Present	10(5.56)	9(10.00)	1(1.11)	0.02*

*Statistically significant value ($P<0.05$)

Clinical Characteristics of patients

Poor oral hygiene was observed in 45(50.00%) of the cases and 20(22.22%) of the controls. Oral lesions were present in 75(83.33%) of the cases and 64(71.11%) of the controls. Orofacial dyskinesia was observed in 17(18.89%) of the cases while none of the controls had facial movement disorder. Hyposalivation was present in 9(10.00%) of the cases and 1(1.11%) of the controls (Table 1).

The most frequent psychotic disorder was schizophrenia 58(64.44). First generation antipsychotic was used in the management of most of the cases, 77(85.56%). Forty-seven cases (52.00%) were on antimuscarinics (Table 2).

Table 2: Clinical diagnosis and findings in the management of cases

CHARACTERISTIC	n (%)
MENTAL DISORDER	
Depression	7 (7.78)
Multiple Psychoactive Substance Use Disorder	1(1.11)
Psychosis	15 (16.67)
Schizophrenia	58 (64.44)
Unspecified Psychotic disorder	4(4.44)
Bipolar Affective Disorder	5(5.56)
ANTIPSYCHOTICS	
First generation	77(85.56)
Second generation	13(14.44)
ON ANTIMUSCARINICS	
Yes	47(52.00)
No	43(48.00)
DURATION OF MENTAL DISORDER	
6 months	7(7.78)

1-2 years	18(20.00)
3-5 years	65(72.22)

Linear alba buccalis was the most frequent oral mucosa lesion seen in patients with psychotic disorder, 55 out of 90 (61.11%) and controls 60(66.67%). Other oral lesions include fissured tongue, frictional keratosis and geographic tongue (Table 3).

Table 3: Oral lesions and dyskinesia in patients with mental disorders and controls

ORAL LESIONS	CASE(N=90)	CONTROLS(N=90)	P VALUE
FISSURED TONGUE (n%)			
Absent	71(78.89)	87(96.67)	
Present	19(21.11)	3(3.33)	<0.001*
FRictionAL KERATOSIS (n%)			
Absent	80(88.89)	87(96.67)	
Present	10(11.11)	3(3.33)	0.08
LINEA ALBA (n%)			
Absent	35(38.89)	30(33.33)	
Present	55(61.11)	60(66.67)	0.44
GEOGRAPHIC TONGUE (n%)			
Absent	87(96.67)	90(100.00)	
Present	3(3.33)	0(0.00)	0.25
OROFACIAL DYSKINESIA (n%)			
Absent	73(81.11)	90(100.00)	
Present	17(18.89)	0(0.00)	<0.001*

*Statistically significant value ($P<0.05$)

Orofacial dyskinesia in patients

Table 3 shows orofacial dyskinesia in patients with mental disorders. Orofacial dyskinesia was observed in 17(18.89%) of the cases and none in the controls. The difference was statistically significant with ($p < 0.001$).

Risk factor for oral lesions and orofacial dyskinesia in psychotic disorders

Table 4 shows that age is associated with developing oral lesions in mental disorders. This was adjusted for sex, alcohol intake and antipsychotic medication. The likelihood of developing oral lesions increased with every one-year increase in age in patients with psychotic disorders (Odds ratio 1.02, 95% C.I (1.00-1.03), $p = 0.04$). Sex, alcohol intake and the type of antipsychotic medication did not have a significant relationship with the likelihood of developing oral lesions.

Another regression model showed that age is associated with developing orofacial dyskinesia in mental disorders (Table 4). This was adjusted for sex and type of antipsychotic medication. The risk of developing orofacial dyskinesia increased with age in patients with psychotic disorders (Odds Ratio 1.096, 95% C.I (1.03-1.16), $p = 0.003$).

Table 4: Risk factors for oral lesions and orofacial dyskinesia in psychotic disorders

RISK FACATOR	ODDS RATIO	95% CONFIDENCE INTERVAL	P VALUE
For Oral Lesions			
Age (Years)	1.02	1.00-1.03	0.04*
Males (Reference Females)	1.81	0.87-3.77	0.11
Alcohol Intake(yes)	0.38	0.12-1.28	0.12
Antipsychotic:1 st generation(Reference 2 nd generation)	1.22	0.42-3.54	0.72
For Orofacial dyskinesia			
Age (years)	1.096	1.03-1.16	0.003*
Male (Reference Female)	1.71	0.53-5.55	0.37
Antipsychotic type:1 st generation(Reference	0.14	0.02-1.81	0.07

2nd generation)

*Statistically significant value (P<0.05)

Antipsychotics and orofacial dyskinesia

Table 5 shows that the type of antipsychotics is a risk factor for development of orofacial dyskinesia in patients with mental disorders. Orofacial dyskinesia was observed in 15(19.48%) cases on first generation antipsychotics while two (15.38%) cases on second generation antipsychotics had orofacial dyskinesia. The observed difference was statistically significant with p<0.001.

Table 5: Antipsychotics as a risk factor for orofacial dyskinesia

ANTIPSYCHOTIC	OROFACIAL DYSKINESIA	OROFACIAL	P VALUE
	ABSENT (N%)	DYSKINESIA PRESENT (N%)	
First Generation	62(80.52)	15(19.48)	
Second Generation	11(84.62)	2(15.38)	
Not on antipsychotic	90(100.00)	0(0.00)	
Total	163(90.56)	17(9.44)	<0.001*

Fisher's exact, *statistically significant

DISCUSSION

Oral lesions with psychosomatic aetiology are still not a sufficiently explained subgroup of diseases, which have long been known in medicine[27]. In addition, there are insufficient explanations for the aetiology of many oral changes, which are believed to have psychosomatic basis[27].

Some studies from South Asia have previously documented the presence of oral lesions in patients with mental disorders. We observed a much higher prevalence than the 39.3% reported by Arjun and colleagues[28]. Furthermore, there is a marked difference in the specific oral lesions reported in their study. While linea alba buccalis and fissured tongue had the highest proportion in this study, leukoplakia and oral submucous fibrosis were the most prevalent oral lesions observed by Arjun and colleagues. This can be attributed to high tobacco and areca nut consumption among the patients with

mental disorders in their study population. In contrast, prevalence of smoking in patients with psychotic disorders in our study was low, 5.56%.

Linea alba is a relatively common asymptomatic alteration of the buccal mucosa[29]. This may explain the high proportion in both the cases and the control in this study. Fissured tongue is characterized by presence of grooves or fissures on the dorsum or lateral surface of the tongue[30]. The etiology remains unclear however, associations with genetics and systemic diseases such as psoriasis[31], diabetes[32], orofacial granulomatosis[33] have been reported. In males, fissured tongue and geographic tongue may be a significant predictor for burning mouth sensation[34]. Geographic tongue is characterized by loss of filiform papillae migrating in a map-like pattern[35]. While the etiology is unknown, risk factor associated with mental health such as stress has been reported and the disorder may be accompanied by anxiety[36]. It is usually asymptomatic and often coexists with fissured tongue. When symptoms occur, it presents as painful burning sensation associated with smoking, spicy foods and fruits which can be a source of distress to the affected individual.

Dangore and colleagues[37] also reported a lower prevalence of 45.3% for oral lesions among individuals with mental disorders. Recurrent aphthous stomatitis, burning mouth syndrome and oral lichen planus were observed more commonly in their study. Suresh and colleagues observed a similar prevalence of 45% for oral lesions in patients with mental disorders[38]. Differences in sociocultural and environmental factors may have been responsible for the markedly different observation in the specific oral lesions in our study compared with other similar studies. The other studies were carried out among an Indian population, with a high prevalence of tobacco use in different forms, chewing of areca nuts and other agents.

Suresh and colleagues[38] and Arjun and colleagues[28] reported a prevalence of 5.17% and 30.30% respectively for oral lesions in patients who have not been diagnosed of any mental disorders in studies that compared oral lesions in individuals with mental disorders and controls. These values are however at variance with the recorded prevalence in this study. Marked aforementioned differences were also observed in specific oral lesions like in the mental health population and similar sociocultural and environmental factors may be responsible. A few risk factors have been identified for oral lesions in individuals with mental disorders. These risk factors are largely life style and habit related. They include tobacco smoking, alcohol intake among others[28]. This study showed an association between increase in age and development of oral lesions in individuals with psychotic disorders compared to the controls.

A Nigerian study reported a prevalence of 27% for tardive dyskinesia in patients with mental health challenges[39]. The higher prevalence may be as a result of the study considering both orofacial and appendicular dyskinesia as a sub syndrome of tardive dyskinesia. Furthermore, a systematic review by Correll and colleagues reported 32.4% and 13.1% prevalence for adults on first generation and second generation antipsychotic respectively [40]. It is noteworthy that most of the literature on dyskinesia did not consider orofacial dyskinesia in isolation but as a sub syndrome of tardive dyskinesia. In addition, a common denominator observed in patients in this study and other literature that reported tardive dyskinesia is that drugs (antipsychotics and neuroleptics) are a common risk factor[41]. Furthermore, the association between the generations of antipsychotics and orofacial dyskinesia observed in this study is consistent with findings in the literature [40].

Dyskinesia is still a major concern in psychiatry. It is most often a side effect of psychotropic drugs, especially the first generation antipsychotics[42]. Although the prevalence is decreasing with the advent of second-generation antipsychotics[43], most psychiatric care receivers in resource limited environment like Nigeria are still treated with the first generation psychotropic drugs because the second generation drugs are more expensive. In addition, the limited coverage of health insurance compels many patients to pay out of pocket and find first generation drugs are more affordable. Similar to first generation antipsychotics, the atypical antipsychotics have some degree of D2 antagonism but more of blockade of 5HT2A receptors[44]. Loose binding to and fast dissociation from D2 receptors is suggested to be the cause of their lower propensity for dyskinesia[45]. All the patients in this study with orofacial dyskinesia did not experience any existing movement disorder prior to commencement of psychotropic medications. Dental problems such as attrition and abfractions are very frequent in dyskinesia[46], as are pain from myalgia, temporomandibular disorders, traumatic lesions, tooth wear and impaired retention of prosthetic devices[47].

This study showed an association between increasing age and the development of orofacial dyskinesia in individuals with mental disorders. This is consistent with findings in other studies wherein ageing was observed as a critical factor for developing dyskinesia[48-51]. The lack of significant gender effect is consistent with previous investigations. However, other risk factors such as female sex, duration of use of antipsychotics, reported by Tarsy and colleagues[52] were not observed in this study. Cognitive impairment, chronicity of psychotropic,[50] diabetes mellitus[53], smoking[54], alcohol [55] and African-American ethnicity[56] are other risk factors. The explanations that have been suggested that associate age with dyskinesia are increased incidence of medical illness with advancing age and neurochemical changes[57-59]. Matson and colleagues suggested decrease in drug metabolism capacity with advancing age as a risk factor for dyskinesia[50].

Although mental health care givers may not be able to diagnose oral lesions, their presence in this group of patients calls for a good understanding of the interaction between oral health and mental health. The lack of coordination between oral physicians and psychiatrists is evident by the few referrals between the two specialists in the center in which this study was carried out. Although the oral lesions observed at the time this study was carried out were mostly not symptomatic, possible long-term systemic association of lesions such as fissured and geographic tongue may require patient follow up.

CONCLUSION

The prevalence of oral lesions present in patients with psychotic disorders was higher than in the controls from the same base. Fissured tongue, frictional keratosis, linea alba and geographic tongue were the oral mucosal lesions seen in both cases and control. However, only the presence of fissured tongue was statistically significant in both groups.

Increasing age was a risk factor for developing oral lesions and orofacial dyskinesia in patients with psychotic disorders. Treatments with first generation antipsychotics were more associated with orofacial dyskinesia than second-generation antipsychotics.

To further establish causality between psychotic disorders and oral lesions, an experimental study is recommended. Moreover, interprofessional collaboration between mental and oral health physicians can be strengthened by the provision of an avenue for more knowledge, insights and resource sharing. This partnership will enhance understanding of professional roles with a view to a more holistic approach in patient management.

LIMITATIONS OF THE STUDY

This study did not quantify the extent to which oral lesions and orofacial dyskinesia exacerbates the mental disorders of the patients. Being a cross sectional study, it cannot lend to discussions on causation.

Consent: All authors declare that written informed consent was obtained from all the patients that participated in this study. A copy of the form is available to the editorial board of this journal.

Ethical Approval: Ethical approval for this study was obtained from the ethics and research committee, Obafemi Awolowo University Teaching Hospitals Complex, Ile-Ife with registration Number: NHREC/27/02/2009a.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

1. Mirza I, Day R, Wulff-Cochrane V, Phelan M (2001) Oral health of psychiatric in-patients. *The Psychiatrist* 25 (4):143-145
2. Sims A (1987) Why the excess mortality from psychiatric illness? *Br Med J (Clin Res Ed)* 294 (6578):986-987
3. Cormac I, Jenkins P (1999) Understanding the importance of oral health in psychiatric patients. *Advances in psychiatric treatment* 5:53-60
4. Stiefel DJ, Truelove EL, Menard TW, Anderson VK, Doyle PE, Mandel LS (1990) A comparison of the oral health of persons with and without chronic mental illness in community settings. *Spec Care Dentist* 10 (1):6-12
5. Barker K, Savage N (2005) Burning mouth syndrome: an update on recent findings. *Aust Dent J* 50 (4):220-223
6. Trombelli L, Zangari F, Calura G (1994) [The psychological aspects of patients with the burning mouth syndrome]. *Minerva Stomatol* 43 (5):215-221
7. Sæmundsson SR, Roberts MW (1996) Oral self-injurious behavior in the developmentally disabled: review and a case. *ASDC J Dent Child* 64 (3):205-209, 228
8. Silva DR, Da Fonseca M (2003) Self-injurious behavior as a challenge for the dental practice: a case report. *Pediatr Dent* 25 (1):62-66
9. De Laat A, Macaluso GM (2002) Sleep bruxism as a motor disorder. *Mov Disord* 17 (S2):S67-S69
10. Manfredini D, Landi N, Fantoni F, Segu M, Bosco M (2005) Anxiety symptoms in clinically diagnosed bruxers. *J Oral Rehabil* 32 (8):584-588
11. Yap AU, Tan KB, Prosthodont C, Chua EK, Tan HH (2002) Depression and somatization in patients with temporomandibular disorders. *The Journal of prosthetic dentistry* 88 (5):479-484
12. Kobayashi RM (1976) Orofacial dyskinesia: Clinical features, mechanisms and drug therapy. *West J Med* 125 (4):277
13. Pakpoor J, Agius M (2014) A review of the adverse side effects associated with antipsychotics as related to their efficacy. *Psychiatr Danub* 26 (Suppl 1):273-284
14. Pekkan G, Kilicoglu A, Algin DI (2010) Treatment of a tardive dyskinesia patient with temporomandibular disorder: a case report. *J Orofac Pain* 24 (2):212
15. Samba D (2012) Gambia Mental Health Report 2012. Ibadan, Nigeria, University College Hospital, Department of Psychiatry. Mental Health Leadership and Advocacy Programme (mhLAP)
16. Feinstein A (2002) Principles of medical statistics. Boca Raton, Fla. CRC:503
17. Woerner MG, Kane JM, Lieberman JA, Alvir J, Bergmann KJ, Borenstein M, Schooler NR, Mukherjee S, Rotrosen J, Rubinstein M, et al. (1991) The prevalence of tardive dyskinesia. *J Clin Psychopharmacol* 11 (1):34-42
18. Gerlach J, Casey DE (1988) Tardive dyskinesia. *Acta Psychiatr Scand* 77 (4):369-378
19. Organization WH, Staff WHO (1995) Application of the International Classification of Diseases to Dentistry and Stomatology: ICD-DA. World Health Organization,

20. Guy W (1976) Abnormal involuntary movement scale (AIMS). ECDEU assessment manual for psychopharmacology 338:534-537
21. Greene JG, Vermillion JR (1964) The simplified oral hygiene index. The Journal of the American Dental Association 68 (1):7-13
22. Navazesh M, Kumar SK (2008) Measuring salivary flow: challenges and opportunities. The Journal of the American Dental Association 139:35S-40S
23. Sreebny L (1992) Saliva: Its role in health and disease. Working group 10 of the commission on oral health, research and epidemiology (CORE). Int Dent J 42:287-304
24. Sim J, Wright CC (2005) The kappa statistic in reliability studies: use, interpretation, and sample size requirements. Phys Ther 85 (3):257-268
25. Goldberg D (1978) Manual of the general health questionnaire. NFER Nelson,
26. Gureje O, Obikoya B (1990) The GHQ-12 as a screening tool in a primary care setting. Social psychiatry and psychiatric epidemiology 25 (5):276-280
27. Kumar NN, Panchaksharappa MG, Annigeri RG (2016) Psychosomatic disorders: An overview for oral physician. Journal of Indian Academy of Oral Medicine and Radiology 28 (1):24
28. Arjun TN, Sudhir H, Sahu R, Saxena V, Saxena E, Jain S (2014) Assessment of oral mucosal lesions among psychiatric inmates residing in central jail, Bhopal, Madhya Pradesh, India: A cross-sectional survey. Indian J Psychiatry 56 (3):265
29. Bhattacharyya I, Chehal HK (2011) White lesions. Otolaryngol Clin North Am 44 (1):109-131
30. Dafar A, Çevik-Aras H, Robledo-Sierra J, Mattsson U, Jontell M (2016) Factors associated with geographic tongue and fissured tongue. Acta Odontol Scand 74 (3):210-216
31. Dawson T (1974) Tongue lesions in generalized pustular psoriasis. Br J Dermatol 91 (4):419-424
32. Guggenheimer J, Moore PA, Rossie K, Myers D, Mongelluzzo MB, Block HM, Weyant R, Orchard T (2000) Insulin-dependent diabetes mellitus and oral soft tissue pathologies: I. Prevalence and characteristics of non-candidal lesions. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology 89 (5):563-569
33. Marcoval J, Viñas M, Bordas X, Jucglà A, Servitje O (2012) Orofacial granulomatosis: clinical study of 20 patients. Oral surgery, oral medicine, oral pathology and oral radiology 113 (4):e12-e17
34. Ching V, Grushka M, Darling M, Su N (2012) Increased prevalence of geographic tongue in burning mouth complaints: a retrospective study. Oral surgery, oral medicine, oral pathology and oral radiology 114 (4):444-448
35. Assimakopoulos D, Patrikakos G, Fotika C, Elisaf M (2002) Benign migratory glossitis or geographic tongue: an enigmatic oral lesion. The American journal of medicine 113 (9):751-755
36. Alikhani M, Khalighinejad N, Ghalaiani P, Khaleghi MA, Askari E, Gorsky M (2014) Immunologic and psychologic parameters associated with geographic tongue. Oral surgery, oral medicine, oral pathology and oral radiology 118 (1):68-71
37. Dangore-Khasbage S, Khairkar PH, Degwekar SS, Bhowate RR, Bhake AS, Singh A, Lohe VK (2012) Prevalence of oral mucosal disorders in institutionalized and non-institutionalized psychiatric patients: a study from AVBR Hospital in central India. J Oral Sci 54 (1):85-91
38. Suresh K, Ganiger CC, Ahammed YA, Kumar MC, Pramod R, Nayak AG, Vibhute N (2014) Psychosocial characteristics of oromucosal diseases in psychiatric patients: Observational study from Indian dental college. N Am J Med Sci 6 (11):570
39. Gureje O (1989) The significance of subtyping tardive dyskinesia: a study of prevalence and associated factors. Psychol Med 19 (1):121-128

40. Correll CU, Leucht S, Kane JM (2004) Lower risk for tardive dyskinesia associated with second-generation antipsychotics: a systematic review of 1-year studies. *Am J Psychiatry* 161 (3):414-425
41. Dolder CR, Jeste DV (2003) Incidence of tardive dyskinesia with typical versus atypical antipsychotics in very high risk patients. *Biol Psychiatry* 53 (12):1142-1145
42. Woerner MG, Alvir JM, Saltz BL, Lieberman JA, Kane JM (1998) Prospective study of tardive dyskinesia in the elderly: rates and risk factors. *Am J Psychiatry* 155 (11):1521-1528
43. Leucht S, Pitschel-Walz G, Abraham D, Kissling W (1999) Efficacy and extrapyramidal side-effects of the new antipsychotics olanzapine, quetiapine, risperidone, and sertindole compared to conventional antipsychotics and placebo. A meta-analysis of randomized controlled trials. *Schizophr Res* 35 (1):51-68
44. Divac N, Prostran M, Jakovcevski I, Cerovac N (2014) Second-generation antipsychotics and extrapyramidal adverse effects. *BioMed research international* 2014
45. Kapur S, Seeman P (2001) Does fast dissociation from the dopamine D2 receptor explain the action of atypical antipsychotics?: A new hypothesis. *Am J Psychiatry* 158 (3):360-369
46. Lumetti S, Ghiacci G, Macaluso G, Amore M, Galli C, Calciolari E, Manfredi E (2016) Tardive dyskinesia, oral parafunction, and implant-supported rehabilitation. *Case reports in dentistry* 2016
47. Myers DE, Schooler NR, Zullo TG, Levin H (1993) A retrospective study of the effects of edentulism on the severity rating of tardive dyskinesia. *The Journal of prosthetic dentistry* 69 (6):578-581
48. Jeste DV, Caligiuri MP, Paulsen JS, Heaton RK, Lacro JP, Harris MJ, Bailey A, Fell RL, McAdams LA (1995) Risk of tardive dyskinesia in older patients: a prospective longitudinal study of 266 outpatients. *Arch Gen Psychiatry* 52 (9):756-765
49. Tenback D, Bakker P (2015) Risk factors for tardive movement disorders in schizophrenia.
50. Matson JL, Fodstad JC, Neal D, Dempsey T, Rivet TT (2010) Risk factors for tardive dyskinesia in adults with intellectual disability, comorbid psychopathology, and long-term psychotropic use. *Res Dev Disabil* 31 (1):108-116
51. Wszola BA, Newell KM, Sprague RL (2001) Risk factors for tardive dyskinesia in a large population of youths and adults. *Exp Clin Psychopharmacol* 9 (3):285
52. Tarsy D, Baldessarini RJ (2006) Epidemiology of tardive dyskinesia: is risk declining with modern antipsychotics? *Mov Disord* 21 (5):589-598
53. Ganzini L, Heintz RT, Hoffman WF, Casey DE (1991) The prevalence of tardive dyskinesia in neuroleptic-treated diabetics: a controlled study. *Arch Gen Psychiatry* 48 (3):259-263
54. Menza MA, Grossman N, Van Horn M, Cody R, Forman N (1991) Smoking and movement disorders in psychiatric patients. *Biol Psychiatry* 30 (2):109-115
55. Dixon L, Weiden PJ, Haas G, Sweeney J, Frances AJ (1992) Increased tardive dyskinesia in alcohol-abusing schizophrenic patients. *Compr Psychiatry* 33 (2):121-122
56. Sramek J, Roy S, Ahrens T, Pinanong P, Cutler NR, Pi E (1991) Prevalence of tardive dyskinesia among three ethnic groups of chronic psychiatric patients. *Psychiatr Serv* 42 (6):590-592
57. Aman MG (1990) Considerations in the use of psychotropic drugs in elderly mentally retarded persons. *J Intellect Disabil Res* 34 (1):1-10
58. Lohr JB, Bracha HS (1988) Association of psychosis and movement disorders in the elderly. *Psychiatr Clin North Am*
59. Kane J, Woerner M, Lieberman J (1985) Tardive dyskinesia: prevalence, incidence, and risk factors. In: *Dyskinesia*. Springer, pp 72-78