

Ecophysiology and vegetative and productive behavior of 'Chardonnay' vines under protected cultivation systems in Serra Catarinense

ABSTRACT

Aims: This study aimed to evaluate the effect of protected cultivation, with anti-hail screen or plastic cover, on phenological behavior, water potential, gas exchange, vegetative growth and vine production under protected cultivation systems in an altitude region of Santa Catarina.

Place and Duration of Study: The experiment was carried out in São Joaquim, SC, in the 2018/19, 2019/20 and 2020/21 harvests, with the cultivar Chardonnay on Paulsen 1103 rootstock.

Methodology: The treatments consist of an uncovered environment (control) and crops protected with white screen anti-hail (4 mm x 7 mm) and clear plastic raffia cover (160 µm).

Results: The anti-hail net did not interfere in the phenological stages, photosynthesis and gas exchange of the plants and reduced the RFA, vegetative growth and production, in relation to the plastic cover, and similar development conditions to the control treatment (without cover). The plastic cover accelerated the maturation process, reduced photosynthetically active radiation (RFA), promoted vegetative growth, delayed leaf senescence and increased productivity in relation to plants without cover, but did not change the photosynthesis of plants and other attributes related to the plants. gas exchange. The highest red to far red ratio was observed in an uncovered system. The water potential of the plants was not affected. Net photosynthesis, gas exchange, CO₂ ratio did not show significant differences. The light saturation point was higher in plants in the cultivation system protected with plastic cover. Chlorophyll a, b, a+b and the chlorophyll a/b ratio were higher in plants under plastic cover in the 2018/19 harvest. For the years 2019/20 and 2020/21, these variables of net photosynthesis, gas exchange and CO₂ ratio did not show differences between treatments, with the exception of the 2020/21 harvest, in which the plastic cover showed a higher value for chlorophylls a + b. Plants under plastic cover showed an anticipation of fruit maturation in relation to the screen and without cover. Plants under plastic cover showed higher values than plants without cover in terms of bunch weight (116%), number of bunches per plant (49%) and weight of bunches per plant (65%). The cultivation protected with anti-hail net maintained these attributes similar to the cultivation without cover.

Keywords: *Altitude wines.; microclimate; wine tourism; hail net; plastic cover.*

1. INTRODUCTION

Although Brazil does not have a tradition in wine production, as with many European countries, there is a considerable area dedicated to the cultivation of grapes and wine production, such as the temperate production zones of Campanha in Rio Grande do Sul, Serra Gaúcha and the Serra Catarinense [1].

The wines produced in the Serra Catarinense region (above 800 m altitude) are highlighted in the national scenario [2]. The factors that make the cultivation of high altitude wine grapes interesting, culturally and economically, are associated with the differentiated flavor and

22 aroma, which, combined with wine tourism, make the culture increasingly attractive to the
23 consumer and profitable to the producer [3].

24 However, places with an altitude of more than 700 m are more prone to hail and late frosts
25 [4], in addition to high rainfall at the time of sprouting and harvest [5], high relative humidity
26 and excessive leaf wetness, conditions that require application of chemical products to
27 control diseases, a factor that affects the quality of wines [6]. Another limiting factor is the
28 anticipation of harvest in many vintages, not allowing the grapes to reach the appropriate
29 ripening period, which affects the productivity and oenological attributes of the wines
30 produced [6, 7].

31 Considering the aspects of climate and grape production in regions of the Serra
32 Catarinense, many producers are adopting cultivation in a protected environment with the
33 use of anti-hail screen [8] or plastic cover [9], aimed at protecting plants from hail and late
34 frosts [10]. However, the impacts of protected cultivation on plant physiology and fruit quality
35 in altitude regions for grape cultivation are still scarce in the literature. The implementation of
36 plastic cover and anti-hail mesh in the Serra Catarinense region requires studies, mainly on
37 factors related to physiological and microclimatic responses, vegetative and productive
38 behavior [11].

39 The use of plastic cover increases the temperature inside the canopy and reduces humidity,
40 evapotranspiration and incident solar radiation [12, 13, 14]. In regions of Rio Grande do Sul,
41 which have a tradition of protected cultivation and the production of quality wines, the works
42 are more advanced regarding the interference of the use of plastic covering. However, in the
43 Serra Catarinense region, little is known about the effects of using this technology.

44 The objective of this work was to evaluate the effects of microclimatic conditions, phenology,
45 gas exchange, light interception, water potential in plants, vegetative and productive
46 behavior of vines managed in an uncovered management system, under cover with an anti-
47 hail screen and transparent plastic tarpaulin in altitude region in the Serra Catarinense.

48

49 **2. MATERIAL AND METHODS**

50

51 The experiment was carried out in the 2018/19, 2019/20 and 2020/21 cycles with the cultivar
52 'Chardonnay', at the Monte Agudo Winery, located in São Joaquim, SC, at 1,264 m altitude,
53 with geographic coordinates 28°14'54' 'S and 49°47'52"W. The vineyard has been implanted
54 for 11 years (2007 to 2018), on Paulsen 1103 rootstock. The conduction system is "Y", with
55 a spacing of 2.90 m between rows and 1.30 m between plants, obtaining a density of 2,870
56 plants per hectare. Pruning is mixed with four sticks per plant and 4 to 6 buds per stick.

57 The climate classification of the region of São Joaquim, SC, is Cfb, according to Köppen
58 [15], with moderate temperature and well-distributed rain, mild summer, occurrence of frosts
59 in winter and autumn, with average temperatures below 20° C, except in summer, when
60 temperatures exceed 28°C. In winter, the average temperature is below 14°C, with minimum
61 temperatures below 8°C.

62 The air under plastic cover had lower values of relative humidity in both seasons. This
63 condition can be attributed to the barrier that the plastic imposes to leaf wetness.

64 The treatments evaluated were the control, without the use of plastic cover, with anti-hail
65 screen and plastic cover.

66 The material used in the cover was plastic braided transparent polyethylene canvas (160 µm
67 thick), waterproofed in the first year (2018) of use in the vineyard. After the grapes were
68 harvested, the material (plastic tarpaulin) was removed, to increase durability, and replaced
69 immediately after pruning in the next agricultural cycle (allowing durability of up to 20 years).
70 The plastic cover was arranged along the planting line, following a north-south orientation,
71 seated on metallic wires and these over steel arches (tunnels with a height of 1.30 m from
72 the plant to the plastic and 3 m wide), fixed on the structure of the “Y” system.

73 The white anti-hail screen, with a mesh opening of 4 mm x 7 mm, fixed in the same
74 arrangement, seated on the same metallic structure used for the plastic, on metallic wires
75 and these on steel arches (tunnels with a height of 1, 30 from the plant to the plastic and 3 m
76 wide), fixed on the structure of the “Y” system (only placed for experimental purposes).

77 Phenology evaluations were carried out from the beginning of sprouting to the end of the
78 cycle, plant water potential, gas exchange, intensity of different wavelengths,
79 photosynthetically active radiation, photosynthetic pigments, vegetative and productive
80 attributes.

81 After performing the winter pruning in the productive cycles, after 14 days, evaluations were
82 carried out to estimate the sprouts of the plants for the phenology calculations. Data were
83 collected from 1 plant of each repetition, at 14-day intervals, throughout the
84 vegetative/productive cycle of the 2018/19 and 2019/20 harvests. Phenology was expressed
85 as a percentage of branches per point of the phenological scale developed by Lorenz et al.
86 [16], on the evaluation dates. In the same plant, the number of branches and total leaves per
87 plant were evaluated.

88 For the microclimatic characterization, the maximum, minimum, average temperatures and
89 relative humidity were obtained using a data logger (NOVUS, LOGBOX-RHT-LCD, country?)
90 at the height of the bunches. The equipment was in the field from August to September in
91 the 2019/20 and 2020/21 harvests.

92 In the evaluations of microclimatic variables in the two seasons (2019/20 and 2020/21),
93 changes caused by the use of plastic covering on the crop rows were observed. Plants
94 under plastic cover had higher maximum, average and minimum air temperatures compared
95 to those uncovered and under screen (Table 1). The average temperature difference
96 between the plants under the uncovered system and the anti-hail screen was 0.2 °C, and 0.5
97 °C between the uncovered system and the plastic cover in the year 2019/20. In the year
98 2020/21, these differences were 0.1 °C in the uncovered system and anti-hail screen, and
99 0.6 °C between the uncovered and the plastic cover, respectively. Study carried out by
100 Pedro Júnior et al. [13] cite higher temperature in the bunch of 'Syrah' vines produced under
101 plastic cover, compared to the uncovered environment. It can be seen, in general, that the
102 plastic cover delays the loss of heat in comparison to the uncovered system and the anti-hail
103 screen. Therefore, temperature directly affects plant development and the hormonal balance
104 of fruit trees [17], as well as the vegetative development and quality of the grapes produced
105 [18].

106 Table 1. Microclimatic variables of maximum, minimum, average temperatures and relative
107 humidity (%) in an uncovered cultivation environment, covered with anti-hail screen and with
108 plastic cover, in the 'Chardonnay' cultivar, from September to October, in the 2019 harvests
109 /20 and 2020/21, São Joaquim, SC.

Microclimatic Variables	Uncovered	Screen	Plastic cover
	Crop 2019/20		

Max. canopy (°C)	34,5	34,2	34,7
T. min. canopy (°C)	7,1	7,1	7,5
T. average canopy (°C)	20,6	20,4	21,1
UR. canopy (%)	79,1	81,4	77,9
Crop 2020/21			
Max. canopy (°C)	32,2	32,1	32,4
T. min. canopy (°C)	8,1	8,1	8,7
T. average canopy (°C)	20,0	20,1	20,6
UR. canopy (%)	81,2	80,6	79,6

110

111 The radiation spectrum was obtained with a spectrophotometer (APOGGEE INSTRUMENTS
 112 – model SS – 110), with detection of wavelengths from 340 to 810 nm, at midday. From the
 113 data obtained, the total radiation available in the control treatment and the radiation available
 114 for plants under anti-hail screen and plastic cover were quantified. The total radiation
 115 (without interference from the physical barrier) was quantified as 100% of the available light,
 116 and the other values calculated as a function of the total radiation, and then, the percentage
 117 of light retained by the physical barrier was determined. The spectroradiometer measured
 118 the amount of light in the ultraviolet (300-390 nm), blue (450-490 nm), green (490-580 nm),
 119 red (620-700 nm), far red (700-750 nm) ranges. The red/far red (V/Vd) ratio was calculated
 120 considering these last two wavelength ranges. Photosynthetically Active Radiation (RFA)
 121 was determined with a cepometer, Licor model leaf area index (LAI-2200C – Licor, USA).

122 The determination of leaf water potential (Ψ_{foliar}) was performed with a pressure chamber
 123 (PMS Instrument Co, model 1000, USA). The evaluations were carried out on leaves
 124 opposite the grape bunches, from 7 am to 3 pm, in December of the 2019/20 and 2020/21
 125 harvests, with intervals of two hours between the evaluations.

126 The number of branches and leaves was estimated per plant. The variables were analyzed
 127 according to the vegetative stage of the plants. Four representative branches were marked
 128 on the plants, with the aid of graduated rulers, and measurements were performed on one
 129 plant by repetition. The branches and leaves were not removed from the plants to avoid
 130 causing damage and injuries. The leaf attributes were quantified through the evaluation of
 131 area, dry mass and specific area, using ten leaves in each repetition. Leaf area (cm^2) was
 132 quantified using a leaf area integrator (Li-Cor, model LI-3100, USA). The dry mass (g) was
 133 quantified on an analytical balance, after drying in an oven at 65°C , with forced air
 134 circulation, for 72 hours. The specific leaf area ($\text{cm}^2 \text{g}^{-1}$) was calculated by dividing the area
 135 (cm^2) by the dry mass (g) of the leaves. The fresh weight of leaves on a plant (kg plant^{-1})
 136 was estimated by the number of leaves on the plant and multiplied by the fresh mass (g).

137 The levels of chlorophyll a, b and total (a+b), a/b and carotenoids ratio were obtained by
 138 collecting three disks of 26.6 mm in circumference and placed in amber glasses (10 mL),
 139 covered with aluminum foil and tape. adhesive, with 7 mL of DMSO (extractor reagent -
 140 dimethyl sulfoxide), incubated at 65°C for two hours. After total chlorophyll extraction, the
 141 liquid was pipetted into Elisa plates, and reading was performed in a microplate reader
 142 (brand, country model), at chlorophyll wavelengths at 649 nm, 665 nm and 480 nm for
 143 chlorophyll a, chlorophyll b and carotenoids, respectively. The levels of chlorophyll a, b and
 144 total and carotenoids were obtained by the formulas: $\text{Chl a } (\mu\text{L}^{-1}) = 12.47 \cdot (665 \text{ nm}) -$
 145 $3.63 \cdot (649 \text{ nm})$. $\text{Chl a } (\text{g L}^{-1}) = ((\text{Chl a} \cdot (\text{DMSO volume}) / 1000)) / \text{leaf disc leaf area } (\text{mm}^2)$
 146 $\cdot 10000$; $\text{Chl b } (\text{g L}^{-1}) = 25.06 \cdot (640 \text{ nm}) - 6.5 \cdot (665 \text{ nm})$, $\text{Chl b} = ((\text{Chl b} \cdot (\text{DMSO}$
 147 $\text{volume}) / 1000)) / \text{leaf area of the sheet } (\text{mm}^2) \cdot 10000$; $\text{Chl a} + \text{b} = 21.44 \cdot (649 \text{ nm}) + 5.97 \cdot (665$
 148 $\text{ nm})$. $\text{Chl a+b } (\text{g L}^{-1}) = ((\text{Chl a} \cdot (\text{DMSO volume}) / 1000)) / \text{leaf disc leaf area } (\text{mm}^2) \cdot 10000$; the

149 a/b ratio by dividing the values of a and b, carotenoids = $(1000 \cdot (470 \text{ nm}) - 2.14 \cdot (\text{Chl a}) -$
150 $70.16 \cdot (\text{Chl b})) / 220$, Carotenoid (g L^{-1}) = $((\text{carotenoids} \cdot (\text{DMSO volume}) / 1000)) / \text{leaf area of}$
151 $\text{leaf disc (mm}^2) \cdot 10000$ adapted [19].

152 Net photosynthesis (A ; $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$), stomatal conductance (g_s); $\text{mol H}_2\text{O m}^{-2} \text{ s}^{-1}$), ratio
153 between CO_2 (C_i/C_a), internal concentration of CO_2 (C_i), quantum yield of photosystem I
154 (FS1) and maximum quantum yield of photosystem II (FSII), relative rate of electron
155 transport (ETR) were generated with an infrared gas analyzer (LI-6400, LI-COR, USA),
156 equipped with an open-top chamber. These evaluations were performed on the color change
157 of the berries.

158 The number of bunches and weight of bunches (kg) per plant were obtained by collecting all
159 bunches from two plants, which were counted and weighed, and the average of these values
160 was obtained. The individual bunch weight (g) was estimated by the total weight of bunches
161 and divided by the number of bunches. Yield (kg ha^{-1}) was estimated by harvesting two
162 plants, and then weighting yield to the total number of plants in one hectare.

163 The design was in a completely randomized design, with three treatments and five
164 replications per treatment, with two plants per replication. The program for statistical analysis
165 was SISVAR 2.0, with LSD test ($p < 0.05$).

166

167 **3. RESULTS AND DISCUSSION**

168

169 A reduction of RFA was observed for plants under plastic cover and anti-hail screen, in
170 relation to uncovered plants, respectively, of 39% and 36% in the 2019/20 crop, between
171 uncovered plants and under anti-hail screen; the difference was 58% and 35% in the
172 2020/21 crop, between uncovered plants and plants under plastic cover (Table 2). The
173 greater reduction of RFA under plastic cover in the second crop may be related to the
174 accumulation of dirt and loss of transparency of the plastic placed in the cultivation line.
175 Although the material is removed at the end of the harvest to increase its durability, the
176 condition of accumulation of residues from the previous year is a relevant factor. Other
177 studies carried out in Brazil also demonstrated the reduction of RFA under plastic cover.
178 Azevedo et al. [20] observed a decrease in the number of grape plants with the cover, at 50
179 cm above the plant canopy. Mota et al. [21], Cardoso et al. [22, 23], Chavarria et al. [24] also
180 found a reduction in RFA under plastic cover. According to Chavarria et al. [25], the
181 reduction of RFA can be 30% in covered vineyards, compared to the uncovered system.
182 This condition is due to the imposition of the physical barrier imposed by the plastic, which
183 resulted in a decrease in the availability of light [26]. According to Leitão et al. [12], the RFA
184 was reduced by 40%, at 50 cm from the canopy, on vines under plastic cover in the São
185 Francisco river valley.

186 The quantity and quality of light incident on the canopy of plants greatly affects the
187 development of vines, as they are directly linked to all physiological processes of plants [27].
188 The amount of available light is essential for the synthesis of photosynthetic pigments in
189 plants, since the chlorophyll content of the leaves is directly linked to the reaction centers of
190 the photosystems, a condition that leads to the growth and development processes of plants
191 [28].

192

193 Table 2. Photosynthetically active radiation (RFA; W m^{-2}) incident on the height of vine
194 bunches of the 'Chardonnay' cultivar, in uncovered cultivation systems, covered with anti-hail

195 screen and with plastic cover, in December, 2019 harvests /20 and 2020/21. Sao Joaquin,
196 SC.

Treatment	Crop 2019/20	Crop 2020/21
Uncovered	2317 a	1613 a
anti-hail screen	1492 b	1142 b
Cob. Plastic	1416 b	872 b
CV %	17,7	21,0

197 Means followed by the same letter, in the columns, did not differ by the LSD test ($p < 0.05$).
198

199 It was observed that the plastic cover was able to intercept the ultraviolet (UV) light incident
200 on the canopy (Table 3). Lights in the blue, red and far red bands were also intercepted by
201 the plastic cover, but at a lower intensity than the UV. Red light is another factor that slows
202 down the leaf senescence process. The reduction of available red light in the canopy allows
203 leaves to remain active for longer periods of time [28], as it presents late leaf senescence.
204 The ratio of red and far red (V/Vd) was higher for plants without plastic cover. Plants under
205 hail net had a lower V/Vd ratio, a condition that causes the plant to have an increase in
206 vegetative growth, changing its characteristics to adapt to the light restriction imposed by the
207 cover [24, 27, 29]. Plants subjected to shading have low respiratory rates and absorb
208 available light, which makes them more efficient in terms of the use of radiation, a condition
209 that reduces the waste of photons in the leaf, due to the smaller number of cells. According
210 to Batista et al. [30], the quality of light available in plant canopy is directly linked to the
211 production of structural genes that contribute to the adaptation of plants to shaded
212 environments.

213

214 Table 3. Light intensity at different wavelengths, and V/Vd ratio, at the height of vine
215 bunches in the 'Chardonnay' cultivar, in an uncovered, covered cultivation system, with anti-
216 hail screen and with plastic cover, in December, 2019/20 harvests,. Sao Joaquin, SC

Radiation attributes	Uncovered	Screen	Plastic cover	CV %
Ultraviolet	76,7 a	76,6 a	12,1 b	11,3
Blue	27,8 a	21,2 b	16,2 c	13,0
Green	67,9 a	37,4 b	40,4 b	21,7
Red (V)	60,8 a	37,4 b	39,1 b	20,3
Far red (Vd)	37,3 a	27,5 b	26,7 b	15,1
V/Vd Ratio	1,63 a	1,29 c	1,47 b	5,92

217 Means followed by the same letter, in the lines, do not differ by the LSD test ($p < 0.05$).
218

219 It was observed in Figure 1 that the water potential of the plants was adequate for the
220 development of the vine, in the three conduction systems, which must be between -0.2 and -
221 0.6 MPa. This condition of no water deficit for plants under plastic cover is explained by the
222 high rainfall that occurs in the region where the work was carried out.
223

224

225 Figure 1. Water potential of leaves in the cultivar Chardonnay, measured at three periods of
226 the day (7:00, 9:00, 11:00, 13:00, 15:00 hours), in the month of December, in the 2019/2019
227 harvests 20 and 2020/21, in uncovered systems, under screen and plastic cover, in the city
228 of São Joaquin, SC.

229

230 For the variables net photosynthesis (A), stomatal conductance (gs), CO₂ (Ci/Ca), CO₂(Ci),
231 Photosystem I (FS I), maximum quantum yield of photosystem II (PhiPS2) and relative rate
232 of electron transport (ETR) there was no difference between treatments (Table 4). These

233 results show that there is no damage to plants grown under plastic cover in terms of
 234 photosynthetic capacity. The quality of light is directly linked to the photosynthetic processes
 235 of plants, an example of this condition according to WANG et al [28] is that reduction in blue
 236 light delays the process of leaf senescence, as there is no degradation of chlorophyll present
 237 in chloroplasts. Plants are able to effectively compensate for the reduced availability of light
 238 for their development. This condition improves the CO₂ assimilation characteristics,
 239 supporting the transport of electrons and maintaining the efficiency of photosystem II. Plants
 240 can change their leaf morphology in terms of the arrangement of chloroplasts [31] and by
 241 changing the thickness of leaf tissues, responding according to the amount of light in the
 242 environment [26]. According to Da Silva et al. [17], the increase in temperature and reduction
 243 in relative humidity can provide a lower rate of net photosynthesis, a condition that directly
 244 reflects on the stages of plant flowering and berry maturation.

245
 246 Table 4. Gas exchange (gs) in vine bunches of the 'Chardonnay' cultivar, in uncovered
 247 cultivation systems, covered with anti-hail screen and with plastic cover, in December,
 248 2018/19 and 2019/20 harvests São Joaquim, SC.

Gas exchange	Uncovered	Screen	Plastic cover	CV %
	Crop 2018/219			
A ($\mu\text{mol CO}_2\cdot\text{m}^{-2}\cdot\text{s}^{-1}$)	13,5 ns	13,5	14,6	8,87
gs ($\text{mol H}_2\text{O}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$)	0,19 ns	0,19	0,20	9,21
Ci/Ca	0,67 ns	0,68	0,66	7,50
Ci	254,9 ns	258,6	248,1	3,32
Fs	574,0 ns	545,5	585,4	8,64
Fv/Fm'	0,52 ns	0,53	0,56	6,21
PhiPS2	0,32 ns	0,35	0,35	12,9
ETR ($\mu\text{mol m}^{-2}\cdot\text{s}^{-1}$)	135,4 ns	144,4	152,7	10,5
Crop 2020/21				
A ($\mu\text{mol CO}_2\cdot\text{m}^{-2}\cdot\text{s}^{-1}$)	14,5 ns	14,4	15,2	9,24
gs ($\text{mol H}_2\text{O}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$)	0,23 ns	0,23	0,24	13,8
Ci/Ca	0,67 ns	0,67	0,67	5,32
Ci	267,1 ns	267,2	266,4	4,82
Fs	706,1 ns	703,3	700,0	28,3
Fv/Fm'	0,52 ns	0,53	0,57	8,62
PhiPS2	0,29 ns	0,30	0,28	16,4
ETR ($\mu\text{mol m}^{-2}\cdot\text{s}^{-1}$)	136,3 ns	149,3	154,4	16,2

249 Means followed by the same letter, in the lines, do not differ by the LSD test ($p < 0.05$). A – Net carbon
 250 assimilation rate. g – Stomatal conductance. Ci/Ca – ratio between indoor and ambient CO₂
 251 concentrations. Internal ci-concentration of CO₂. Fs- photosystem I. Fv/Fm'- maximum quantum yield
 252 of photosynthesis II. PhiPS2 - photosynthesis II. ETR - relative rate of electron transport. *means
 253 followed by the same letter do not differ statistically by the LSD test (5% probability). *CV % -
 254 coefficient of variation.
 255

256 Chlorophyll a contents and chlorophyll a/b ratio in leaves were higher for plants under plastic
 257 cover in the 2018/19 crop year. For chlorophyll b, the highest content was obtained in plants
 258 under hail net and plastic cover (Table 5). Plants with higher chlorophyll a/b ratios, according
 259 to Wang et al. [28] and Hairmansis et al [32], present greater disorders in chloroplasts. This
 260 condition explains the need for adaptation of cell arrangements in leaves to maintain their
 261 photosynthetic capacity. The highest levels of chlorophyll b in shaded environments is
 262 related to the evaluation of plants and their adaptation to environmental changes [33]. In the
 263 2019/2020 crop year there was no difference between treatments for all variables related to
 264 chloroplast pigments. In the 2020/21 crop year, there was a difference only for the
 265 chlorophyll a+b variable, which was higher in plants under plastic cover, compared to plants
 266 under hail and uncovered screens. Kong et al. [34], in a study carried out with the quality of
 267 light, temperature and photosynthesis in cherry tomatoes, observed that plants subjected to
 268 higher temperatures had higher values of chlorophyll a, b and a+b. The data found in this
 269 work for the protected cultivation of vines corroborate those presented by Kong et al. [34].
 270 Work carried out by Chavarria et al. [26] also showed higher levels of photosynthetic
 271 pigments in vines produced under plastic cover. According to Streit et al. [31], plants
 272 subjected to high light intensities can undergo the process of photoinhibition, as a way of
 273 protecting the plant from excess light and, consequently, reducing chlorophyll levels.
 274 According to Wang et al. [26], plants subjected to a higher incidence of blue light tend to
 275 reduce the production of chlorophyll in the leaves, a condition observed in Table 5, which
 276 shows plants without cover present greater availability of blue light and, consequently, lower
 277 production of chlorophyll. The levels of carotenoids in the leaves did not differ between
 278 treatments (Table 5).

279 Table 5. Levels of photosynthetic pigments in vines in the 'Chardonnay' cultivar, in
 280 uncovered cultivation systems, covered with anti-hail screen and with plastic cover, in
 281 December, 2018/19, 2019/20 and 2020/21 harvests. Sao Joaquin, SC

Photosynthesizers pigments	Uncovered	Screen	Plastic cover	CV%
Crop 2018/19				
Chlorophyll a (g L ⁻¹)	2,50 c	3,14 b	3,23 a	2,02
Chlorophyll b (g L ⁻¹)	1,33 b	1,60 a	1,57 a	2,45
Chlorophyll a+b	3,83 b	4,74 a	4,80 a	1,41
a/b ratio	1,88 b	1,97 b	2,06 a	3,20
Carotenoids (g L ⁻¹)	0,55 ns	0,59	0,56	7,36
Crop 2019/20				
Chlorophyll a (g L ⁻¹)	1,93 ns	1,75	2,08	28,8
Chlorophyll b (g L ⁻¹)	1,17 ns	1,13	1,25	29,4
Chlorophyll a+b	3,09 ns	2,88	3,33	35,7
a/b ratio	1,66 ns	1,54	1,66	31,7
Carotenoids (g L ⁻¹)	0,38 ns	0,31	0,38	37,8
Crop 2020/21				
Chlorophyll a (g L ⁻¹)	1,35 ns	1,22	1,43	34,4
Chlorophyll b (g L ⁻¹)	0,81 ns	0,74	0,82	30,0
Chlorophyll a+b	2,16 ab	1,97 b	2,25 a	20,6
a/b ratio	1,66 ns	1,64	1,75	30,5

7	135 ns	130	128	6,8
8	210 a	205 a	197 b	2,2
9	225 b	221 b	240 a	2,1
Crop 2019/2020				
0	52 ns	56	52	8,7
1	81 a	81 a	72 b	6,1
5	109 a	109 a	83 b	10,6
6	140 b	138 b	125 a	4,1
7	148 b	148 b	133 a	3,6
8	199 b	207 a	196 b	5,5
9	221 b	221 b	255 a	2,0

321 *(0)-budding, (1)-leaf development, (5)-inflorescence appearance, (6)-flowering, (7)-fruit development,
 322 (8)-fruit maturation, (9)-beginning of dormancy. *Evaluations carried out only in the 2018/19 and
 323 2019/20 harvests. *Means followed by the same letter do not differ statistically by the LSD test (5%
 324 probability). *CV (%) - coefficient of variation.

325

326 There was no difference between treatments for the number of branches per plant in the
 327 2018/19, 2019/20 and 2020/21 crop years, and in the 2019/20 year there was no difference
 328 for specific leaf area (AFE). The number of leaves per plant and the PF of leaves per plant
 329 were higher for plants treated with plastic cover in the three seasons. The AFE in the
 330 2018/19 and 2020/21 harvests was higher in plants conducted under plastic cover, differing
 331 from plants conducted in an uncovered system and under hail net (Table 7). According to
 332 Salem et al. [38] and Chavarria et al. [25], vines under plastic cover have higher AFE. This
 333 condition is directly linked to the increase in temperature and the availability of water [42].
 334 Another factor is the quality of light incident on the canopy, since the greater vegetative
 335 growth under the plastic cover is related to the lower V/Vd ratio, as well as the greater
 336 availability of blue and red radiation, which falls on the canopy and favors the synthesis of
 337 pigments. photosynthetic agents [28].

338 Table 7. Vegetative attributes measured in December, in the 2018/19, 2019/20 and 2020/21
 339 harvests, in uncovered systems, under screen and plastic cover in the 'Chardonnay' cultivar,
 340 in the city of São Joaquim, SC.

Vegetative attributes	Crop 2018/19			CV %
	Uncovered	Screen	Plastic cover	
Number of branches plant ⁻¹	26 ns	25	27	8,8
No. of leaves plant ⁻¹	623 b	531 c	731 a	7,9
PF of leaves plant ⁻¹ (kg)	3,277 a	2,915 b	4,102 a	9,7
AFE (cm g ⁻¹)	14 b	15 b	18 a	11,0
Crop 2019/20				
Number of branches plant ⁻¹	14 ns	14	15	7,8
No. of leaves plant ⁻¹	399 b	381 b	472 a	7,3
PF of leaves plant ⁻¹ (kg)	1,223 b	1,332 b	1,793 a	8,1
AFE (cm g ⁻¹)	15 ns	18	16	21,0
Crop 2020/21				
Number of branches plant ⁻¹	26 ns	25	27	8,8
No. of leaves plant ⁻¹	614 b	514 b	726 a	6,9

PF of leaves plant ⁻¹ (kg)	1,948 b	1,638 b	3,139 a	15,5
AFE (cm g ⁻¹)	15 b	15 b	22 a	3,5

341 *Number-number. *PF – fresh weight. AFE – specific leaf area. Means followed by the same letter do
 342 not differ statistically by the LSD test (5% probability). CV % - coefficient of variation.

343
 344 Plants grown under plastic cover had a higher number of clusters per plant compared to
 345 plants in an uncovered system and under screen, in the 2018/19 harvest. In the 2019/20 and
 346 2020/21 crops, plants under plastic cover and anti-hail screen did not differ from each other,
 347 however, both covered environments provided a greater number of clusters per plant
 348 compared to the uncovered system. In the three seasons of study, the weight of bunches per
 349 plant and productivity were higher in plants under plastic cover, in relation to plants under
 350 screen and in an uncovered system, which there was no difference between the last two
 351 systems of conduction. In the 2019/20 and 2020/21 crop years, the weight of individual
 352 bunches was higher for plants in a plastic cover system compared to the system under
 353 screen and uncovered, which showed no differences between them. However, in the
 354 2018/19 crop year, plants under plastic cover and under hail net did not show differences
 355 between them, and there was no difference between the screen cover and uncovered
 356 system (Table 8). In the 2018/19 crop, no differences were observed for plant bunch weight
 357 between plants grown under plastic cover and screen.

358 The gain in number of bunches per plant under plastic cover, compared to the uncovered
 359 system, was 28%, 17% and 50% in the 2018/19, 2019/20 and 2020/21 crops, respectively.
 360 For plants conducted under anti-hail nets, the gains, in relation to the uncovered system,
 361 were 3%, 17%, and 29%, in the 2018/19, 2019/20 and 2020/21 harvests, respectively. For
 362 the weight of bunches (g) plants under plastic cover, in relation to uncovered, they showed
 363 an increase of 10%, 49% and 30%, in the 2018/19, 2019/20 and 2020/21 harvests,
 364 respectively. For the anti-hail net, gains were lower, 6%, 11% and 0.9%, compared to
 365 uncovered, in the 2018/19, 2019/20 and 2020/21 harvests, respectively. The results of the
 366 present work corroborate those of Pedro Júnior and Hernades [41], who reported higher
 367 values for number of bunches, weight of bunches per plant and individual weight of bunches
 368 in plants grown under plastic cover.

369 These differences between the numbers of bunches can be attributed to the way of
 370 management in the pruning, conduction and thinning of bunches. Another important factor
 371 was the occurrence of late frost in November 2019, a condition that caused the death of
 372 shoots and defoliation in up to 90% in plants conducted in an uncovered system and under
 373 an anti-hail screen. This condition directly affected the productivity of the 2019/20 crop, and
 374 provided a reduction in production in the 2020/21 crop, as the physiological condition and
 375 the amount of reserve were compromised for the following year. The plastic cover provided
 376 productivity increases of 49%, 116% and 65%, compared to uncovered, in the 2018/19,
 377 2019/20 and 2020/21 harvests, respectively. The anti-hail net increased productivity by 17%,
 378 18%, 60%, compared to uncovered, in the 2018/19, 2019/20 and 2020/21 harvests,
 379 respectively.

380 The productivity gains corroborate the results found by Azevedo et al. [20] with grapes
 381 produced under plastic cover in the São Francisco River Valley, and for grapes produced in
 382 Bento Gonçalves [40]. Pedro Júnior and Hernandez [41] also observed higher yields in
 383 grapes produced under plastic cover, in summer and winter crops. These authors attributed
 384 the productivity gains to the lower incidence of diseases and lower defoliation in the plants.

385 Table 8. Productive attributes of vines in the cultivar 'Chardonnay', in uncovered cultivation
 386 systems, covered with anti-hail net and with plastic cover, in December, 2018/19, 2019/20
 387 and 2020/21 harvests. Sao Joaquin, SC.

Productive attributes	Uncovered	Screen	Plastic cover	CV %
Crop 2018/19				
No. of plant bunches ⁻¹	31 b	32 b	43 a	11,1
Weight bunches kg plant ⁻¹	3,9 b	4,6 b	6,4 a	15,0
bunch weight (g)	130,8 b	138,2 ab	145,3 a	6,77
Productivity (Mg.ha ⁻¹)	11,221 b	13,460 b	18,482 a	15,0
Crop 2019/20				
No. of plant bunches ⁻¹	4,7 b	6,2 a	5,9 a	11,0
Weight bunches kg plant ⁻¹	0,08b	0,08b	67,3a	52,0
bunch weight (g)	83,9 b	67,2 b	128 a	16,7
Productivity (Mg.ha ⁻¹)	0,228 b	0,251 b	2,158a	56,0
Crop 2020/21				
No. of plant bunches ⁻¹	19 b	37 a	38 a	21,0
Weight bunches kg plant ⁻¹	2,1 b	0,9 b	6,1 a	34,0
bunch weight (g)	111 b	112 b	159 a	8,9
Productivity (Mg.ha ⁻¹)	6,104 b	2,441 b	17,449 a	34,2

388 Means followed by the same letter, in the lines, do not differ statistically by the LSD test (5%
389 probability). CV % - coefficient of variation.
390

391 4. CONCLUSION

392
393 The objective of the work was "to evaluate the effect of protected cultivation on phenology,
394 water relations, gas exchange, vegetative growth and production in 'Chardonnay' vine."
395 Thus, it must conclude on how the plastic cover and the screen influenced phenology, water
396 relations, gas exchange, vegetative growth and production in 'Chardonnay' vines.
397

398 COMPETING INTERESTS

399
400 Authors have declared that no competing interests exist.
401

402 AUTHORS' CONTRIBUTIONS

403
404 This work was carried out in collaboration between all authors. Authors CVTA, CAS and
405 KCS designed the study and wrote the protocol. Authors CLF, AL and ASH performed the
406 statistical analysis and wrote the first draft of the manuscript. Author KCS managed the
407 analyses of the study and the translation of the manuscript. All authors read and approved
408 the final manuscript.
409

410 **COMPETING INTERESTS DISCLAIMER:**

411

412 **AUTHORS HAVE DECLARED THAT NO COMPETING INTERESTS EXIST. THE**
413 **PRODUCTS USED FOR THIS RESEARCH ARE COMMONLY AND**
414 **PREDOMINANTLY USE PRODUCTS IN OUR AREA OF RESEARCH AND**

415 **COUNTRY. THERE IS ABSOLUTELY NO CONFLICT OF INTEREST BETWEEN**
416 **THE AUTHORS AND PRODUCERS OF THE PRODUCTS BECAUSE WE DO**
417 **NOT INTEND TO USE THESE PRODUCTS AS AN AVENUE FOR ANY**
418 **LITIGATION BUT FOR THE ADVANCEMENT OF KNOWLEDGE. ALSO, THE**
419 **RESEARCH WAS NOT FUNDED BY THE PRODUCING COMPANY RATHER IT**
420 **WAS FUNDED BY PERSONAL EFFORTS OF THE AUTHORS.**
421

422 REFERENCES

- 423
424
425
- 426 1. Chelotti, M.C.; Medeiros, R. M. V. CARTOGRAPHIES OF VITICULTURE IN MINAS
427 GERAIS: from the genesis in the South to the expansion to the North. *InterEspaço: Journal*
428 *of Geography and Interdisciplinarity*, 2021;e202106-e202106. DOI: 10.18764/2446-
429 6549.e202106.
 - 430 2. Felippeto, J.; Caliari, V.; Brighenti, E.; Nesi, C.N. Thermal requirement and
431 physicochemical characteristics of different vine varieties in São Joaquim, SC. *Agropecuária*
432 *Catarinense*, 2021;34:61-66. DOI: 10.52945/rac.v34i1.1056.
 - 433 3. Ferri, G.K. History and viticulture project: innovative teaching and sustainable
434 socioeconomic development in Serra Catarinense. *Revista GepesVida*, 2019;1:59-66.
 - 435 4. Caldana, N.F.S.; Da Silva, G.M.F.; Martelocio, A.C.; Nitsche, P.R., Caramori, P.H.
436 Characterization of hailstorm occurrences and their socioeconomic impacts in the state of
437 Paraná. *Agrometeors*, 2019;27:271-284. DOI: 10.31062/agrom.v27i2.26455.
 - 438 5. Brighenti, A.F.; Brighenti, E.; Bonin, V.; Rufato, L. Phenological characterization and
439 thermal requirement of vine varieties in São Joaquim, Santa Catarina-Brazil. *Rural Science*,
440 2013;43:1162-1167. DOI: 10.1590/S0103-84782013005000082.
 - 441 6. Brighenti, A.F., Malinovski, L.I., Stefanini, M., Vieira, H.J., Silva, A.L.DA. Comparison
442 between the wine-growing regions of São Joaquim - SC, Brazil and San Michele All'Adige -
443 TN, Italy. *Brazilian Journal of Fruticulture*, 2015;37:281-288. DOI: 10.1590/0100-2945-
444 093/14.
 - 445 7. Martins, W.A.; Santos, S.C.; Jara, R.S., Souza, J.L.A.C.; Galvão, J.R.; Biscaro, G. A.
446 Phenology and thermal demand of blackberry cv. Tupy. *Journal of Agricultural Sciences*,
447 2019;42:720-730. DOI: 10.19084/rca.17529.
 - 448 8. Bosco, L.C.; Bergamaschi, H.; Cardoso, L.S.; De Paula, V.A., Marodin, G.A.B.;
449 Nachtigall, G. R. Apple production and quality when cultivated under anti-hail cover in
450 Southern Brazil. *International Journal of Biometeorology*, 2015;59:773-782. DOI
451 10.1007/s00484-014-0893-6.
 - 452 9. Pedro Jr, M.J.; Hernades, J.L. Fungal diseases in vines cultivated in the open and under
453 waterproof cover. *Summa Phytopathologica*, 2020;46:155-160. DOI: 10.1590/0100-
454 5405/203640.
 - 455 10. Hawerroth, F.J.; Nachtigall, G. R. Autumn and winter weather conditions and their
456 influences on the 2016/17 apple crop in the region of Vacaria, RS. Bento Gonçalves, RS:
457 Embrapa Grape and Wine, 2016. (Embrapa Grape and Wine. Technical Communiqué, 190).

- 458 11. De Oliveira, B.; Egypt, R.; Laureano, O.; Castro, R.; Pereira, G.E.; Silva, J.M.R. Climate
459 effects on physicochemical composition of Syrah grapes at low and high-altitude sites from
460 tropical grown regions of Brazil. *Food Research International*, 2019;121:870 – 870. DOI:
461 10.1016/j.foodres.2019.01.011.
- 462 12. Leitão, M.V.B.R.; Azevedo, P.V.; Lima, P.C.S.; Oliveira, G.M.; Santos, C.A.C. Influence
463 of plastic covering on the microclimate in vineyards in the San Francisco River Valley
464 Region. *Brazilian Journal of Meteorology*, 2017;32:399-407. DOI: 10.1590/0102-
465 77863230007.
- 466 13. Pedro Jr, M.J.; Hernades, J.L.; Santos, A.O.; Loureiro, A.C. Microclimate, production and
467 composition of 'Syrah' must grown under plastic cover. *Journal of Agroveterinary Sciences*,
468 2019;18:34-46. DOI: 10.5965/223811711832019374.
- 469 14. Holcman, E., Sentelhas, P.C., Spósito, M.B., Conceição, A.A.F., Couto, H.T.Z. Yield and
470 quality of grapes cultivated under plastic coverings. *Journal of Phytopathology*, 2019;2:1-11.
471 DOI: 10.1111/jph.12786.
- 472 15. Koppen, W. Die Wärmezonen der Erde, nach der Dauer der heissen, gemässigten und
473 kalten Zeit und nach der Wirkung der Wärme auf die organische Welt betrachtet.
474 *Meteorologische Zeitschrift*, 1884;1:215-226.
- 475 16. Lorenz, D.H.; Eichhorn, K.W.; Bleiholder, H.; Klose, R.; Meier, U.; Weber, E.
476 Phenological growth stages of the grapevine (*Vitis vinifera* L. ssp. *vinifera*) - codes and
477 descriptions according to the extended BBHC scale. *Australian Journal of Grape and Wine*
478 *Research*, 1995;1:100-103. DOI: 10.1111/j.1755-0238.1995.tb00085.x.
- 479 17. Da Silva, T.R.G.; Da Costa, M.L.A.; Farias, L.R.A.; Dos Santos, M.A.; De Rocha, J.J.;
480 Silva, J.V. Abiotic factors in plant growth and flowering. *Society and Development*,
481 2021;10:234-243. DOI: 10.33448/rsd-v10i4.13817.
- 482 18. Maranhao, K.R.; Reis, L.L.; Brito, M.R.; Benjamin, R.S.C. Influence of water deficit on
483 the ecophysiology characteristics of vines cv. Brs Violeta and Pinot Noir. *PesquisAgro*,
484 2020;3: 52-64. DOI: 10.33912/pagro.v3i1.657.
- 485 19. Wellburn, A. R. The spectral determination of chlorophyll a and b, as well as total
486 carotenoids, using various solvents with spectrophotometers of different resolution. *Journal*
487 *of Plant Physiology*, 1994;144:307-313. DOI: 10.1016/S0176-1617(11)81192-2.
- 488 20. Azevedo, P.V.; Dos Santos, C.A.C.; Lima, P.C.S.; Leitão, M.M.V.B.; Oliveira, G.M.;
489 Abrahão, R. Effect of the plastic cover on the productivity and profitability of vineyard in the
490 São Francisco River Valley, Brazil. *Brazilian Journal of Meteorology*, 2020;35:81-88. DOI:
491 10.1590/0102-7786351008.
- 492 21. Mota, C.S.; Amarante, C.V.T.; Of the Santos, H.P.; Albuquerque, J.A. Water availability,
493 solar radiation and photosynthesis in 'Cabernet Sauvignon' vines under protected cultivation.
494 *Brazilian Journal of Fruticulture*, 2009;31:432-439. DOI: 10.1590/S0100-
495 29452009000200017.
- 496 22. Cardoso, L.S.; Bergamaschi, H.; Comarim, F.; Chavarria, G.; Marodin, G.A.B.; Dalmago,
497 G.A.; Santos, H.P. Dos, Mandelli, F., Micrometeorological alterations in vineyards by the use
498 of plastic coverings. *Brazilian Agricultural Research*, 2008;43:441-447. DOI: 10.1590/S0100-
499 204X2008000400001.

- 500 23. Cardoso, L.S.; Bergamashi, H.; Comin, F.; Chavarria, G.; Marodin, G.A.B.; Dalmago,
501 G.A.; Dos Santos, H.P.; Mandelli, F. Patterns of solar radiation interception in vineyards with
502 and without plastic cover. *Brazilian Journal of Fruticulture*, 2010;32:161-171. DOI:
503 10.1590/S0100-29452010005000029.
- 504 24. Chavarria, G.; Tomm, G.O.; Muller, A.; Mendonca, H.F.; Mello, N.; Betto, M.S. Leaf area
505 index in canola cultivated under variations in spacing and seeding density. *Rural Science*,
506 2011;41:2084-2089. DOI: 10.1590/S0103-84782011001200008.
- 507 25. Chavarria, G.; Santos, H.P.D.; Mandelli, F.; Marodin, G.A.B.; Bergamaschi, H.; Cardoso,
508 L. S. Phenological characterization and thermal requirement of the cultivar Moscato Giallo
509 under plastic cover. *Brazilian Journal of Fruticulture*, 2009;31:119-126. DOI: 10.1590/S0100-
510 29452009000100018
- 511 26. Chavarria, G.; Santos, H.P. Dos, Castro, L.A.S.De Marodin, G.A.B., Bergamaschi, H.
512 Anatomy, chlorophyll content and photosynthetic potential in grapevine leaves under plastic
513 cover. *Brazilian Journal of Fruticulture*, 2012;34:661-668. DOI: 10.1590/S0100-
514 29452012000300003.
- 515 27. Taiz L.; Zeiger, E. *Plant physiology*: 5th ed. Porto Alegre: Editora Artmed, 2015.
- 516 28. Wang, S.; Wang, X., Shi, X.; Wang, B.; Zheng X.; Wang, H.; Liu, F. Red and blue lights
517 significantly affect photosynthetic properties and ultrastructure of mesophyll cells in
518 senescing grape leaves. *Horticultural Plant Journal*, 2016;2:82–90. DOI:
519 10.1016/j.hpj.2016.03.001.
- 520 29. Kerbauy, G.B. *Plant Physiology*, 2nd ed. Rio de Janeiro: Guanabara Koogan, 2008.
- 521 30. Batista, DS, Felipe, SHS, Silva, TD, De Castro, KM, Rodrigues, TCM, Miranda, NA,
522 Rios-Rios, AM, Faria, DV, Fortini, EA, Chagas, K., Silva, GT, Xavier, A., Arenciba, AD,
523 Otoni, WC Light quality in plant tissue culture: does it matter? *In Vitro Cellular &*
524 *Developmental Biology – Plant*, 2018;12:1-21. DOI: 10.1007/s11627-018-9902-5.
- 525 31. Streit, N.M.; Canterle, L.P.; Canto, M.W.; Hecktheuer, L.H.H. The chlorophylls. *Rural*
526 *Science*, 2005;35:748-755.
- 527 32. Hairmansis, A., Yullianida, Y., Hermanasari, R., Lestari, P. Development of Shading
528 Tolerant Rice Varieties Suitable for Intercropping Cultivation in Agroforestry Systems. *E3S*
529 *Web of Conferences*, 2021;305:1-5. DOI: 10.1051/e3sconf/202130507001.
- 530 33. Sonobe, R. Hirono, Y. Oi, A. Quantifying chlorophyll a and b content in tea leaves using
531 hyperspectral reflectance and deep learning. *Remote Sensing Letters*, 2020;11:933–942.
532 DOI: 10.1080/2150704X.2020.1795294.
- 533 34. Kong, L.; Em, Y.; Jiao, X.; Liu, X.; Xu, Z. Interactive regulation of light quality and
534 temperature on cherry tomato growth and photosynthesis. *Environmental and Experimental*
535 *Botany*. 2021;182:104326. DOI: 10.1016/j.envexbot.2020.104326.
- 536 35. Alonso, F.; Chiamolera, F.M.; Hueso, J.J.; Gonzalez, M.; Cuevas, J. Heat unit
537 requirements of “Flame Seedless” table grape: A tool to predict its harvest period in
538 protected cultivation. *Plants*, 2021;10:1-13. DOI: 10.3390/plants10050904.

- 539 36. Zhang, H.; Gao, D.; Li, D.; Li, X. Studies on developments of quality physiology of peach
540 in greenhouse. *Chinese Agricultural Science Bulletin*, 2005;28: 286–288.
- 541 37. Martínez-Gómez, P.; Rahimi Devin, S.; Salazar, J.A., López-Alcolea, J.; Rubio, Martínez-
542 García, P.J. Principles and prospects of prunus cultivation in greenhouse. *Agronomy*,
543 2021;11:1-9. DOI: 10.3390/agronomy11030474.
- 544 38. Salem, E.H.; Gouda, F.E.Z.M.; Abdel-Raman, M.A.A. SVU Effects of plastic covering
545 and potassium fertilization source on growth and fruiting of flame seedless grapevines.
546 *International Journal of Agricultural Sciences*, 2021;3:89-100. DOI:
547 10.21608/svuijas.2021.65024.1083.
- 548 39. Duarte Júnior, J.B.; Manechini, W. Cultivars of *Coffea arabica* L, in different planting
549 densities after implantation and the frost event. *Brazilian Journal of Development*,
550 2021;7:42497-42511. DOI:10.34117/bjdv7n4-626.
- 551 40. Comin, F., Bergamaschi, H., Heckler, B.M., Dos Santos, H.P., Alba, D., Saretta, E.
552 Microclimate and production of 'Niagara Rosada' vines in organic cultivation under plastic
553 cover. *Brazilian Journal of Fruticulture*, 2012;34:152-159. DOI: 10.1590/S0100-
554 29452012000100021.
- 555 41. Pedro Junior, MJ, Hernandes, JL, Blain, GC, Rolim, G. de S., Microclimate in 'Niagara
556 rosada' vineyards in different management systems during winter and summer crops,
557 *Revista Brasileira de Fruticultura*, 2013 ;35:151-158. DOI: 10.1590/S0100-
558 29452013000100018.
- 559 42. Giorgi, E.G.; Sadras, V.O.; Keller, M.; Peña, J.P. Interactive effects of high temperature
560 and water deficit on Malbec grapevines. *Australian Journal of Grape and Wine Research*,
561 2019;25:345-356. DOI: 10.1111/ajgw.12398.