

# Assessment of the potential for methane emissions caused by the livestock sector in the region of Kankan, Guinea

## ABSTRACT

Man has always organized his living space near animal farms. Nevertheless, the population growth and the healthiness due to the droppings of these animals have motivated work on the recovery of animal waste. This study follows this logic. It consists in the evaluation of the potential of methane emissions due to the livestock sector in the Administrative Region of Kankan. The methodological approach adopted for this study consists in carrying out a field survey for the census of animal herds, the use of data from different livestock services and the formulas have enabled us to make an estimate of methane emissions from the sector. livestock in the region. The results obtained during this work relate to: the average daily quantities of cow dung and chicken droppings in the Kankan region are: cow dung (4.45 kg) and chicken droppings (0.015 kg) ; the number of animal livestock by prefecture of the region, i.e. a bovine population of the region (1572132); the daily regional potential of biogas and energy respectively (763588.006 m<sup>3</sup>/d) and (3497233.069 kWh/d) and annual regional methane emissions (108908054 kg/year). The results obtained during this work are a first estimate of methane emissions due to the livestock sector in the Kankan region.

*Keywords: Assessment, potential, emissions, methane, breeding*

## 1. INTRODUCTION

The study of the causes and effects of climate change has given rise to a large number of scientific works in various fields. A key point is now the study of the coupling between human activities and climate change, in particular in order to limit the harmful impacts for our societies [1]. In 2013, the Intergovernmental Panel on Climate Change (IPCC) highlighted the effect of human activities on the increase in greenhouse gas (GHG) concentrations and therefore average surface temperatures (the temperature at the surface of the Earth would have increased by 0.85°C since 1850). Since the beginning of the industrial era, atmospheric concentrations of nitrous oxide (N<sub>2</sub>O), carbon dioxide (CO<sub>2</sub>) and methane (CH<sub>4</sub>) have increased by 15%, 30% and 150% respectively [2].

Faced with this environmental threat, the developed countries agreed in 1997 to establish a plan to reduce greenhouse gas (GHG) emissions, this is the Kyoto Protocol. The report of the Intergovernmental Panel on Climate Change (IPCC) confirms and reinforces the certainty of the existence of an increase in the greenhouse effect due to human activities [3]. The consequences of climate change due to GHGs are multiple, we can cite: the melting and disappearance of glaciers, the reduction of water resources, the health consequences and the rise in sea level [4, 5].

Like any human activity, livestock farming exerts a strong influence on the natural environment that surrounds it, this impact can be positive and/or negative for the ecosystems involved by the provision of benefits and pressures on the environment by emissions of GHGs throughout the chain. There are three main GHGs emitted in the context of livestock-related activities, which are: carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>) and nitrated nitrogen (N<sub>2</sub>O). About 44% of GHG emissions from the livestock sector are made up of CH<sub>4</sub> [6, 7].

In Guinea, livestock farming is the second most important activity in the rural world. It is a growth-enhancing sector that contributes substantially to food security and the fight against poverty. The national livestock census in 2017 gave: cattle (6407000); sheep and goats (459400). The poultry population is estimated at 28400000 poultry of local varieties and 1500000 hens of improved strains in semi-intensive poultry farms [8]. Livestock are an important source of CH<sub>4</sub> emissions due to their numbers, their ruminant digestive systems, in which enteric fermentation takes place and the management of their excreta (dung, manure, slurry, etc.).

The global anthropogenic emission (63%) of the total CH<sub>4</sub> emissions comes mainly from swamps, ruminant breeding, rice cultivation, household waste dumps, oil and gas operations [9]. CH<sub>4</sub> emissions from manure management are generally less significant than enteric emissions. The largest emissions

are associated with animal management operations in confined spaces, where manure is treated using liquid systems. Two-thirds of the CH<sub>4</sub> from ruminant farming comes from enteric fermentation and one-third from animal waste. The average annual values of potential CH<sub>4</sub> emissions due to livestock are: dairy cows (90 kg/year), growing cattle (65 kg/year) and cattle from 2 to 6 years old (51 kg/year) [10,11].

This study falls within this perspective. Its objective is to contribute to the assessment of methane emissions due to the livestock sector in the administrative region of Kankan.

## 2. MATERIAL AND METHOD

### 2.1. Presentation of the study area

The administrative region of Kankan is located 781 km from the capital Conakry. It is the largest administrative region of Guinea, it covers an area of 72145 km<sup>2</sup>, with five prefectures (Kankan, Kérouané, Kouroussa, Mandiana and Siguiré). The sub-Saharan climate is characterized by the alternation of two seasons (dry and rainy) with temperatures varying from 25°C to 41°C and rainfall varying between 1100 and 1800 mm of water per year [12]. The population of the Kankan region is estimated at 2 097 257 inhabitants in 2016, with an average density of 28 inhabitants per km<sup>2</sup>. The prefecture of Siguiré is the most populated in the region and in Guinea, with 724 631 inhabitants, including 360 147 women (49.70%) [13].

### 2.2. Process of methane emissions due to livestock

CH<sub>4</sub> emissions from manure management are generally less significant than enteric emissions. The largest emissions are associated with animal management operations in confined spaces, where manure is treated using liquid systems. Two-thirds of the CH<sub>4</sub> from ruminant farming comes from enteric fermentation and one-third from animal waste [14]. In a suckler cattle farm, enteric methane is responsible on average for 59% of greenhouse gas emissions [15]. Fig. 1 shows the process of enteric methane formation.




Fig.1. Formation of enteric methane [15]

The management of manure from breeding sites to pastures through storage and spreading is the most impactful item of the livestock operation from an environmental point of view. These effluents are responsible for non-enteric CH<sub>4</sub> emissions.

The method for estimating CH<sub>4</sub> emissions from livestock consisted of categorizing livestock, assessing populations and knowing the characteristics of the diet. The equations are applied seasonally to assess livestock feed consumption, while taking into account certain performance parameters of the different animal categories, which are: average weight, average weight gain per day, average mature weight, average number of working hours per day, feeding conditions, average temperature of the rearing environment, average daily milk production, and feed digestibility rate [16].

Gross energy (GE) is the energy needed by an animal to survive and support activities such as growth, lactation and gestation. This energy consumption for cattle is calculated by relation 1 [11].

$$EB = \left[ \frac{(EN_s + EN_a + EN_l + EN_{tra} + EN_g)}{TES} + \left( \frac{EN_{cce}}{TEC} \right) \right] \frac{DA\%}{100} \quad (1)$$

With :

EN<sub>s</sub>: Net energy necessary for survival in MJ/d; NE<sub>a</sub>: Net energy necessary for the activities in MJ/d; EN<sub>l</sub>: Net energy required for lactation in MJ/d; EN<sub>tra</sub>: Net energy required for work in MJ/d; EN<sub>g</sub>: Net energy required for gestation in MJ/d; EN<sub>cce</sub>: Net energy required for growth in MJ/d; TES: Rate of net energy available in food for survival in %; TEC: Net energy rate in the feed available for growth in %; DA%: Digestible energy expressed as a percentage of gross energy. These different parameters are defined as follows [17, 18].

*The Net Energy Needed for Survival* is the amount of energy needed to keep the animal in balance (the body neither loses nor gains energy). It is determined by relation 2.

$$EN_s = C_f_i (PV)^{0,75} \quad (2)$$

PV: Average live weight of the animal in (kg); C<sub>f<sub>i</sub></sub>: Coefficient varying for all animal categories in MJ/d.kg (C<sub>f<sub>i</sub></sub> = 0.322 for cows outside the lactation period, C<sub>f<sub>i</sub></sub> = 0.386 and C<sub>f<sub>i</sub></sub> = 0.370 for bulls).

*Net energy needed for activities* is the energy that animals need for food, drink and shelter. It is based on dietary conditions rather than the characteristics of the diet itself. This energy is calculated by relation 3.

$$EN_a = C_a \cdot EN_s \quad (3)$$

With: C<sub>a</sub> the activity coefficient, corresponding to the animal's food conditions (C<sub>a</sub> = 0.36 for large open pastures).

The Net energy needed for growth is the energy of weight gain. It is determined by the relationship.

$$EN_{c_{ce}} = 22,02 \cdot \left( \frac{PV}{C.P.M} \right)^{0,75} PP^{1,097} \quad (4)$$

Or :

C: Coefficient equal to 0.8 for females; 1.0 for castrates and 1.2 for bulls;  
MP: Mature live weight of adult female of moderate body condition in kg;  
PP: Average weight gain per day in kg/d.

*The net energy needed for lactation* in cattle is expressed as a function of the quantity of milk produced and the fat content in % equal to 4%. It is determined by relation 5.

$$EN_l = \text{lait} \cdot (1,47 + 0,40 \cdot \text{mat\_gra}) \quad (5)$$

Milk: Quantity of milk produced in kg/d,  
Mat<sub>gra</sub>: Milk fat content in % of weight.

*The Net Energy Needed for Work* is the energy needed to pull the cattle. This net energy necessary for work is determined by relation 6.

$$EN_{tra} = 0,10 \cdot EN_s \cdot t \quad (6)$$

Where: t is the average number of working hours per day.

Net energy needed for gestation, the energy needed for gestation lasting an average of 281 days per year, is estimated at 10% of NEs. Relation 7 makes it possible to evaluate this energy.

$$EN_g = C_g \cdot EN_s \quad (7)$$

Where: C<sub>g</sub> is the gestation constant for cattle, it is 0.10.

*The rate of net energy available in food for survival* in relation to the digestible energy consumed (TES) is calculated by relation 8.

$$TES = [1,123 - (4,092 \cdot 10^{-3} \cdot DA\%) + [1,126 \cdot 10^{-5} \cdot (DA\%)^2] - \left( \frac{25,4}{DA\%} \right)] \quad (8)$$

*The rate of net energy* in the feed available for growth in relation to the digestible energy consumed (TEC) is calculated by relation 9.

$$TEC = [1,164 - (5,160 \cdot 10^{-3} \cdot DA\%) + [1,308 \cdot 10^{-5} \cdot (DA\%)^2] - \left( \frac{37,4}{DA\%} \right)] \quad (9)$$

Feed consumption in units of dry matter (DMC) per day (kg/d), is estimated by dividing the BE by the energy density of the feed. A default value of 18.45 MJ/kg dry matter can be used if specific feed information is not available. The CMS value should be in the range of 2 to 3% of the body weight of mature or growing animals. For high-producing dairy cows, consumption may exceed 4% of body weight [19]. Relations 10 and 11 allow us to evaluate this consumption.

For growing and end-of-life cattle, we have:

$$CMS = PV^{0,75} \cdot \left[ \frac{(0,2444 \cdot EN_{ma} - 0,0111 \cdot EN_{ma}^2 - 0,472)}{EN_{ma}} \right] \quad (10)$$

For mature cattle we have:

$$CMS = PV^{0,75} \cdot \left[ \frac{(0,0119 \cdot NE_{ma}^2 + 0,0119)}{EN_{ma}} \right] \quad (11)$$

Where: EN<sub>ma</sub> is the estimated net dietary energy concentration of the diet.

### **2.2.1. Potential for CH<sub>4</sub> emissions from enteric fermentation**

Herbivores produce methane as a byproduct of enteric fermentation. The amount of methane emitted depends on the type of digestive tract, the age, the weight of the animal and the quality and quantity of food consumed. The potential for CH<sub>4</sub> emissions due to enteric fermentation is calculated by relationships 12 and 13.

$$Em_{CH_4} = FE_{(T)} \cdot \left( \frac{N_{(T)}}{10^6} \right) \quad (12)$$

$$TEm_{CH_4} = \sum_i E_i \quad (13)$$

Or :

TE<sub>mCH<sub>4</sub></sub>: Annual methane emissions due to enteric fermentation in (Gg CH<sub>4</sub>/year);

EF<sub>(T)</sub> : Emission factor for the livestock category defined in (kg CH<sub>4</sub>/head year);

N<sub>(T)</sub> : Number of heads of livestock species/category T; T: Livestock species/category;

E<sub>i</sub> : Emissions of livestock category and sub-category i.

The EF emission factor for each animal category is given by relationship 14.

$$FE = \left[ \frac{EB \cdot \left( \frac{Y_m}{100} \right) \cdot 365}{55,65} \right] \quad (14)$$

Or :

EB: Gross energy consumption in MJ/head.day;

Y<sub>m</sub>: CH<sub>4</sub> conversion factor, this is the percentage of gross energy in the feed converted to CH<sub>4</sub> and the value 55.65 (MJ/kg CH<sub>4</sub>) represents the value

### **2.2.2. Potential for CH<sub>4</sub> emissions from manure management**

The decomposition of manure under anaerobic conditions, during storage, processing and application to pasture produces CH<sub>4</sub>. The different steps used to assess non-enteric or manure management CH<sub>4</sub> emissions consist of:

- Gather data corresponding to livestock populations based on their characteristics;
- Use default values or develop country-specific emission factors for all livestock subgroups in terms of kg CH<sub>4</sub>/animal/year;
- Multiply the emission factors of the livestock subgroup by the population of the subgroup;
- Add emissions from all defined livestock species to determine national emissions.

Relationship 15 is used to evaluate CH<sub>4</sub> emissions due to manure management for a defined population, in Gg CH<sub>4</sub>/year.

$$CH_{4fumer} = \sum_{(T)} \frac{(FE_{(T)} \cdot N_{(T)})}{10^6} \quad (15)$$

Or :

EF<sub>(T)</sub>: Emission factor for the livestock category defined in kg CH<sub>4</sub>/head year;

N<sub>(T)</sub>: Number of heads of livestock species/category T in the country;

T: Livestock species/category.

The value of the MCF of a system varies according to the way of managing the manure and the climate, it varies from 0 to 100%. The evaluation of the FCM is thus done using relation 16.

$$FE_{(T)} = (SV_{(T)} \cdot 365) \cdot \left[ B_{O(T)} \cdot 0,67 \text{kg/m}^3 \cdot \sum_{S,K} \frac{FCM_{S,K}}{100} \cdot GF_{(T,S,K)} \right] \quad (16)$$



1	Kankan	23281	15228265	105000	5250	113834	910672	16144187
2	Kouroussa	294137	19 118 905	794	39.7	174614	1396912	20515857
3	Mandiana	326462	21 220 030	2 500	1025	225247	1801976	23023031
4	Siguiri	372411	24 206 715	100000	5000	195151	1561208	25772923
5	Kérouané	344841	22 414 665	95350	4767.5	129078	1032624	23452057
TOTAL		1572132	102188580	321644	16082.2	837924	6703392	108908054




Fig. 3. Number of livestock by prefecture





Fig. 4. Potential for CH4 emissions from the livestock sector by species



**Fig. 5. Potential annual CH<sub>4</sub> emissions due to the livestock sector by prefecture**

The diagrams in fig. 3, 4 and 5 show the potential annual CH<sub>4</sub> emissions due to the livestock sector by prefecture in the region. The results obtained show that the greatest quantity of emissions is recorded in Siguiiri (25772923 kg/year), followed respectively by Kérouané (23452057 kg/year), Mandiana (23023031 kg/year), Kouroussa (20515857 kg/year), and Kankan (16144187 kg/year). With total annual emissions in the region of 108908054 kg/year. This result is a first estimate of the contribution to greenhouse gases of the livestock sector in the Kankan region.

#### 4. CONCLUSION

This study consisted in carrying out tests to assess the potential for methane emissions due to the livestock sector in the Administrative Region of Kankan. The results of this work show that the greatest amount of CH<sub>4</sub> emissions is recorded in Siguiiri, followed respectively by Kérouané, Mandiana, Kouroussa, and Kankan. Thus, the management and recovery of this animal waste would make it possible to locally reduce methane emissions.

#### REFERENCES

- [1] DAMON Gildas David Farid, (2017). Modélisation de la cinétique de gazéifications étagée de la biomasse tropicale : cas des balles de riz et des rafles de maïs. Thèse de doctorat de l'université d'Abomey Calavi du Bénin et de l'Université de Technologie de Compiègne, 174p.
- [2] HOUGHTON J.T., CALLANDER B.A., VARNEY S.K. EDS., (1992). Climate change. The supplementary report to the IPCC scientific assessment, IPCC Scientific Assessment Working Group. Cambridge University Press, Cambridge, 200 p.
- [3] GIEC (2001). Climate Change, (2001) : Impacts, adaptation and vulnerability. Contribution of Working Group II to the third assessment report of IPCC. Cambridge University Press, Cambridge, 850 p.
- [4] Andrew V, Hambaliou B., Anna C., Robert J. G., Martin N., Claudia W. R., Ray D., Doug M. Potential Methane Emission Reductions for Two Manure Treatment Technologies. Environmental Technology 2017; 23.
- [5] Browne J. D., Allen E., Murphy J. D. Evaluation of the biomethane potential from multiple waste streams for a proposed community scale anaerobic digester. Environmental Technology, 2013
- [6] GIEC. (2007). Climate Change 2007 : Mitigation. Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. B. Metz, O.R. Davidson, P.R. Bosch, R. Dave & L.A. Meyer, eds. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- [7] FAO. (2006). Livestock's long shadow - Environmental issues and options, by H. Steinfeld, P. J. Gerber, T. Wassenaar, V. Castel, M. Rosales & C. de Haan. Rome.
- [8] Ansoumane Sakouvogui, Madeleine Kamano, Mamby Keita, Assessment of the energy potential of pig dung by the production of biogas in the urban municipality from N'Zérékoré in Guinea. International Journal of Multidisciplinary Research and Growth Evaluation, Volume 2; Issue 4; July-August 2021; page No. 374-376.
- [9] Popova M., Morgavi D.P., Doreau M., Martin C., (2011). Production de méthane et interactions microbiennes dans le rumen. INRA Prod. Anim. 5, pp. 447 - 460.

- [10] Bodjui O. A., Loissi K., et Moussa B. (2017). Evaluation of the Biogas Production Potential by Anaerobic Digestion of Fermentable Agricultural Residues in Côte d'Ivoire. *International Journal of Waste Resources*, 7(4).
- [11] G.E.I.C. : Groupes d'Experts Intergouvernemental sur l'Evolution du Climat. Rapport disponible sur : <https://www.ipcc.ch/>
- [12] Rapport, Programme Conjoint des Nations Unies pour la région administrative de Kankan, avril 2013. 44p.
- [13] Faya Oulare, Ansoumane Sakouvogui, Sékou Fatoumata Conde, Mamby Keita, Study of the construction and testing of a storage collector-type solar water heater at Julius Nyerere University in Kankan, Guinea, *International Journal of Advanced Engineering and Technology* ISSN: 2456-7655, Volume 5, Issue 1, 2021; Page No. 16-21.
- [14] Bernard Seguin, Jean-François Soussan (2008)., Emissions de gaz à effet de serre et changement climatique : causes et conséquences observées pour l'agriculture et l'élevage. *Courrier de l'environnement de l'INRA* n°55, 13p.
- [15] Tubiello F., Salvatore M., Córdor Golec R.D., Ferrara A., Rossi S., Biancalani R., Federici S., Jacobs H., Flammini A., (2014). Agriculture, Forestry and Other Land Use Emissions by Sources and Removals by Sinks. ESS Working, Paper No. 2, FAO, Rome.
- [16] Dolle J.-B., Moreau S., Brocas C., Gac A., Raynal J., Duclos, A., (2015). Elevage de ruminants et changement climatique. Paris. Ed. Institut de l'Elevage. 24 pages.
- [17] Thomas TURINI., (2016). Les impacts environnementaux de la production de viande : Cas des ruminants. Journées Nationales Groupements Techniques Vétérinaires - Nantes 9 p.
- [18] Gac A., Béline F. Bioteau T., Maguet K., 2007. A French Inventory of Gaseous Emissions (CH<sub>4</sub>, N<sub>2</sub>O, NH<sub>3</sub>) from Livestock Manure Management Using a Mass-Flow Approach. *Livestock Science* 112, 252-260.
- [19] Gibbs, M.J., Conneely, D., Johnson, D., Lassey, K.R. and Ulyatt, M.J. (2002). CH<sub>4</sub> emissions from enteric fermentation. In : Background Papers: IPCC Expert Meetings on Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories, p 297–320. IPCC-NGGIP, Institute for Global Environmental Strategies (IGES), Hayama, Kanagawa, Japan.
- [20] Ulyatt, M.J., Lassey, K.R., Shelton, I.D. and Walker, C.F. (2005). Methane emission from sheep grazing four pastures in late summer in New Zealand. *New Zealand Journal Agricultural Research* 48 : 385-390
- [21] Yvan Chouinard, 65ème Congrès de l'Ordre de Agronomes du Québec, Production et émission du Méthane et du Gaz Carbonique par les ruminants, Université Laval, septembre 2016, 10 p.
- [22] R., Burch, and P.K. Loader, Investigation of Pt/Al<sub>2</sub>O<sub>3</sub> and Pd/Al<sub>2</sub>O<sub>3</sub> catalysts for the combustion of methane at low concentrations. *Applied Catalysis B : Environmental*, 2010, pp. 149-164.

UNDER PEER REVIEW