

FACTORS ASSOCIATED WITH SURGICAL TREATMENT OUTCOME OF PSOAS MUSCLE ABSCESS

ABSTRACT

Background: Psoas abscess is a rare condition but one with potential for severe morbidity. Many factors may be implicated in the treatment outcome of the condition and efforts must be directed to identify, understand and control the factors for a better prognosis.

Objective: To identify the factors associated with surgical treatment outcome of psoas muscle abscess.

Methodology: This descriptive cross-sectional analysis was conducted upon a sample of 58 patients (chosen via non-probability, consecutive sampling) scheduled for surgical treatment for psoas muscle abscess (aged 18 to 60 years) at Dept. of General Surgery at Liaquat University Hospital, Hyderabad & Jamshoro. The data was recorded onto a pre-structured questionnaire comprising of inquiries pertaining to basic biodata, socio-demographic details, presenting complaints, hematological parameters, site, type, etiology and microbiology of the abscess and treatment outcome in terms of resolution of abscess cavity analyzed via SPSS v.16.0.

Results: Most common presentation of patients was pain in flanks, followed by fever. Inflated WBC count and high acute phase response was noted among most patients; with C-reactive protein and ESR levels amounting to twice the normal range in all patients. The abscess was commonly right sided (81%) and secondary (82.7%) in origin. The commonest isolated organism was Mycobacterium TB and consequently spinal TB was the commonest etiological factor. Resolution of abscess cavity was noted among 77.5% of the cases with a greater proportion of recovery falling in line non-tuberculous etiology, less altered hematological indices and primary psoas muscle abscess.

Conclusions: After careful consideration, it can be concluded that secondary psoas muscle and that with tuberculous etiology have a poorer surgical outcome, thus greater care and more effective surgical techniques may be employed to achieve complete resolution of abscess.

Keywords: Mycobacterium TB, Psoas Abscess, Etiology, Treatment Outcome and Secondary Abscess.

INTRODUCTION

Psoas abscess is a bothersome condition with unspecific presentation and insidious onset, resulting in diagnostic and treatment delays and resultantly, a high level of morbidity i.e. destruction of at least a part of the psoas muscle. ^[1, 2] Initially believed to be rare, the condition is on the rise and especially due to HIV, I/V drug use, and immunosuppressant therapy; the incidence had risen to cases per 10,000 hospital admissions in the last decade. ^[3] Studies as early as 1992 have reported a 400% increase in prevalence every 5 years. These studies also report that epidemiological characteristics vary depending on the underlying cause. Primary psoas abscesses account for approximately 30% of cases worldwide. More than 90% occur in developing or tropical countries. 83% of primary cases occur in patients < 30 years. Secondary psoas abscesses are more common, particularly in developed countries and affect the elderly more often. ^[4-10] Through the surgical management (drainage) is simple and often successful, complications such as avascular necrosis of the femoral head, osteomyelitis, cellulitis of the thigh, and septic arthritis of the hip may arise. Other complications may include iliac vein thrombosis, pulmonary embolism,

hydro-nephrosis, renal failure and further dissemination of organisms especially in immunocompromised patients. Many factors may be implicated in the treatment outcome of the condition and efforts must be directed to identify, understand and control the factors for a better prognosis. [11-13]

MATERIAL AND METHODS

This descriptive – cross-sectional analysis was conducted upon a sample of 58 patients (chosen via non-probability, consecutive sampling) scheduled for surgical treatment for psoas muscle abscess (aged 18 to 60 years) at Dept. of General Surgery at Liaquat University Hospital, Hyderabad & Jamshoro. Consenting adult patients (of both genders) aged 18 to 60 years presenting with psoas muscle abscess (more than 5 cm) were included in the study. Non-consenting patients and patients suffering from with severe co- morbidities like uncontrolled diabetes mellitus, uncontrolled hypertension chronic, HCV and HBV were excluded from the sample. The data was recorded onto a pre-structured questionnaire comprising of inquiries pertaining to basic biodata, socio-demographic details, presenting complaints, hematological parameters, site, type, etiology and microbiology of the abscess and treatment outcome in terms of resolution of abscess cavity analyzed via SPSS v.16.0.

RESULTS

Total of 58 patients of Psoas muscle abscess were evaluated. The mean age of study subjects were 38.5 ± 12.5 (range from 18-60 years).

In current study there were 72.4% (n=42) males 27.5% (n=16) females

In this study most common presentation of patients was pain in all 100% (n=58) subjects from which 50% (n=29) had pain in flanks, 34.4% (n=20) in groin and 15.5% (n=9) in abdomen, followed by fever 72.4% (n=42). Other symptoms included poor health condition 70.6% (n=41), Palpable mass in 43.1% (n=25) from which 22.4% (n=13) was Groin lump and 20.6% (n=12) was Abdominal lump, limitation of hip movement seen in 20.6% (n=12).

Table 1: The factors affected treatment outcome as shown below.

Outcome		Gender		Age	
		M (42)	F (16)	< 35 (21)	≥ 35 (33)
Drainage in 1 st Attempt	Complete	34	11	19	26
	Incomplete	08	05	02	11
Mean Resolution Time (Days)		8.42	12.11	7.02	14.83
Pain Relief	1 st POD	13	02	08	06
	2 nd POD	21	04	05	07
	3 rd POD	06	08	04	12
	At Discharge	02	02	04	08
Complete Resolution (Abscess Cavity)		32	13	18	27

Duration (Hospital Stay)	< 3days	29	14	15	28
	> 3 days	13	02	06	05

Outcome		Tuberculosis		Type	
		Yes (39)	No (19)	Primary (10)	Secondary (48)
Drainage in 1st Attempt	Complete	28	17	09	36
	Incomplete	11	02	01	12
Mean Resolution Time (Days)		13.71	6.40	7.81	14.92
Pain Relief	1st POD	12	11	09	15
	2nd POD	25	08	01	15
	3rd POD	02	00	00	13
	At Discharge	00	00	00	05
Complete Resolution (Abscess Cavity)		29	16	10	35
Duration (Hospital Stay)	< 3days	25	18	07	36
	> 3 days	14	01	03	12

DISCUSSION

In present study a total of 58 patients of Psoas muscle abscess were evaluated. The mean age of study subjects were 38.5 ± 12.5 (range from 18-60 years). Results were comparable with A study from India by Dave BR et al the mean age was 36.5 ± 12.7 (range 18-63 years). Whereas F. Pombo et al has given mean age 35 years (ranged from 19 to 61 years), Elbadrawi AM et al, 32 (range 21 to 55), Aboobakar R, et al, 32 years (range from 10-70 years), Ye F. et al. 38.5 ± 8.7 years (age range from 20 to 63 years). [3, 5, 7, 8, 11]

In current study there were 72.4% (n=42) males 27.5% (n=16) females. Results were comparable with a study from India by Dave BR et al, who has reported in his study that there were 21(72.4%) males and 8 (27.5%) females. Whereas Wong et al has reported 27 (64.2%) males and 15 (35.7%) females, Yadav RP et al, 22 (61.1%) males and 14 (38.9%) females, Y.J Kim et al 63 (54.3%) were male and 53 (45.3%) female. Another study from turkey by Tarhan H, has reported 6 (40%) patients women and 9 (60%) men. Alvi AR, has giving in his study 4 (66.6%) male and 2 (33.3%) female, whereas Tabrizian P, has reported 32 (52.4%) men and 29 (47.5%) women. [1-3, 6, 10, 12, 13]

In this study most common presentation of patients was pain in all 100% (n=58) subjects from which 50% (n=29) had pain in flanks, 34.4% (n=20) in groin and 15.5% (n=9) in abdomen,

followed by fever 72.4% (n=42). Other symptoms included poor health condition 70.6% (n=41), Palpable mass in 43.1% (n=25) from which 22.4% (n=13) was Groin lump and 20.6% (n=12) was Abdominal lump, limitation of hip movement seen in 20.6% (n=12). Tabrizian P reporting in his study that non-specific symptoms were found in most patients. The most common initial symptoms, including abdominal pain, other gastrointestinal tract complaints, and lower extremity pain.

Only 26% (16 of 61) of patients were initially seen with fever (38.5°C), Dave B R et al, reporting in his study Presenting features included back pain (n = 29), radicular pain (n = 6), fever (n = 8), weight loss (n = 15), anorexia (n = 10), walking difficulty (n = 15), spasm (n = 29), range of movement restriction (n = 29), groin mass (n = 5), and pseudo flexion deformity of hip (n = 8). Aboobakar R, et al, also reporting the commonest presenting symptom was unilateral flank pain (50%) and the commonest sign was unilateral flank tenderness (70%). [3, 5, 13]

CONCLUSION

After careful consideration, it can be concluded that secondary psoas muscle and that with tuberculous etiology have a poorer surgical outcome, thus greater care and more effective surgical techniques may be employed to achieve complete resolution of abscess.

REFERENCES

1. Kim Y.J, Yoon J.H, Kim S.I, Wie S.H, Kim Y.R. Etiology and outcome of iliopsoas muscle abscess in Korea; changes over a decade. *Intern J Surg* 2013;11(2) :105-9.
2. Wong, PL Ho, SK Lam. Retrospective review of clinical presentations, microbiology, and outcomes of patients with psoas abscess. *Hong Kong Med J* 2013;19(2):416-23 .
3. Dave B R, Kurupati RB, Shah D, Degulamadi D, Borgohain N, Krishnan A et al. Outcome of percutaneous continuous drainage of psoas abscess: A clinically guided technique. *Indian J Orthop.* 2014;48(1): 67–73.
4. Hsieh MS, Huang SC, Loh EW, Tsai C-A, Hung Y-Y, Tsan YT et al. Features and treatment modality of iliopsoas abscess and its outcome: a 6year hospital-based study. *BMC Infectious Diseases* 2013; 13(1):578.
5. Aboobakar R , Cheddie S , Singh B. Surgical management of psoas abscess in the Human Immunodeficiency Virus era, *Asian Journal of Surgery* 2016;10(4):1-5.
6. Tarhan H, Çakmak Ö, Türk H, Can E, Un S, Zorlu F. Psoas Abscess: Evaluation of 15 Cases and Review of the Literature. *Journal of Urological Surgery* 2014;1: 32-35-73
7. Pombo F, Martín-Egaña R, Cela A, Díaz J, Linares-Mondéjar P, Freire M. Percutaneous Catheter Drainage of Tuberculous Psoas Abscesses, *Acta Radiologica*, 1993;34(4): 366-68
8. Elbadrawi AM, Morsi AM, Elkhateeb TM. Drainage of Pyogenic Sacro-Iliac Joint Infection Using a Percutaneous Technique (Case Series of 13 Patients). *MOJ Orthop Rheumatol.* 2017; 8(2):3-6

9. Rafiq K, Iqbal M, Chodhry AM. Psoas abscess (Presentation, asseesment and management) *Annals*. 2002; 8(1):19-21
10. Yadav RP, Agrawal CS, Adhikary S, Kumar M, Regmi R, Amatya R et al. Iliopsoas abscess: Analysis and perspectives from an endemic region of Eastern Nepal. *Kath Uni Med J*. 2007; 5(4): 497-500
11. Ye F, Zhou Q, Feng D. Drainage of tuberculous psoas abscess. *Med Sci Monit*, 2017; 23(2): 5374-81
12. Alvi AR, Rehman ZU, G Nabi ZR. Pyogenic psoas abscess: case series and literature review. *Tropical Doctor* 2010; 40: 56–58
13. Tabrizian P, Scott QN, Greenstein A, singh UR, Celia M, Divino et al. Management and Treatment of Iliopsoas Abscess. *Arch Surg*. 2009;144(10):946-49

UNDER PEER REVIEW