

Case Report on Alcohol Dependence WITH DELIRIUM TREMORS

Abstract

Production :-

Alcohol is a natural product made by reacting lactose fermentation with yeast spores. Because it includes calories with no nutritional value, alcohol is used as a source of energy. Different sugar sources are used for fermentation to create a variety of alcoholic beverages.

Patient Information

Patient Present History:-

On September 3rd, 2019, a 24-year-

old male patient was admitted to AVBRH with a specific complaint of abdominal pain for the past three years. Mr. Suraj began drinking alcohol on a daily basis. Initially, he only drank at night, but later he began drinking during the day.

Clinical finding :-

Mood irritable, social withdrawal, mental confusion, seizure, and vomiting

Diagnostic evaluation:-

Hb - 11.9mg%, MCV - 87cum/micron, MCH- 28.9pico/gm, Total Bilirubin- 1.47

USG- His appearance was natural on cement. There was no localised lesion or obvious mass lesion in the pelvis.

KEYWORDS:-

Alcohol Dependence Syndrome with Delirium Tremor

INTRODUCTION

Alcohol is a natural product made by reacting lactose fermentation with yeast spores. Because it includes calories with no nutritional value, alcohol is used as a source of energy. Different sugar sources are used for fermentation to create a variety of alcoholic beverages. History of the Patient - On May 5, 2021, a 32-year-old male patient was admitted to AVBRH with the primary complaint of. Since the age of two

A previous psychiatric illness (DSM - IV and ICD - 10) in which a person is physically or mentally reliant on alcohol is known as alcohol dependency. A chronic condition in which a person desires alcoholic beverages and is unable to control his or her consumption. A person with this disease often has to drink higher doses to achieve the name effect and may have symptoms

KEY WORDS :-

Patient Information

Patient Present History :-

On September 3rd, 2019, a 24-year-old male patient was admitted to AVBRH with a specific complaint of abdominal pain for the past three years. Mr. Suraj began drinking alcohol on a daily basis. Initially, he only drank at night, but later he began drinking during the day. On a daily basis, he consumes 2 to 3 quarters. Eye opener drinking and abdominal discomfort are also reported by the patient. Intermittently, he is in excruciating discomfort.

| Content | Results |
|--------------------------------|-------------------------------------|
| Mood and affect is impaired | Unstable and irritable mood |
| Content of thought is impaired | Delusions of persecution is present |
| Attention is intact | he count the number backwards |
| Concentration is intact | correct result |
| Judgment is intact | he giving correct answer |

Table 1. Mental Status Examinations

| Pathology | Results |
|-----------------|----------------|
| Hb % | 11.9mg % |
| MCV | 87Cum / micron |
| MCH | 28.9Pico / gm |
| Total Bilirubin | 1.47 |

Table 2. Blood Investigation Report

USG - His appearance was natural once more. There was no localised lesion or obvious mass lesion in the pelvis.

The first impression is mild hepatomegaly.

A thorough examination of the patient's background and mental state indicates that he or she is truly suffering from a racial oppression hallucination.

The first impression is mild hepatomegaly.

A thorough examination of the patient's background and mental state indicates that he or she is truly suffering from a racial oppression hallucination.

Therapeutic intervention -

Psychopharmacology

Antipsychotic and anticraving medications were given to the patient. Tab Neurobiantfort 2ml OD, inj. Lorazepam 2mg HS, inj. Thiamine 200mgIM

Treatment for a short period of time

Acute psychotic disorders necessitate prompt hospitalisation, whether in a mental facility or a crisis line. These syndromes are used to treat a variety of medical conditions. The attending physician makes the choice to admit a patient to the hospital.

Continuation Treatment- *After a specific period of time has elapsed, the effectiveness of anti-craving medicine for psycho-pharmacotherapy is usually evident. Improved sleep, agitation regression, anxiety and paranoia reintegration, and the removal of psychotic symptoms were all seen in the first six weeks. If the patient's condition does not improve, another antipsychotic medication will be prescribed, or the first dose will be raised. A worsening of symptoms, severe side effects, or an insufficient pharmaceutical response are all common reasons for cognitive treatment.* Treatment with opioids, mood stabilisers (lithium or valproate), or an anticonvulsant medicine (carbamazepine) may be prescribed if mood disorders or cyclic breakouts emerge. The distinction between post-narcoleptic depression and the development of an affective illness is crucial (schizoids). Prevention of recurrence Consider the likelihood that symptoms will resurface during the first two years of starting treatment, which should be continued for another one to two years after recovery. Periodic checkups and effective therapeutic therapy, including social and psychological counselling, are necessary during this long-term follow-up

Nursing management :-

Nursing care was provided according to the requirements in order to maintain personal hygiene and avoid further complications.

The patient, as well as his relatives, should be elicited as much information as feasible.

The behaviour pattern included posturing, psychomotor activity, and hygiene.

Check to make sure the person is not hurt.

Take notice of the patient's emotional state and affect to see if it's appropriate.

Assessed to discover if his thoughts were delusory.

Determine the severity of the patient's hand tremor and suggest upper-extremity training.

Follow-up and Results.

At the time of release, the patient's condition was satisfactory. His main complaint when he was admitted to the hospital was hand tremor, abdominal pain, nausea, vomiting, and headache. His condition has improved since he received treatment. Their family was informed about the medication regimen and personal hygiene instructions, and they were told to return in 10 days for a routine examination to see how the sickness was progressing.

Discussion

Syndrome of Alcohol Dependence The Edwards and Gross alcohol - dependence syndrome (ADS) formulations from 1976 are included in the main classification systems for alcohol - use disorder (DSM - IV and ICD - 10). However, the criteria for distinguishing alcohol dependence (AD) from substance dependency (AA) in DSM - IV and hazardous use in ICD - 10 continue to be problematic. Furthermore, these worries stem from a shift in historical perspectives and problems associated with alcohol intake. Alcohol use disorders (AUDs) are characterised in terms of their historical evolution; improvements in the recognition of drug issues including alcohol. ⁽⁴⁾

The amount of alcohol that can be detected is different. The degree to which alcohol is linked to such events varies. The Stroke fraction is influenced by alcohol, coronary heart disease, liver cirrhosis, and a variety of cancers. The amount of alcohol that causes various disorders varies. (7) Alcohol consumption is the fourth leading cause of preventable death in the United States (after smoking, high blood pressure and obesity). Hazardous alcohol use killed nearly 3 million people in 2016, accounting for 5.3 percent of all deaths, according to a 2018 WHO estimate. (8) Men are responsible for the majority of such deaths over the world. Ghogare had conducted study on the effects of alcoholism on the brain. In a study, Joshi et al compared total self-stigma in patients with schizophrenia and alcoholism. Kelkar and Parisha studied the prevalence of substance abuse among schizophrenia patients. (11) Patel et al looked on the frequency of alcohol-related disorders in hospitalised male patients. Clinical comorbidity, severity, and liver enzyme impairment in alcohol-dependent patients were reported by Gedam et al. (12). 13 In rural Central India, Jonas et al. investigated the prevalence of depression, suicidal ideation (13) and alcohol and nicotine consumption.

Strength

A 24-year-old male patient tolerates all medications well and responds effectively to the hospital's therapeutic therapy within seven days of receiving it.

Informed Consent

Patients and their relatives were notified before this case was adopted, and informed consent was obtained from both the patient and the relatives.

CONCLUSIONS

Patient The patient was admitted to the hospital for 10 days with abdominal pain, nausea, and vomiting as the chief diagnosis. Tremor might also be felt in both hands. A member of the health team had begun rapid therapy now that the patient's condition had improved.

REFERENCES

- [1] Coffelt, T.A., Bauer, B.D. and Carroll, A.E., 2013. Inpatient characteristics of the child admitted with chronic pain. *Pediatrics*, 132(2), pp.e422 - e429.
- [2] Freud, S. (1917). *Metapsychologie* Ergänzung Zur Traumlehre. In *Gesameltewerke*.
- [3] Fuechtner, V., 2011. *Berlin Psychoanalytic: Psychoanalysis and Culture in Weimar Republic Germany and Beyond* (Vol. 43). Univ of California Press.
- [4] Lanczik, M., 1992. Karl Ludwig Kahlbaum (1828 - 1899) and the emergence of psychopathological and nosological research in German psychiatry. *History of Psychiatry*, 3(9), pp.53 - 58.
- [5] Bleuler, E. (1911). *Dementia praecoxoder Gruppe der Schizophrenien*. In *Aschaffenburg*
- [6] Aschaffenburg, G., 1911. *Handbuch der Psychiatrie*: hrsg. von G. Aschaffenburg (Vol. 4, No. 1). Deuticke.
- [7] American Psychiatric Association (2004). *Practice guideline for the treatment of patients with schizophrenia*. Supplement to the *American Journal of Psychiatry*, 161.
- [8] Janicak, P., Davis, J., Preskorn, S., et al. (1997). *Management of acute psychosis*. In *Principles and practice of psychopharmacotherapy* (eds. P.G. Janicak, J.M. Davis, S.H. Preskorn, and F.J. Ayd) (2nd edn), pp. 110–39. Williams and Wilkins, Baltimore, MD.
- [9] Ting - Kai Li1, Brenda G Hewitt, the Alcohol Dependence Syndrome, 30 Years Later: A Commentary. *The 2006 H. David Archibald Lecture*, Epub 2007 Aug 4.
- [10] Alcohol - Use Disorders: Diagnosis, Assessment and Management of Harmful Drinking and Alcohol Dependence. <https://www.ncbi.nlm.nih.gov/books/NBK65500/>
- [11] Ghogare, Ajinkya Suresh Rao. "study of cognitive impairment in patients of alcohol use disorder." *Indian journal of psychiatry* 61, no. 9, 3 (January 2019): S583–84.
- [12] Joshi, Amey, Lipsy Modi, and Imran Ali Shivji. "Comparison of Total Self - Stigma between Schizophrenia and Alcohol Dependence Patients." *INDIAN JOURNAL OF PSYCHIATRY* 60, no. 5, 1 (February 2018): 125–26.
- [13] Kelkar, Parisha. "prevalence of substance abuse in patients of schizophrenia." *Indian journal of Psychiatry* 61, no. 9, 3 (January 2019): S472–73.
- [14] Patel, Tejas, V, Mehul J. Brahmabhatt, and Ganpat K. Vankar. "Prevalence of Alcohol Use Disorders in Hospitalised Male Patients." *Archives of Psychiatry and Psychotherapy* 20, no. 4 (December 2018): 47–55. <https://doi.org/10.12740/APP/99147>.

- [13] Gedam, Sachin Ratan, AjabDhabarde, Pradeep S. Patil, Animesh Sharma, Kanika Kumar, and Vijay Babar. "Psychiatric Comorbidity, Severity of Dependence and Liver Enzymes Dysfunction among Alcohol Dependent Individuals: A Cross - Sectional Study from Central Rural India." *Journal of Clinical and Diagnostic Research* 13, no. 4 (April 2019): VC1–5. <https://doi.org/10.7860/JCDR/2019/40368.12759>.
- [14] Jonas, Jost B., Vinay Nangia, Marcella Rietschel, Torsten Paul, Prakash Behere, and Songhomitra Panda - Jonas. "Prevalence of Depression, Suicidal Ideation, Alcohol Intake and Nicotine Consumption in Rural Central India. The Central India Eye and Medical Study." *PLOS ONE* 9, no. 11 (November 19, 2014). <https://doi.org/10.1371/journal.pone.0113550>.

UNDER PEER REVIEW