

EFFECTS OF COVID-19 ON CARDIOVASCULAR PATIENTS: A SYSTEMATIC REVIEW

ABSTRACT

Aim of Object-During the COVID-19 pandemic, the entire world is experiencing a mortality situation; most people are battling against the corona virus, but some individuals have already suffered from cardiovascular problems.

For improved patient care, adequate information and comprehension of the relationship between cardiovascular disorders and COVID-19 is required. The dominant clinical manifestations of the corona virus infection are on the respiratory system. In this instance, the acute cardiac injury is the most often reported cardiac abnormality, in which the degree of cardiac output is increased, troponin levels rise, and mostly it is found in about 8% to 12% of patients. The involvement of viral cardiomyocytes and systemic inflammation is the most prevalent mechanism for cardiac damage. The corona virus attaches itself and enters through angiotensin converting enzyme-II.

Discussion and Conclusion-Recent articles on COVID-19 have revealed nothing regarding these individuals' cardiac vascular manifestations. This is a critical component of all that has a big influence on COVID-19 patients' cardiovascular systems. To fully comprehend the method and effects, more study is required.

Keywords: COVID-19, Cardiovascular disease, Cardiac Troponins, Angiotensin Converting Enzyme-II (ACE-II).

1. Introduction

Corona viruses are a wide group of viruses that may infect both animals and humans. Several corona viruses have been linked to respiratory infections in humans, ranging from the common cold to more serious illnesses including Middle East Respiratory Syndrome (MERS) and Severe Acute Respiratory Syndrome (SARS) (SARS). This pandemic is challenging task for everyone especially healthcare sector around the world. Symptomatic phase of this virus is very dangerous but even in asymptomatic phase of Covid-19 has the ability to infect others and transmit in community at high infective rate which make the Covid -19 a world pandemic. Before the outbreak in Wuhan, China, in December 2019, no one had heard of this new virus or illness.^{1,2,3}

The signs and symptoms might range from a simple case of the flu to pneumonia. When infected individuals have non-communicable conditions such as cardiovascular disease or diabetes, these symptoms become much worse. If individuals are already having cardiovascular disease then they are more vulnerable to COVID 19 and worsen condition in the cardiovascular disease. The primary goal of this study is to offer an overview of various cardiovascular diseases and their manifestations in COVID 19 patients, as well as the influence of COVID 19 on patients with cardiovascular illness before and after treatment. Since scientists are continuously working to understand the new symptoms of COVID-19.⁴

2. Pathogenic considerations

The seven species of SARS CoV-2 (RNA beta corona virus) produce infection in humans and 4 species mainly produce flu like syndrome but the other 3 species (SARS, MERS and COVID-19) cause severe illness in humans. SARS CoV-2 primarily attack in the respiratory system but cardio vascular system also affected in several ways.⁵

Angiotensin-converting enzyme (ACE-2) is a key player in the neurohumoral control of the cardiovascular system, both in good health and in disease. SARS CoV-2 binds to ACE-2 receptors in the lungs and heart, altering the ACE-2 signalling pathway and causing pulmonary and cardiac issues.⁶ COVID-19 causes systemic inflammation and increase the level of cytokines which leads to multiple organ failure. In

studies, it has cleared that severe COVID-19 illness has high level of cytokines.⁷ Acute respiratory illness reduce the oxygen demand supply which cause acute myocardial injury.⁸

Systemic inflammation can create prothrombotic milieu and worsen the symptoms. Precipitation of plaque ruptures by systemic inflammation and increased shear stress due to increase coronary blood supply cause the acute myocardial infarction.⁹ COVID 19 is treated with a variety of antiviral medications, corticosteroids, and other treatments, although these have negative effects on the cardiovascular system. Electrolyte imbalance is common in cardio vascular disease patients and it can occur in any systemic condition, causing arrhythmias.¹⁰ Hypokalemia increases susceptibility to injury to various tachyarrhythmias.¹¹

3. Cardiovascular complications

COVID-19 cause myocardial injury and myocarditis with increases in troponin level which is due to hypoxia, Increased cardiac physiologic stress or direct myocardial injury.¹² In one study myocarditis was also recognizes with increase viral loads and related casualties were due to myocarditis.¹³ Acute Myocardial infarction (AMI) and atherosclerotic plaque disruption increases in systemic inflammation.¹⁴

It has been proved in patients with an initial stage of corona virus infection that 23% of AHF patients and 33% of cardiomyopathy patients were found among the COVID-19 positive cases.¹⁵ Another study discovered that heart failure was present in 24% of patients, and that it was associated to a higher risk of mortality.¹⁶

Patient infected with COVID 19 are also in risk of getting Venous Thromboembolic event (VTE).¹⁷ Factors which contribute in developing VTEs are systemic inflammation, coagulation abnormality, multi organ dysfunction.¹⁸ Another study suggests that there is abnormality in coagulation pathway in COVID 19 patients with increased level of D-dimer.¹⁹ If the level of D-dimer was found to be greater than 1mcg/mL, then it may be the major reason behind the mortality in COVID 19 patients.^{20,21}

Myopericarditis as an etiology for Cardiovascular collapse- The SARS CoV 2 can cause myocarditis and pericarditis that sometime may or may not be associated with pneumonia.²² Myocarditis may be identified using specialist imaging using Cardiac magnetic resonance and contrast enhancement, which is caused

by events such as necrosis, scarring, and myocardial edoema.²³ The focal or global Myocardial inflammation are the results of myocarditis. Focal myocarditis can mimic an acute coronary syndrome, or may present with acute chest pain or angina and can result in coronary angiography emergency.²⁴

Right ventricular failure as a cause of cardiovascular collapse- Patients with severe COVID 19 are at a significant risk of developing adult respiratory distress syndrome. COVID 19 patients have a higher risk of developing deep venous thrombosis and acute pulmonary embolism. In individuals with severe COVID 19, factors that may compromise right ventricular function include vasoplegic shock myocarditis and acute coronary syndrome.²⁵

4. Medication interactions

Newly medications interact broadly with other CV drugs including anticoagulants, antiplatelet and statin. In the treatment of COVID-19, many of agents may have interaction. They show interaction with oral antiplatelet drugs. Protease inhibitor like Lopinavir/Rotinavir inhibit CYP3A4 metabolism. The active metabolites of Clopidogrel is formed by CYP2C19 but inhibition CYP3A4 may also reduce the effective dosage of Clopidogrel.²⁶

5. Discussion

Cardiovascular patients are particularly vulnerable in the event of a COVID-19 pandemic, as SARS CoV-2 may exacerbate their condition. Some drugs are used to treat this illness; however studies have shown that these treatments can have negative impacts on the health of individuals who are already suffering from cardiovascular disease.

6. Conclusion

COVID-19 infection is associated with the cardiovascular disease like myocardial injury and myocarditis, VTE, AMI, Heart failure. Medications that are used to treat COVID 19 also have potential adverse effects on heart and circulatory system. When treating COVID 19 patients, it is essential for persons with cardiovascular disease to be informed of the contraindications of medications, as well as for clinicians to be aware of these side effects or consequences.

8. References

1. Ministry of Health and Family Welfare, Government of India, <https://www.mohfw.gov.in>.
2. The world health organization(WHO). Coronavirus disease (COVID-19) situation report. Available at: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>.
3. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease (COVID-19) outbreak in China: summary of a report of 72314 cases from the Chinese center for disease Control and prevention. *J Am Med Assoc*. 2020.
4. Wu Q, Zhou L, Sun X, et al. Altered lipid metabolism in recovered SARS patients twelve years after infection. *Sci Rep*. 2017;7: 9110.
5. Xiong TY, Redwood S, Prendergast B, Chen M. Coronaviruses and the cardio-vascular system: acute and long-term implications. *Eur Heart J*. 2020.
6. Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with novel coronavirus-infected pneumonia in Wuhan, China. *J Am Med Assoc*. 2020.
7. Li B, Yang J, Zhao F, et al. Prevalence and impact of cardiovascular metabolic diseases on COVID-19 in China. *Clin Res Cardiol*. 2020.
8. Lippi G, Plebani M. Laboratory abnormalities in patients with COVID-2019 infection. *Clin Chem Lab Med*. 2020.
9. Driggin E, Madhavan MV, Bikdeli B, et al. Cardiovascular Considerations for Patients, Health Care Workers, and Health Systems During the Coronavirus Disease (COVID-19) Pandemic. *J Am Coll Cardiol*. 2020; 20: 4637-4.
10. Chen D, Li X, Song Q, Hu C, Su F, Dai J. Hypokalemia and clinical implications in patients with coronavirus disease. 2019; (COVID-19). *medRxiv*. 2020.
11. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020.
12. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020; 395.
13. Chen C, Zhou Y, Wang DW. SARS-CoV-2: a potential novel etiology of fulminant myocarditis. *Herz*. 2020.

14. Xu Z, Shi L, Wang Y, et al. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *Lancet Respir Med*. 2020; 8(4):420-422.
15. Zhou F, Yu T, Du R et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020; 28:1054-1062.
16. Inciardi RM, Lupi L, Zaccone G, et al. Cardiac involvement in a patient with coronavirus disease 2019 (COVID-19). *JAMA Cardiol*. 2020.
17. Olejniczak M, Schwartz M, Webber E, et al. Viral myocarditis – incidence, diagnosis and management. *J Cardiothorac Vasc Anesth*. 2020.
18. Driggin E, Madhavan MV, Bikdeli B, et al. Cardiovascular considerations for patients, health care workers, and health systems during the coronavirus disease 2019 (COVID-19) pandemic. *J Am Coll Cardiol*. 2020.
19. Clerkin KJ, Fried JA, Raihelkar J, et al. Coronavirus disease 2019 (COVID-19) and cardiovascular disease. *Circulation*. 2020.
20. Zheng YY, Ma YT, Zhong JY, et al. COVID-19 and the cardiovascular system. *Nat Rev Cardiol*. 2020.
21. Xiong TY, Redwood S, Prendergast B, et al. Coronaviruses and the cardiovascular system: acute and long-term implications. *Eur Heart J*. 2020.
22. Inciardi RM, Lupi L, Zaccone G, et al. Cardiac involvement in a patient with coronavirus disease 2019 (COVID-19). *JAMA Cardiol*. 2020.
23. Olejniczak M, Schwartz M, Webber E, et al. Viral myocarditis – incidence, diagnosis and management. *J Cardiothorac Vasc Anesth*. 2020.
24. Friedrich MG, Sechtrm U, Schulz-Menger J, et al. Cardiovascular magnetic resonance in myocarditis: a JACC white paper. *J Am Coll Cardiol*. 2009; 53: 1475-1487.
25. Libby P, Simon D. Inflammation and thrombosis: the clot thickens. *Circulation*. 2001; 103: 1716-1720.
26. Behnood B, Mahesh V, David J, Taylor C. COVID-19 and Thrombotic or Thromboembolic Disease: Implications for Prevention, Antithrombotic Therapy and Follow-Up. *Journal of the American college of cardiology*. 2020; 75:23.

UNDR PEER REVIEW