

ANTIBACTERIAL, ANTIFUNGAL AND ANTIHELMINTHIC PROPERTIES OF ETHANOLIC, METHANOLIC AND WATER EXTRACTS OF POLLEN

Abstract

Microorganisms and helminthes can cause serious diseases in humans as well as in animals. The use of antimicrobial and antihelminthic drugs have created selective pressure and caused resistance to antibiotics used against them, thus it necessitates the use of honey bee's derived natural products. One such bee derived product is pollen, collected by worker honey bees from the flowering plants and modify it by adding its salivary secretions. The present study embodies use of pollen as antimicrobial and antihelminthic substance. Among microorganisms 4 Gram (+ve) bacteria; (*Bacillus subtilis*, *Staphylococcus epidermidis*, *Staphylococcus aureus*, *Streptococcus pneumoniae*) and 3 Gram (-ve) bacteria; (*Escherichia Coli*, *Pseudomonas aeruginosa*, *Salmonella enteric*) and 2 yeasts (*Candida albicans* and *Saccharomyces cerevisiae*) were used and the methodology used was disc diffusion assay and broth dilution method. The antihelminthic effect was observed among amphistomes via bioassay method under *in vitro* conditions. For observations three types of pollen extracts (ethanolic, methanolic and water extract) were prepared and positive controls used were; Ampicillin for antibacterial, Amphotericin B for antifungal and Albendazole for anti-helminthes. The antimicrobial activities were determined by measuring the zones of inhibition diameters in millimeters after 24 hours of incubation at optimum temperature for each microbe and also by broth dilution method. Results obtained showed that the water extract of pollen was found to be most effective against bacteria used in the present study where; Gram (+ve) bacteria were more susceptible as compared to the Gram (-ve) bacteria. It was also observed that among yeasts; *Saccharomyces cerevisiae* was more susceptible towards ethanolic extract of pollen while *Candida albicans* showed more inhibitions towards water extract of pollen. Results also demonstrated that none of the extracts of pollen was found to be effective against Helminthes (amphistomes) used in the present study.

Keywords: Pollen, antimicrobial, antihelminthic, disc diffusion, bioassay.

Introduction

The problem of bacterial resistance is growing very fast and the use of medicinal plants as antimicrobial and antihelminthic is a common practice. The recent increase in the popularity of alternative medicine and natural products has renewed interests in bee derived products as potential natural remedies. Among them bee pollen is one such valuable product having many pharmacological and medicinal properties through direct action against microorganisms or synergism with antimicrobial drugs. Bee pollen, also known as 'The Life Giving Dust' is collected by worker honey bees from the flowering plants. Worker bees have possessed pollen baskets (corbiculaor) on their hind legs which facilitate pollen transportation to the hive with the aid of salivary secretions. Bee pollen is regarded as functional foods for their high nutritional values as they are rich source of proteins, essential amino acids, sugars, fatty acids, vitamins, macro and microelements and are also rich in polyphenolic compounds¹. They also contain a wide variety of other health promoting compounds present in functional foods, such as prebiotics, probiotics, fibre, lignans, triterpenes, carotenoids, bioactive peptides and organic acids¹⁻⁵. Therefore it is considered as an excellent substitute of antibiotics⁶ and implemented as complementary medicine for a

large variety of impaired health conditions^{4, 7, 8}. Bee pollen exhibits antibacterial and antifungal activity^{1,3,9-12}.

The present study embodies results of investigations undertaken to evaluate the antimicrobial and antihelminthic properties of pollen by using *in vitro* methods. Determination of these activities was done by disc diffusion method and broth dilution method against pathogenic and non pathogenic bacteria and yeasts. In disc diffusion method, the microorganisms were screened for their susceptibility towards pollen (extracted in ethanol, methanol and water), applied on the disc of agar plate at the concentration range of 1.562-300mg/disc and for seeing antihelminthic activity water extract of pollen was used.

Material and Methods

Collection of Pollen: Pollen was collected from the pollen basket of worker honey bees returning to the hive with pollen loads, by installing a pollen trap at the entrance of the beehive.

Preparation of the pollen extract: The extract of pollen was prepared by following the method of Nagai¹³. For this 3g of fresh bee pollen was taken. It was suspended and extracted by shaking with 10 volumes of water at room temperature. The suspension was centrifuged at 10,000g in a refrigerated centrifuge for 20min. Supernatant fraction was collected and filtered. The filtrate was freeze dried. The powder of the extract was dissolved in ice-cold distilled water and filtered through 0.22 μ PTFE membrane for sterilization and was stored at cool place for use in experiment.

Helminths (Amphistomes): *Gastrothylax crumenifer*, were obtained from large intestine of sheep/goat procured from local slaughter house.

Microorganisms: Microorganisms such as bacteria (*Staphylococcus aureus*: MTCC No- 1144, *Staphylococcus epidermidis*: MTCC No-9040, *Streptococcus pneumoniae*: MTCC No- 2672, *Salmonella enterica*: MTCC No-3231, *E. coli*: MTCC No-2314, *Bacillus subtilis*: MTCC No-2435, *pseudomonas* MTCC No-3465 and fungi (*Candida albicans* (Yeast): MTCC No- 4748, *Saccharomyces cerevisiae* (Yeast): MTCC No- 3090) were procured from IMTECH (Institute of Microbial Technology) Sector-39, Chandigarh, India. The organisms were maintained in suitable/respective media (agar plates at 4⁰C). The strains were checked biochemically prior to usage.

***In vitro* anti helminthic activity of bee products**

Worm motility inhibition assay was employed for the evaluation of anti helminthic activity of bee pollen under *in vitro* conditions. The *in vitro* anti helminthic activity was conducted at three different

concentrations (100, 300, 500 mg/ml) to determine the inhibitory effect of bee pollen extracts on amphistome worms. Mature amphistome worms (*Gastrothylax crumenifer*) were collected from the large intestine of sheep/goat procured from local slaughter house (Fig.1). The worms were washed in phosphate buffered saline (PBS pH 7.2) and then suspended in PBS. Albendazole dissolved in 1% DMSO and diluted in PBS at concentrations of 5, 10 and 15 µg/ml and PBS alone served as positive and negative control respectively. There were three replicates for each treatment concentration. Ten vigorously motile worms were placed in each petri dish containing test solutions and observations were made at 15, 30, 60 and 120 min intervals for cessation of motility by gross visual motility of worms as index for anti helminthic activity. After exposure to different treatments, the worms were put in lukewarm PBS for 30 min in order to confirm mortality.



Fig.1. Amphistomes (*Gastrothylax crumenifer*) from stomach of sheep/goat.

Results and Discussion

The natural products of both plant and animal origin have wide range of pharmacological activities such as; antimicrobial, anti-inflammatory and anti-modulatory¹⁴⁻¹⁷. Currently the use of natural products has increased due to reduction in number of effective antibiotics provided by the pharmaceutical industries as well as due to the increasing drug resistance among microorganisms¹⁸⁻²². Microbes are evolving various mechanisms of antibiotics resistance and in some cases become multi drug resistant²³⁻²⁶. Therefore, a systematic study was carried out to evaluate the antimicrobial potential of pollen against a range of Gram (+ve) and Gram (-ve) bacteria as well as in yeasts by disc diffusion assay and broth dilution method and against amphistome by bioassay method under *in vitro* conditions.

In vitro antimicrobial activity of Pollen

Disc diffusion method:

Antibacterial activity of pollen was evaluated for three types of extracts *viz.* ethanolic, methanolic and water. Organisms initially selected for antimicrobial studies with pollen were nonpathogenic Gram (+ve)

and Gram (-ve) bacteria viz. *Bacillus subtilis*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Streptococcus pneumoniae*. Thereafter putative pathogenic Gram (+ve) bacteria viz. *Staphylococcus aureus*, *Staphylococcus epidermidis* and Gram (-ve) bacteria viz. *Salmonella enterica* which were screened for the inhibitory activity of pollen by disc diffusion method and broth dilution method. The stock solutions of pollen were made at a concentration of 300mg/ml. These were serially diluted to obtain the concentration of 300mg/ml, 200mg/ml, 100 mg/ml, 50 mg/ml, 25 mg/ml, 12.5 mg/ml, 6.25 mg/ml, 3.125 mg/ml and 1.562 mg/ml. Agar plates were made and 25-50 μ l of each organism was uniformly spread on the plates. Inoculum was always prepared fresh 24-48hrs prior to start of the experiment. Then 25 μ l of all the above mentioned concentrations of each product was applied on separate agar plates and incubated at their respective growth conditions. After 24-48 hrs clear zones of inhibitions of culture growth around the discs having pollen were measured. Results obtained are shown in Tables 1-10. The effectiveness of pollen was also compared with standard antibiotic as positive controls, such as; ampicillin (antibacterial), Amphotericin B (antifungal) and Albendazole (anti-helminthes).

Pollen:

(a). Ethanolic Extract:

The values observed for ethanolic extract of pollen for Gram (+ve) bacteria such as *Staphylococcus epidermidis* varied from 15.18 \pm 0.78-17.33 \pm 0.57mm, the zones of inhibition observed for *Staphylococcus aureus* varied from 13.50 \pm 0.89-18.5 \pm 1.20mm and for *Bacillus subtilis* from 11.00 \pm 0.63-12.55 \pm 1.22mm at concentrations ranging from 50-300mg/ml. For *S. epidermidis*, *S. aureus* and *Bacillus subtilis* no inhibition zones observed at concentrations ranging from 1.562-25mg/ml. In case of *Streptococcus pneumoniae* the zones of inhibition ranged from 9.89 \pm 0.16-13.46 \pm 1.49mm at concentrations from 100-300mg/ml and no inhibition zones were observed from 1.562-50mg/ml of ethanolic extract of pollen as shown in Table 1. The inhibition zones shown by ethanolic extract of pollen against Gram (-ve) bacteria such as *E. coli* varied from 8.00 \pm 1.15-10.28 \pm 0.33mm at concentrations from 50-300mg/ml. The values observed for *Pseudomonas aeruginosa* varies from 7.5 \pm 1.02-8.5 \pm 1.07mm at concentrations from 200-300mg/ml, the zones of inhibition observed against *Salmonella enterica* varied from 8.6 \pm 1.00-13.0 \pm 1.02mm at concentration range of 100-300mg/ml of ethanolic extract of pollen (Table 2). The zone of inhibition obtained against *Candida albicans* was 7.5 \pm 1.02mm at 300mg/ml and no inhibitions were observed at concentration lower than this, which showed that for *Candida albicans* the ethanolic extract of pollen was not much effective. The inhibition zones observed against *Saccharomyces cerevisiae* ranged from 13.50 \pm 0.89-16.2 \pm 1.30mm at concentrations varying from 100-300mg/ml of ethanolic extract of pollen. Here the results obtained showed that *Saccharomyces cerevisiae*

was more sensitive as compared to *Candida albicans* to ethanolic extract of pollen used in the present studies (Table 3).

Table 1. Antimicrobial activity of ethanolic extract of pollen against Gram (+ve) bacteria					
Gram (+ve) Bacteria					
Ethanolic extract of propolis		<i>Bacillus subtilis</i>	<i>Staphylococcus epidermidis</i>	<i>Staphylococcus aureus</i>	<i>Streptococcus pneumoniae</i>
S. No	(mg/ml)	Zones of inhibitions (mm)			
1.	1.562-25	NI	NI	NI	NI
2.	50	11.00±0.63*	15.18±0.78	13.50±0.89	NI
3.	100	10.85±1.16	15.40±0.98	14.2±0.30	9.89±0.16
4.	200	11.05±1.57	17.35±1.00	16.2±1.30	11.99±0.29
5.	300	12.55±1.22	17.33±0.57	18.5±1.20	13.46±1.49

*All the values are expressed as mean ± S.D (n=5)

Table 2. Antimicrobial activity of ethanolic extract of pollen against Gram (-ve) bacteria				
Gram (-ve) Bacteria				
Ethanolic extract of propolis		<i>Escherichia Coli</i>	<i>Pseudomonas aeruginosa</i>	<i>Salmonella enterica</i>
S. No	(mg/ml)	Zones of inhibitions (mm)		
1.	1.562-25	NI	NI	NI
2.	50	8.00±1.15*	NI	NI
3.	100	8.10±0.86	NI	8.6±1.00
4.	200	9.20±0.71	7.5±1.02	9.4±2.01
5.	300	10.28±0.33	8.5±1.07	13.0±1.02

*All the values are expressed as mean± S.D (n=5)

Table 3. Antimicrobial activity of ethanolic, methanolic and water extracts of pollen against yeast							
Pollen extracts		<i>Candida albicans</i>			<i>Saccharomyces cerevisiae</i>		
		WEP	EEP	MEP	WEP	EEP	MEP
S. No	mg/ml	Zones of inhibitions (mm)					
1.	1.562-6.25	NI	NI	NI	NI	NI	NI
2.	12.5	NI	NI	NI	8.70±0.34*	NI	NI
3.	25	NI	NI	NI	10.43±0.33	NI	NI

4.	50	NI	NI	NI	11.50±0.08	NI	NI
5.	100	NI	NI	NI	12.65±0.24	13.50±0.89	8.6±0.96
6.	200	9.8±0.06	NI	NI	14.43±0.43	14.2±0.30	8.22±1.02
7.	300	11.8±0.32	7.5±1.02	NI	14.05±1.54	16.2±1.30	10.54±2.01
*All the values are expressed as mean± S.D (n=5)							

(b). *Methanolic extract:*

The values observed for methanolic extract of pollen for Gram (+ve) bacteria such as *Staphylococcus epidermidis* varied from 9.55±0.47-11.25±0.53mm at range of concentrations from 100-300mg/ml. The zones of inhibition observed for *Staphylococcus aureus* varied from 8.6±1.00-12.4±.098mm at range of concentrations from 50-300mg/ml of methanolic extract of pollen; for *Bacillus subtilis* it was 8.43±1.39-9.72±1.09 mm at range of concentrations from 100-300mg/ml. The values for *Streptococcus pneumoniae* ranged from 10.29±0.39-11.26±1.19mm at concentrations from 200-300mg/ml and no inhibition zones observed from 1.562-100mg/ml of methanolic extract of pollen as shown in Table 4. The present studies were in agreement with Sramkova²⁷ where the antimicrobial effect of pollen samples was tested by using the agar well diffusion method and the results showed that the most sensitive bacteria to ethanolic extract of poppy pollen was *Staphylococcus aureus*. The zones of inhibition observed for methanolic extract of pollen against Gram (-ve) bacteria such as *E. coli* varied from 7.025±0.83-8.925±0.44mm at range of concentrations from 100-300mg/ml. No inhibition zones were observed from 1.562-50mg/ml of methanolic extract of pollen. Moreover *Salmonella enterica* was found to be insensitive against all the concentrations 1.562-300mg/ml used for evaluating the antibacterial activity of pollen methanolic extract. The value observed against *Pseudomonas aeruginosa* was 8.56±0.98mm at 300mg/ml of methanolic extract of pollen and no inhibitions was observed at lower concentrations as shown in Table 5. From the results obtained it was concluded that highest inhibition was found against *E. coli* and least against *Salmonella enterica* with all the extracts. Aboude⁹ also obtained similar results by analyzing samples of bee bread and bee pollen from different aromatic and medicinal plants and observing them for their antimicrobial activities. They observed that Gram (+ve) bacteria were more sensitive to bee bread and bee pollen than Gram (-ve) bacteria. Further results obtained for yeasts revealed that there were no zones of inhibition found against *Candida albicans* by using the entire range of concentrations (1.562-300mg/ml) of the methanolic extract of pollen, while *Saccharomyces cerevisiae* showed inhibitions ranging from 8.6±0.96-10.54±2.01mm at concentrations varying from 100-300mg/ml of methanolic extract of pollen thus concluding that *Saccharomyces*

Cerevisiae was more sensitive to methanolic extract of pollen as compared to *Candida albicans* as shown in Table 3. It has previously been reported that pollen was more effective in water extract as compared to the ethanolic and methanolic extracts²⁸. The present study showed highest inhibition was observed with water extract of pollen against Gram (+ve) organisms than ethanolic extract of pollen and least by the methanolic extract of pollen.

Table 4. Antimicrobial activity of methanolic extract of pollen against Gram (+ve) bacteria					
Gram (+ve) Bacteria					
Methanolic extract of pollen		<i>Bacillus subtilis</i>	<i>Staphylococcus epidermidis</i>	<i>Staphylococcus aureus</i>	<i>Streptococcus pneumoniae</i>
S. No	(mg/ml)	Zones of inhibitions (mm)			
1.	1.562-25	NI	NI	NI	NI
2.	50	NI	NI	8.6±1.00*	NI
3.	100	8.43±1.39	9.55±0.47	8.22±1.02	NI
4.	200	8.78±1.00	10.20±0.43	10.54±2.01	10.29±0.39
5.	300	9.72±1.09	11.25±0.53	12.4±.098	11.26±1.19
*All the values are expressed as mean ± S.D (n=5)					

Table 5. Antimicrobial activity of methanolic extract of pollen against Gram (-ve) bacteria				
Gram (-ve) Bacteria				
Methanolic extract of pollen		<i>Escherichia Coli</i>	<i>Pseudomonas aeruginosa</i>	<i>Salmonella enterica</i>
1.	1.562-50	NI		
2.	100	7.025±0.83*	NI	NI4z
3.	200	8.175±0.59	NI	NI
4.	300	8.925±0.44	8.56±0.98	NI
*All the values are expressed as mean± S.D (n=5)				

(c) Water extract:

The zones of inhibition observed for water extract of pollen against Gram (+ve) bacteria such as *Staphylococcus epidermidis* was observed to be in the range of 9.80±0.28-16.00±0.32mm at concentrations ranging from 12.5-300 mg/ml. No inhibition zones were observed at lower concentrations. The values observed for *Staphylococcus aureus* were from 8.70±0.34-15.49±1.51mm at concentration

range of 6.25-300mg/ml and there was no inhibition at lower concentrations. The range of inhibition observed for *Streptococcus pneumoniae* was from 8.90 ± 0.27 - 11.45 ± 1.09 mm at concentrations ranging from 25-300mg/ml, no inhibitions were observed at concentrations less than 25mg/ml of water extract of pollen. The range of inhibitions observed for *Bacillus subtilis* was 8.90 ± 0.47 - 11.25 ± 0.53 mm at concentrations from 25-300mg/ml water extract of pollen. No inhibitions against *Bacillus subtilis* were observed at concentrations of 1.562-12.5mg/ml water extract of pollen (Table 6). The antimicrobial activity observed by using water extract of pollen against Gram (-ve) bacteria showed no inhibition zones at concentrations ranging from 1.562-6.25mg/ml. The most sensitive Gram (-ve) bacteria was found to be *E. coli* and the values observed varied from 7.98 ± 0.41 - 11.25 ± 0.44 mm at concentration ranging from 12.50-300mg/ml. The zones of inhibition observed for *Salmonella enterica* ranged from 6.0 ± 1.47 - 10.8 ± 0.77 mm at concentration varying from 12.50-300mg/ml. The values observed for *Pseudomonas aeruginosa* were from 9.8 ± 0.06 - 15.0 ± 0.08 mm at 100-300mg/ml for water extract of pollen as shown in Table 7. The results obtained against *Candida albicans* varied from 9.8 ± 0.06 - 11.8 ± 0.32 mm at concentrations ranging from 200-300mg/ml and no inhibitory effect of water extract of pollen was observed at lower concentrations. For *Saccharomyces cerevisiae* results obtained showed zones of inhibition ranging from 8.70 ± 0.34 - 14.05 ± 1.54 mm at concentrations 12.5-300mg/ml. No inhibition zones were observed at concentrations less than 12.5mg/ml of water extract of pollen. Data obtained suggested that *Saccharomyces cerevisiae* was more sensitive as compared to *Candida albicans* to the water extract of pollen used in the present studies (Table 3).

Table 6. Antimicrobial activity of water extract of pollen against Gram (+ve) bacteria					
Gram (+ve) Bacteria					
Water extract of pollen		<i>Bacillus subtilis</i>	<i>Staphylococcus epidermidis</i>	<i>Staphylococcus aureus</i>	<i>Streptococcus pneumoniae</i>
S. No	(mg/ml)	Zones of inhibitions (mm)			
1.	1.562-3.125	NI	NI	NI	NI
2.	6.25	NI	NI	$8.70\pm 0.34^*$	NI
3.	12.5	NI	9.80 ± 0.28	10.43 ± 0.33	NI
4.	25	8.90 ± 0.47	10.23 ± 1.08	11.50 ± 0.08	8.90 ± 0.27
5.	50	10.13 ± 0.34	11.35 ± 0.44	12.65 ± 0.24	9.63 ± 0.74
6.	100	10.55 ± 0.66	13.18 ± 0.60	14.43 ± 0.43	10.85 ± 0.56
7.	200	11.13 ± 0.73	14.58 ± 0.21	14.05 ± 1.54	10.19 ± 0.99

8.	300	11.25±0.53	16.00±0.32	15.49±1.51	11.45±1.09
*All the values are expressed as mean ± S.D (n=5)					

Table 7. Antimicrobial activity of water extract of pollen against Gram (-ve) bacteria				
Gram (-ve) Bacteria				
Water extract of pollen		<i>Escherichia Coli</i>	<i>Pseudomonas aeruginosa</i>	<i>Salmonella enterica</i>
S. No	(mg/ml)	Zones of inhibitions (mm)		
1.	1.562-6.25	NI	NI	NI
2.	12.5	7.98±0.41*	NI	6.0±1.47
3.	25	8.65±0.37	NI	6.5±0.56
4.	50	9.88±0.57	NI	8.30±0.55
5.	100	10.0±0.49	9.8±0.06	9.2±1.29
6.	200	10.05±0.31	11.8±0.32	10.0±0.76
7.	300	11.25±0.44	15.0±0.08	10.8±0.77
*All the values are expressed as mean± S.D (n=5)				

These studies were in agreement with Aboude⁹ who reported antimicrobial activities of bee bread and bee pollen against bacterial strains isolated from pathological conditions in man. They revealed that Gram (+ve) bacteria are more sensitive to bee bread and bee pollen than Gram (-ve) bacteria. This was also supported from the studies of Pascoal²⁹, where antimicrobial activity of bee pollen against Gram (+ve) was being more sensitive as compared to Gram (-ve) bacteria. In his studies *Staphylococcus aureus* was the most sensitive and *Candida glabrata* was found to be most resistant of the microorganisms studied. Sramkova⁷³ determined antioxidant and antibacterial activity of monofloral bee pollen samples against pathogenic bacteria. The antimicrobial effect of pollen samples were tested by using the agar well diffusion method. The most sensitive bacteria to the poppy pollen ethanolic extract was *Staphylococcus aureus* (70%) The most sensitive bacteria to rape bee pollen methanolic extract (70%) and sunflower ethanolic extract (70%) was *Salmonella enterica*.

Broth dilution method:

For determination of inhibitory concentrations of the honey bee product pollen against the organisms listed previously and to study the effect of a range of concentrations of different extracts on the growth of an organism, experiments were done with broth dilution method. Organisms were grown in presence of pollen at concentrations ranging from 3mg/ml-60mg/ml. Growth of Gram (+ve) and Gram (-

ve) non-pathogenic bacteria viz. *Bacillus subtilis*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Streptococcus pneumoniae* was measured at late log phase. Then pathogenic Gram (+ve) bacteria viz. *Staphylococcus aureus*, *Staphylococcus epidermidis* and Gram (-ve) bacteria viz. *Salmonella enterica* were screened separately for the inhibitory activity of honey bee products by broth dilution assay. Growth of each organism was measured at late log phase by taking O.D. at 600nm (Table 8). Determination of antimicrobial activity by broth dilution method for bee pollen revealed that there is concentration dependent decline in growth of organisms under study. Therefore this concludes the antimicrobial properties of pollen. The antimicrobial properties observed for bee pollen could be due to cell wall lyses and plasma membrane degradation, which leads to a loss of potassium ions and the damage, caused provoking cell autolysis³⁰.

Table: 8 Optical density observed against Gram (+ve) and Gram (-ve) bacteria with water extract of pollen:

Pollen	(O.D) for Gram (+ve) Bacteria				(O.D) for Gram (-ve) Bacteria		
	<i>Bacillus subtilis</i>	<i>Staphylococcus epidermidis</i>	<i>Staphylococcus aureus</i>	<i>Streptococcus pneumoniae</i>	<i>Escherichia coli</i>	<i>Pseudomonas aeruginosa</i>	<i>Salmonella enterica</i>
Control	0.99	1.55	1.45	1.65	1.68	1.47	1.42
3mg/ml	0.850	1.42	1.36	1.53	1.60	1.20	1.34
7.5mg/ml	0.81	1.31	1.21	1.46	1.43	1.16	1.21
15mg/ml	0.79	1.22	1.16	1.31	1.35	1.09	1.15
30mg/ml	0.65	0.99	0.98	1.20	1.22	0.96	1.08
60mg/ml	0.50	0.82	0.88	1.11	1.15	0.88	0.78

Table 9 In vitro anti helminthes activity:

Bee products	O.D for <i>Candida albicans</i>						O.D for <i>Saccharomyces cerevisiae</i>					
	Control	3m g/ml	7.5 mg/ml	15 m g/ml	30 m g/ml	60 m g/ml	Cont rol	3mg/ml	7.5m g/ml	15m g/ml	30mg/ml	60mg/ml
Pollen	2.25	1.85	1.79	1.72	1.65	1.44	2.65	2.57	2.32	2.10	1.90	1.78

Parasitic infections have always been a major concern to the medical field and among them amphistomes are the prominent causal agents of diseases in humans as well as in animals especially sheep and goat, which ultimately cause considerably economic losses to livestock industry and hence to the economic development of a country. Over the past few years, medical science has led many milestones in parasitological research but still efficient products to control helminthes are yet to discover. Moreover the drugs used for this purpose has caused resistance considerable toxicity to humans beings through foods derived from livestock causing serious

health hazards³¹. This makes the necessity of use of natural products as antihelminthic drugs. Among them use of medicinal plants continues to be the most fruitful approach towards antihelminthic drugs³². The present study was undertaken to evaluate anti helminthes activity of pollen by Petri dish method³³, in comparison with a standard drug Albendazole, against amphistome (*Gastrothylax crumenifer*) parasitizing the large intestine of sheep/goat through *in vitro* studies by the worm motility inhibition assay.

Bee products Extracts	Concentrations	15min	30min	60min	120min
Water extract of pollen	100mg/ml	10	9	8	8
	300mg/ml	9	8	8	7
	500mg/ml	9	8	9	8
Positive control (Albendazole)	5µg/ml	8	6	2	0
	10µg/ml	8	5	2	1
	15µg/ml	8	5	3	0
Negative control	(Normal saline only)	9	8	6	4

Water extracts of pollen was used for this study as this was observed to be the most effective for microorganisms tested during the *in vitro* study. Mortality was observed after every 15, 30, 60 and 120 min in the entire test group. The data obtained is presented in Table 10. Perusal of the results obtained revealed that pollen at the highest concentration tested (500mg/ml) after completion of 120min of the experiment did not give any more mortality than the negative control (3 and 4 live amphistome respectively) and was therefore not effective in controlling the parasite. The positive control using Albendazole, however at much lower concentration (5, 10, 15µg/ml) was able to arrest the parasite almost cent percent at the end of the experiment. Results therefore suggested that bee products were not potent anti helminthes agents and are not suitable for application against amphistome; *Gastrothylax crumenifer*.

Conclusion

Pollen: Zone of inhibition measurement done for the organisms studied showed that *Staphylococcus aureus* was most susceptible to ethanolic and methanolic extracts of pollen, followed by *S. epidermidis* and *Streptococcus pneumonia*. *Staphylococcus epidermidis* was also inhibited by water extract of pollen. In yeast, higher inhibition was observed for *Saccharomyces cerevisiae* than *Candida*

albicans with ethanolic extract of pollen while higher inhibition was observed with water extract of pollen against *Candida albicans*

In vitro antihelminthic activity: Amphistomes (*Gastrothylax crumenifer*) obtained from the gut of sheep/goat were taken as test organism. The entire range of bee products at all concentrations used in the present study did not show any effect different from the negative control on the mortality of amphistome. The positive control using Albendazole was very effective even at much lower concentrations.

References

1. Didaras NA, Karatasou K, Dimitriou TG, Amoutzias GD, Mossialos D, Antimicrobial Activity of Bee-Collected Pollen and Beebread: State of the Art and Future Perspectives. *Antibiotics* (Basel). 9 (2020) 811. doi:10.3390/antibiotics9110811.
2. Isidorov V, Isidorova A, Szczepaniak L, Lazarek U, Gas chromatographic-mass spectrometric investigation of the chemical composition of beebread. *Food Chem.* 115 (2009) 1056.
3. Bakour M, Fernandes A, Barros L, Sokovic M, Ferreira I, Badiia L, Bee bread as a functional product: Chemical composition and bioactive properties. *LWT.* (2019) 109.
4. Märgäoan R, Strant M, Varadi A, Topal E, Yücel B, Cornea-Cipcigan M, Campos MG, Vodnar DC, Bee Collected Pollen and Bee Bread: Bioactive Constituents and Health Benefits. *Antioxidants.* 8 (2019) 568.
5. Kostic AŽ, Milincic DD, Bara'c MB, Ali Shariati M, Tešić ŽL, Peši'c MB, The Application of Pollen as a Functional Food and Feed Ingredient-The Present and Perspectives. *Biomolecules.* 10 (2020) 84.
6. Abdelnour SA, Abd El-Hack ME, Alagawany M, Farag MR, Elnesr SS, Beneficial impacts of bee pollen in animal production, reproduction and health. *J. Anim. Physiol. Anim. Nutr.* 103 (2019) 477.
7. Komosinska-Vassev K, Olczyk P, Kazmierczak J, Mencner L, Olczyk K, Bee Pollen: Chemical Composition and Therapeutic Application. *Evid. Based Complement. Altern. Med.* (2015) 297425.
8. Kieliszek M, Piwowarek K, Kot A, Błazejak S, Chlebowska-Smigiel A, Wolska I, Pollen and bee bread as new health-oriented products: A review. *Trends Food Sci. Technol.* 71 (2018) 170.
9. Abouda Z, Zerdani I, Kalalou I, Faid M and Ahami MT, The antibacterial activity of Moroccan bee bread and bee pollen (fresh and dried) against pathogenic bacteria. *Research Journal of Microbiology.* 6 (2011) 376.
10. Graikou K, Kapeta S, Aligiannis N, Sotiroudīs G, Chondrogianni N, Gonos E, Chinou I, Chinou I, Chemical analysis of Greek pollen-antioxidant, antimicrobial and proteasome activation. *Chem. Cent. J.* 5 (2011) 33.

11. Morais M, Moreira L, Feás X, Estevinho LM, Honeybee-collected pollen from five Portuguese Natural Parks: Palynological origin, phenolic content, antioxidant properties and antimicrobial activity. *Food Chem. Toxicol.* 49 (2011)1096.
12. Kaškonienė V, Adaškevičiūtė V, Kaškonas P, Mickienė R, Maruška A, Antimicrobial and antioxidant activities of natural and fermented bee pollen. *Food Biosci.* 34 (2020) 100532.
13. Nagai T, Nagashima T, Myoda T, Inoue R. Preparation and functional properties of extracts from bee bread. *Nahrung.* 2004 48(3):226-9. doi: 10.1002/food.200300421.
14. Abu-Al-Basal M A. Healing potential of *Rosmarinus officinalis* L. on full-thickness excision cutaneous wounds in alloxan-induced-diabetic BALB/c mice. *Journal of Ethnopharmacology.* 131 (2010), 443-450.
15. Zhao L, La V D and Grenier D. Antibacterial, antiadherence, antiprotease and anti inflammatory activities of various tea extracts: potential benefits for periodontal diseases. *Journal of medicinal food.* 16 (2013): 428-436.
16. Mansourian, A., Boojarpour, N., Ashnagar, S., Momen Beitollahi, J. & Shamshiri, A. R. The comparative study of antifungal activity of *Syzygium aromaticum*, *Punica granatum* and nystatin on *Candida albicans*; an in vitro study. *Journal of Medical Mycology* 24, e163- 461 168, doi:10.1016/j.mycmed.2014.07.001 (2014).
17. Hashemipour, M. A., Tavakolineghad, Z., Arabzadeh, S. A., Iranmanesh, Z. & Nassab, S. A. 463 Antiviral activities of honey, royal jelly, and Acyclovir against HSV-1. *Wounds : a 464 compendium of clinical research and practice* 26, 47-54 (2014).
18. McGrath, L. J., Becker-Dreps, S., Pate, V. & Brookhart, M. A. Trends in antibiotic treatment of acute otitis media and treatment failure in children, 2000-2011. *PLoS One* 8, e81210, doi:10.1371/journal.pone.0081210 468 PONE-D-13-29268 (2013).
19. Filius, P. M. & Gyssens, I. C. Impact of increasing antimicrobial resistance on wound management. *American Journal of Clinical Dermatology* 3, 1-7, doi:030101 (2002).
20. Hwang, A. Y. & Gums, J. G. The emergence and evolution of antimicrobial resistance: Impact on a global scale. *Bioorganic & Medicinal Chemistry* doi:S0968-0896(16)30261-9 10.1016/j.bmc.2016.04.027 (2016).
21. Falagas, M. E. et al. Outcome of infections due to pandrug-resistant (PDR) Gram negative bacteria. *BMC Infectious Diseases* 5, 24, doi:1471-2334-5-24 10.1186/1471-2334-5-24 (2005).
22. Bathoorn, E. et al. Emergence of pan-resistance in KPC-2 carbapenemase-producing *Klebsiella pneumoniae* in Crete, Greece: a close call. *Journal of Antimicrobial Chemotherapy* 71, 1207-1212, doi: 10.1093/jac/dkv467 (2016).
23. Vilacoba, E. et al. Widespread dispersion of the resistance element tet(B)::ISCR2 in XDR *Acinetobacter baumannii* isolates. *Epidemiology & Infection* 144, 1574-1578, 507 doi:S0950268815002897 10.1017/S0950268815002897 (2016).
24. Weterings, V. et al. An outbreak of colistin-resistant *Klebsiella pneumoniae* carbapenemase-producing *Klebsiella pneumoniae* in the Netherlands (July to December 511 2013), with inter-institutional spread.

European Journal of Clinical Microbiology & Infectious Diseases 34, 1647-1655, doi:10.1007/s10096-015-2401-2 513 10.1007/s10096-015-2401-2 (2015).

25. Iyamba, J. M., Wambale, J. M., Lukukula, C. M. & za Balega Takaisi-Kikuni, N. High prevalence of methicillin resistant staphylococci strains isolated from surgical site infections in Kinshasa. *Pan African Medical Journal* 18, 322, doi:10.11604/pamj.2014.18.322.4440.
26. Kulkova, N., Babalova, M., Sokolova, J. & Krcmery, V. First report of New Delhi metallo beta-lactamase-1-producing strains in Slovakia. *Microbial Drug Resistance* 21, 117-120, doi:10.1089/mdr.2013.0162 (2015).
27. Sramkova A, Kuo SM and Popova, Antioxidant and antibacterial property of mono-floral pollen sample against pathogenic bacteria. *Food chemistry*. 9 (2013) 34.
28. Devi A, Kumar NR and Kaur J, Evaluation of the antioxidative potential of Bee Products: Pollen and Bee Bread against *Staphylococcus aureus* Infected Balb/c mice. *Journal of Chemical and Pharmaceutical Sciences*.4 (2016).
29. Pascoal A, Rodrigues S, Teixeira A and Estevinho LM, Biological activities of commercial bee pollens: Antimicrobial, antimutagenic, antioxidant and anti-inflammatory. *Food and Chemical Toxicology*. 63 (2014) 233.
30. Mirzoeva OK, Grisjanin RN and Calder PC, Antimicrobial action of propolis and some of its components: the effects on growth, membrane potential and motility of bacteria. *Microbiology Research*. 152 (1997) 239.
31. Anbu, J., Murali,A., Sathiya, R., Saraswathy, GR., Azamthulla, M. *In Vitro* Anthelmintic Activity of Leaf Ethanolic Extract of *Cassia Alata* and *Typha Angustifolia*. *MSRUAS-SAS Tech Journal*. 2018; 14(2): 41-44.
32. Ghangale GR, Tushar M and Jadhav ND, *In vitro* antihelminthic activity of alcoholic extract of *Allivum Sativum* against rumen amphistome. *Veterinary boukra World*. 2 (2009) 385.
33. Aggarwal R, Suri M and Bagai U, *In vitro* anthelmintic effect of ethanolic and aqueous extracts of *calotropis procera*, *azadirachta indica* and *punica granatum* on *trichuris globulosa*, an intestinal nematode of sheep. *European Journal of Biomedical and Pharmaceutical sciences*. 3 (2016) 575.