

Role of Medicinal Plants in the Management of Diabetes Mellitus: Future Prospective

Abstract:

Diabetes mellitus is a worse metabolic condition in which level of glucose and lipid increased in blood circulation due to insufficient insulin functionality that may lead to worse the complication like atherosclerosis, hyperlipidemia, gangrene, limb amputation, kidney failure, and many more. Furthermore, high concentration of glucose in blood stream can directly produce reactive oxygen species which ultimately more worsen the conditions like hyperlipidemia, atherosclerosis etc. This disorder is expected to affect 9.3% of the global population (46.3 crore people) in 2019, increasing to 10.2% (57.8 crore) by 2030 and increasing 10.9 % (700 crore) by 2045. Medicinal plants are used as medicine to treat many ailments since a long time ago. But now they have been rediscovered with having specific constituents to treat diseases and many more yet to discover. There are so many active phytoconstituents of plants which are being extracted, purified, tested and formulated for patient convenience. In the present review, various plants used to treat diabetes mellitus due to have fewer side effects as compared to allopathic treatment (oral hypoglycemic drugs) are described. Many plants with hypoglycemic effect of many plants have been reported, and the mechanisms of action of these plants with these hypoglycemic behaviors are being examined through in vitro and in vivo studies. Some of these medicinal plants and their active constituents related to antidiabetic activity are discussed in this review.

Keyword: Diabetes mellitus, reactive oxygen species, low density lipoprotein, high density lipoprotein, ethanolic extract, aqueous extract, in-vitro, in-vivo.

INTRODUCTION

Diabetes mellitus is known as a severe metabolic condition marked by high glucose level in circulation, hyperlipemia, hyperaminoacidemia, and hypoinsulinemia, which occurs in a reduction in insulin secretion and operation. It's linked to the progression of minor and major vascular disorders including nephropathy, cerebrovascular disease, neuropathy and cardiovascular disease. This disease is related to a poorer quality of life as well as an increased risk of death and morbidity. Long-term hyperglycemia plays a major role in onset up to worsening of micro and macrovascular complications [1]. Diabetes is expected to affect 9.3% of the global population (46.3 crore people) in 2019, increasing to 10.2% (57.8 crore) by 2030 and 10.9% (70 crore) by 2045. The incidences are greater in urban areas (10.8%) than rural areas (7.2%), and in high-income countries (10.4%) than low-income countries (7.2%). One-half of those with diabetes (50.1%) are unaware that they have the disease [2]. Insulin and various oral antidiabetic drugs such as sulfonylureas, biguanides, alpha-glucosidase inhibitors, and glinides are the currently available treatment for DM. Medications are costly and hard to acquire in developing countries. Due to the adverse effects which are associated with oral hypoglycemic drugs (therapeutic agents) for the treatment of diabetes mellitus, nowadays herbal medicines are becoming increasingly popular [3]. As a result, conventional herbal remedies extracted from plants are primarily used and play a specific and effective role in the treatment of DM. [1] The hypoglycemic effect of many plants used as anti-diabetic treatments has been confirmed, and the mechanisms of these plants' behaviour are being examined. Herbal or plant-based ingredients are high in phenolic content, terpenoids, flavonoids, coumarins, and other constituents that help lower blood sugar levels [4].

Traditional therapeutics and alternative remedies derived from traditional medicinal plants are also discussed in this study. Orthodox medicines extracted from widely accessible medicinal plants carry a lot of promise for the development of new anti-diabetic medications. WHO (World Health Organization) has compiled a list of 21,000 medical herbs used around the world [5]. A list of medicinal plants with anti-diabetic and other positive properties, as well as herbal medicines used in diabetes care, has been compiled. The data for this study were retrieved from Web of Science, PubMed, and Google Scholar. Web of Science was chosen as search engine because it is the most widely accepted and frequently used database for analysis of scientific publications.

MEDICINAL PLANTS WITH ANTIDIABETIC POTENTIAL

Allium cepa

Allium cepa is widely known as onion, a spice plant that belongs to the Liliaceae family. It has been used in the prevention of several diseases since earlier civilizations. Bora and Sharma

described a comprehensive review on therapeutic potential of *Allium cepa*[6]. Hypoglycemic activity is one of *Allium cepa*'s most significant effects in diabetes mellitus, among its many health benefits. (S-methyl cysteine) Sulphur compounds and (quercetin) flavonoids are largely responsible for *Allium cepa*'s hypoglycemic activity, which helps in reducing glucose, lipids in circulation, oxidative stress. Onion extracts have been shown to have hypoglycaemic and hypolipidemic impact. In preliminary clinical trials, slices of *Allium cepa* were found to be safe to eat by diabetic patients and to have adequate hypoglycemic function[7]. Silver nanoparticles of *Allium cepa* were also prepared by green synthesis for diabetes treatment. UV-visible spectroscopy, FT-IR, and SEM were used to classify the synthesised silver nanoparticles. The particles have higher levels of α -amylase and inhibitory activities of α -glucosidase, according to the in vitro reports. Besides that, it had higher antioxidant activity and lower cytotoxicity. The green synthesised silver nanoparticles were reported to be an effective phyto-medicine for the treatment of DM[8].

Azadirachta indica

Azadirachta indica, commonly named as neem is a tropical and semi-tropical tree local to the Indian subcontinent. The ethanolic components of plant leaves were studied and found that plant enriched in antioxidant and antidiabetic flavonoid, and these were nicotiflorin, rutin, isoquercitrin, quercitrin and astragalol[9]. The use of flower extract enhanced functional recovery, especially in the areas of motor and sensory functions. Malondialdehyde levels were significantly reduced, while superoxide dismutase activity and axon density were significantly increased [10]. Reports suggested that the extract of plant's leaves decrease the level of glucose, cholesterol, and triglycerides in serum but there was an increase in weight of the body, tissue antioxidants, vascular endothelial growth factor contents and total collagen [11].

***Aegle marmelos* Correa-**

Aegle marmelos is commonly named as Bael is a fruit tree with fruits belonging to the Rutaceae family, widely grown all over the world. This plant has been getting prominence popularity due to its immense classical medicinal usage. Bael fruit has flavonoids, fibers, terpenoids, carotenoids, coumarins, phenolics and alkaloids[12]. The essential oil of *Aegle marmelos* also revealed that it has inhibitory activities against many important enzymes which are related to DM and has mild antioxidant ability[13]. In streptozotocin-induced diabetic rats, the effect of a chloroform extract of the plant was observed. Total and specific advanced glycation end products, protein carbonyl formation, and collagen solubility tests were used to assess antiglycation activity in vitro[14].

***Allium sativum*-**

A part of the Alliaceae family, *Allium sativum* L. (garlic) is commonly used as a worthwhile ingredient and a common cure for a range of infections and diseases. *Allium sativum* is grown almost everywhere in the world and it seems to have arisen in Asia before expanding to other areas of the globe. *Allium sativum* is a major source of S-binding components, the most significant of which is alliin. *Allium sativum* L. has a characteristic odour, flavour, and biological and medicinal properties owing to volatiles like allicin and sulphur compounds (lipid-soluble) like diallyl disulphide, ajoene, diallyl trisulphide, diallyl sulphide, and dithiols [15]. Dyslipidemia, a major risk factor for cardiovascular disease, is often linked to diabetes. According to existing data, *Allium sativum* has a good reputation for treating diabetic dyslipidemia. When compared to placebo, monitor, or baseline, the groups treated with garlic had significantly lower serum TG (19 to 36%) and fasting blood glucose (FBG, 18 to 56%). In patients who are suffering from

Type 2 DM, mixing garlic with metformin increased glycemic regulation and lipid profile[16].

***Aloe barbadensis* Miller-**

Aloe was formerly classified as part of the Liliaceae tribe, but it is now classified as a separate family, Aloaceae. Its origins are in South and East Africa, as well as the Mediterranean. It has over 400 species and can be found all over the world, though it is mainly found in subtropical areas[17]. Aloe-emodin, aloin, aloesin and emodin, are the major active constituents of aloe. Aloe-emodin, in particular, has emerged as a potential antimicrobial, antidiabetic, cytotoxic, cardioprotective and also anti-inflammatory and skin protective activity[18].

***Aloe ferox* L-**

Aloe ferox L., a perennial succulent commonly named as cape aloe that is belonging to the family Liliaceae and is described as having massive stemless leaves with a narrow edge and spongy dense scimitar leaves. In vitro experiments revealed that leaf of *Aloe ferox* extracts inhibited α -glucosidase but had no effect on α -amylase inhibition. Isoferulic acid-3-glucuronide, oleic acid, and stearic acid were found in *Aloe ferox* leaf powder. Isoferulic acid, cinnamic acid, and were present in *Aloe ferox* water extract. Reports revealed that *Aloe ferox* leaf powder showed promise in preventing weight gain and possibly having a hypoglycemic impact in animals who fed on high lipid diet possibly due to inhibitory activity of α -glucosidase that helped in the regulation of the glucose in circulation[19]. Supplementing with *A. ferox* resulted in a modest rise in serum insulin but no improvement in end-point plasma glucose levels. *A. ferox*, on the other hand, reported that there was a clinically significant improvement in the status of a diabetic person[20].

***Ficus religiosa*-**

F. religiosa commonly named as sacred fig and peepul isa member of the Moraceae family. There are some countable species of trees found with the largest varieties in Southeast Asia, tropical South America, and Australia[21]. It is reported that with the multiple organs of this tree Diabetes and many more diseases have all been treated. The use of *F. religiosa* bark extract showed in a substantial decrease in glucose in circulation in rats, with the effect being more articulated at 50 and 100 mg/kg than at 25 mg/kg. This has also been shown to have a critical antilipid peroxidative effect on diabetic rat's pancreas. The findings suggest that bark aqueous extract has anti-diabetic effects[22].

***Momordica charantia*-**

Momordica charantia (Family: Cucurbitaceae) is a tropical and subtropical plant, commonly named as bitter melon or bitter melo, balsam pear or karela[23]. It is grown in China, India, East Africa, and Central and South America. *M. charantia* is primarily used for food, the whole plant, especially the seeds and fruit, has long been used as classical medicine for the treatment of DM and its complications[24]. Major changes were reported in the absorption of glucose (61%) and release of adiponectin (75%) that were correlated with the drug mixture water extract and insulin compared to control doses[25]. The anti-adipose activity of *M. charantia* extract is contradicted by its inhibitory effect on leptin expression[23]. The important chemical constituents of *M. charantia* are: 1) heteropolysaccharides 2) proteins and peptides 3) Cucurbitanes and cucurbitacins: terpenoids and saponins; 4) flavonoids and phenolic extracts; 5) essential oils, sterols, amino acids and fatty acids. Water makes up 93.2 percent of a fruit's dry weight, while protein and lipids make up 18.02 and 0.76 percent, respectively[26]. Bitterness in bitter-melon is

caused by triterpene glycosides, specifically momordicoside K and momordicoside L. *M. charantia* juice showed significant antidiabetic capacity[27].

Lagenaria siceraria

The fruit of the bottle gourd (*Lagenaria siceraria*) belongs to the cucurbitaceae family[28]. Bottle gourd has been reported as an anti-diabetic and anti-cancer agent in some trials[29]. Administration of pulp powder showed a substantial low ($p < 0.01$) in triglyceride and level of VLDL-c, as well as a significant rise ($p < 0.01$) in HDL-c. These findings indicate that bottle gourd pulp powder (DBPP) has therapeutic potential and may show improved effects if ingested for a longer period of time.[28] Saponins and flavonoids are reported in *Lagenaria siceraria* fruit extract. Saponin has the capability to bind with cholesterol in the lumen of the intestine, preventing it from being absorbed, and it also boosts lipoprotein lipase function[30]. The fruit extract of *Lagenaria siceraria* is sometimes used as a potent Nutraceutical having important effects like lowering the lipid level in circulation and also showed antioxidant effects, and can be useful as a symptomatic measure for people who are at risk of cardiovascular disease[31]. Fruits of *L. siceraria* possess outstanding antioxidant effect and α -amylase inhibitory effect[32].

Ocimum sanctum

Ocimum sanctum also known as Holy Basil, is a Southeast Asian Ayurvedic herb with a long history of cultural use[33]. In *O. sanctum*, many chemical compounds are reported. The major compounds include eugenol methyl ether, caryophyllene, germacrene D, α -elemene, and copaene. The key constituent in *O. sanctum* is eugenol methyl ether[34]. Methanol extract and its effective fractions (ethyl acetate/butanol) have been shown antidiabetic activity due to polyphenolic compounds[35][36].

Eugenia jambolana

It is commonly known as black plum or Jamun belonging to Myrtaceae family. It is a large evergreen tree found throughout the Indian subcontinent[37]. The seed of *Eugenia jambolana* was reported to contain a variety of biologically active chemical constituents, including saponin, flavonoids, triterpenoids, gallic acid, glycosides and ellagic acid[38]. *E. jambolana* has been reported the antidiabetic and anti-oxidant property against streptozotocin-induced diabetic rats[39]. In experimental, diabetes induced rats gave treatment with ethanolic extract of kernel at a mentioned dose per body weight substantially improved glucose, urea and cholesterol in circulation that boosted tolerance of glucose and whole protein and liver glycogen levels[40].

Pterocarpus marsupium

The *Pterocarpus marsupium*, also known as the Indian Kino Tree or Vijyasara, Bija, belongs to the family of Fabaceae and is found throughout India[41]. Pterostilbene, (-) epicatechin, pterosupin, marsupin, tannins, and other bioactive principles found in the heart wood of this woody tree are responsible for its anti-diabetic function[42]. The results of an experimental study design for diabetic neuropathy screening the animals revealed that diabetic animals had neuroprotective behaviour. A study also reported that *P. Marsupium* pretreatment led to substantial lowering the level of glucose[43].

Trigonella foenum

It is a dicotyledonous plant with an annual harvest, commonly known as fenugreek and methi in

hindi belonging to the Papilionaceae subfamily of the Fabaceae family[44]. Trigonella foenum seeds were high in saponins, notably dioscin or diosgenin, which had been isolated and screened for α -glucosidase inhibitory activity in vitro, confirming natural anti-diabetic agents with minor toxicity[45]. (9Z,12Z)-N-((3R,4R,5S)-4,5-Dimethyl-2-oxotetrahydrofuran-3-yl) octadeca-9,12-dienamide (N55), a positive GLP-1 signalling modulator obtained from fenugreek seeds was reported as effective for treatment of diabetes[46].

Gymnema sylvestre

Gymnema sylvestre, commonly known as gurmar, a vulnerable and slow-growing plant belongs to Apocynaceae family and has historically been used to treat a number of diseases. It is found in India, Africa, Australia, and China as a wild herb[47]. The main chemical constituents of *G. sylvestre* are gymnemic acids, which are triterpenoid saponins and are believed to have anti-diabetic properties due to their biological activity[48]. Gymnemic acids possess anti-sweetener, anti-diabetic, and anti-inflammatory potential. Gymnemic acid molecules have an atomic structure as molecules of glucose. These molecules bind to taste binding site, blocking sugar molecules in food from inducing them and thereby reducing sugar cravings[49]. Further studies were shown that triterpene glycoside inhibit pancreatic amylase, glucosidase, sucrase, and maltase activity which were derived from *Gymnema sylvestre*[50].

Carica papaya

Carica papaya is a small herbaceous plant in the Caricaceae family. This plant is commonly grown for its edible, tasty fruit, which has a high nutritious value and is easy to digest[51]. Ethanolic extract of *C. papaya* leaf reported to have highest concentrations of tannins and steroids and both steroids and quinones were detected in the largest concentrations in chloroform extract[52]. It was also reported that the aqueous extract of *C. papaya* was found to retain body weight by activating the few residual β -cells, resulting in insulin release and decreased liver glycogen content[51].

Costus igneus

Costus igneus, commonly name insulin plant, Fiery Costus or Spiral Flag, is a herbaceous plant belongs to the family of Costaceae. It is grown in India to treat DM[53]. Three main proteins present in this plant are; Aglycin, Viglycin and ILP (insulin like protein) that directly stimulate IRs (Insulin Receptor) and promote secondary stimuli until all glucose uptake are done[54]. ILP act via insulin signaling pathway was further demonstrated by Mansi R. Hardikar et al through studied mechanism of action by experimenting on L6 myotubes (immortalized rat skeletal myoblast cell line)[55]. A stable and effective silver coated nanoparticles were also prepared and characterized for oral drug delivery[56]. It was also reported that *Costus igneus*-ZnO nanoparticles were more effective than simple leaf extract of *C. igneus*[57].

Cinnamomum zeylanicum

This traditional medicine is obtained from bark of trees belonging to the family of Lauraceae and also believed that plant has many health benefits such as reducing triglyceride, LDL, total cholesterol[58]. chemical profiling of aq. Extract showed total phenolic content and proanthocyanidin content in which cinnamaldehyde and cinnamic acid were main chemical constituent having α -glucosidase inhibitory and α amylase inhibitory activity[59]. Phenolic extract of plant is responsible for antioxidant and free radical scavenging activity also[60].

Zingiber officinale

It consists of fresh and dried roots of plant belongs to family of zingiberaceae, cultivated in india, South East Asia, Mexico and other parts of the world[61]. Zingerone was important chemical constituent and also found to be more antioxidant than BHA[62]. Ginger also showed the inhibitory action against alpha glucosidase enzyme which is helpful to treat type 2 DM.[63]. Methanolic extract of plant rhizomes found to be effective in weight management, reduction in higher lipid content and glucose level[64]. In vivo studies suggested that daily consumption of 1.2 g of ginger for 90 days effectively treat DM type 2 patients by decreasing blood sugar, total cholesterol and LDL[65].

Berberis aristata

It is spinous herb commonly known as Daruhaldi, Darhald, chitra or Indian beri beri, belonging to the family Berberidaceae. It is widely distributed throughout the Himalayan region[66]. Root bark of the plant contained protoberberine alkaloid like oxyberberine, karachine, berberine and berbamine & flavor contained polyphenolic flavonoids like rutin[67]. Meta-analysis studies showed that plant has the acitivity to reduce LDL, total cholesterol and glucose and also have the effect to increase the level of HDL[68].

Rubia cordifolia

It is herbaceous climbing plant with thin red bark, commonly known as majith, belonging to the family of Rubiaceae[69]. Tripathi YB et al found Rubiadin, dihydroxy anthraquinone, is main active chemical constituent has the antioxidant activity. Aq.root extract was found to have antidiabetic activity with lowering serum triglyceride level effect[70].

Jasminum grandiflorum

It is the medicinal plant commonly known as Cameli, Jati or Spanish jasmine, belonging to family Oleaceae (olive family) used to treat many ailments[71]. Phenolic compounds were active chemical constituents of dried flower bud has the antioxidant activity[72]. Ethanolic extract of flower has diabetic wound healing capacity by impairing angiogenesis. Further it has maximum wound breaking strength through maximizing collagen synthesis, that also showed antioxidant activity[73]. In vivo and *in vitro* studies showed an antioxidant and anti-inflammatory effect of methanolic extract of leaves[74].

Mangifera indica

Mangifera indica, commonly known as Mango, belongs to the family Anacardiaceae. Traditionally, parts of plants used to treat many ailments including diabetes type 2 to some extent[75]. there are so many active chemical constituents including polyphenols, phenolic acid and various flavonoids are reported as free radical scavenger activity[76]. Aq. Extract of leaves of plant has significant effect to reduce the high glucose level, when fed simultaneously only, it suggested that it may be due to hindrance in absorption of glucose in intestine[77]. It has been also reported that extract from acid hydrolysis has maximum antioxidant property as compared to the ethanolic extract[78].

DISCUSSION

In present study, we focused on medicinal plants which are very effective to treat Diabetes mellitus. Diabetes is increasing sharply to crisis point (also with obesity), with disastrous consequences including accelerated atherosclerosis, gangrene and limb amputation, kidney failure, neuropathy, and blindness. Adult brains need glucose as a supply of energy, and blood glucose monitoring represents the need to retain sufficient fuel supplies in the face of reduced food consumption and fluctuating metabolic demands. Extra calories are stored in the form of glycogen or fat. During fasting, the accumulated energy must be metabolized in a controlled way. Insulin is the most powerful regulatory hormone. Excess calories are stored as glycogen or fat. During fasting these energies stored need to be metabolized in a regulated manner. Insulin induction is stimulated by elevated blood glucose levels. Insulin release is influenced by the amount of glucose consumed. Insulin secretion is more efficient when glucose is given orally rather than intravenously. In diabetics, glucose has a lower ability to secrete insulin. The prevalent β cell is found in the center of individual islet, which is covered by a mantle of A cells, δ and PP cells. B cells also secrete islet amyloid polypeptide or amylin, a peptide that delays gastric emptying and inhibits insulin by inducing glycogen breakdown in striated muscle and C peptide. Glucagon works in opposition to insulin, increasing blood glucose and stimulating muscle protein breakdown. Insulin and glucagon synthesis is inhibited by somatostatin. It's widely spread outside the pancreas, and it's also released from the hypothalamus, inhibiting pituitary gland growth hormone release.

In diabetic patient there is an increment in mobilization of fatty acid i.e. lipolysis from adipose tissues to blood circulation that ultimately cause the high level of ketone bodies in circulation. As this fatty acids have both structural and functional requirements in every cell but in excess amount it may lead to disease that is treated by some medicinal plants like *Trigonella foenum-graecum*[45], *Eugenia jambolana*[38], *Lagenaria Siceraria*[31], *Carica papaya*[51].

In DM, high concentration of glucose may directly cause can directly make reactive oxygen species that lead to disparity among enzymatic and non-enzymatic antioxidant protection and also intensify the metabolism of glucose by a way of polyol (sorbitol) pathway as a result in increased production of reactive oxygen species (ROS). ROS can trigger low-density lipoprotein (LDL) oxidation, which results in the formation of ox-LDL, that is not identified by the LDL receptor but can also be grabbed by scavenger receptors in macrophages, contributing to foam cell formation, atherosclerotic plaques and complications of DM[79]. Some reports suggest that some plants like *Allium sepa*[8], *Azadirachta indica*[9], *Aegle marmeloscorrea*[13] having antioxidant ability to treat the complications of DM. Many medicinal plants are reported to have anti-diabetic activity but many of them have don't have much data regarding mechanism of action and clinical trial, therefore it is suggested that a lot of scope is there on medicinal plants for finding mechanism of action and further clinical trial for benefits of human being.

NOTE:

The study highlights the efficacy of "herbal medicine" which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

REFERENCES

- [1] C. M. Chetty, "Herbal Medicines for Diabetes Mellitus : A Review," vol. 2, no. 3, pp. 1883–1892, 2010.
- [2] P. Saeedi *et al.*, "Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045 : Results from the International Diabetes Federation Diabetes Atlas , 9 th edition," *Diabetes Res. Clin. Pract.*, vol. 157, p. 107843, 2019, doi: 10.1016/j.diabres.2019.107843.
- [3] K. Platel and K. Srinivasan, "Plant foods in the management of Diabetes mellitus: Vegetables as potential hypoglycaemic agents," *Nahrung - Food*, vol. 41, no. 2, pp. 68–74, 1997, doi: 10.1002/food.19970410203.
- [4] M. Jung, M. Park, H. Lee, Y.-H. Kang, E. Kang, and S. Kim, "Antidiabetic Agents from Medicinal Plants," *Curr. Med. Chem.*, vol. 13, no. 10, pp. 1203–1218, 2006, doi: 10.2174/092986706776360860.
- [5] C. Dwivedi and S. Daspaul, "Antidiabetic Herbal Drugs and Polyherbal Formulation Used For Diabetes: A Review," *J. Phytopharm. JPHYTO*, vol. 2, no. 23, pp. 44–51, 2013.
- [6] K. Bora and A. Sharma, "Phytoconstituents and therapeutic potential of allium cepa linn.- A Review," *Pharmacogn. Rev.*, vol. 3, no. 5, pp. 170–180, 2009.
- [7] M. Sajid, H. Akash, K. Rehman, and S. Chen, "SC," *Nutrition*, 2014, doi: 10.1016/j.nut.2014.02.011.
- [8] D. Jini and S. Sharmila, "Green synthesis of silver nanoparticles from Allium cepa and its in vitro antidiabetic activity," *Mater. Today Proc.*, vol. 22, no. xxxx, pp. 432–438, 2020, doi: 10.1016/j.matpr.2019.07.672.
- [9] C. Vergallo, E. Panzarini, and L. Dini, "High performance liquid chromatographic profiling of antioxidant and antidiabetic flavonoids purified from Azadirachta indica (neem) leaf ethanolic extract," *Pure Appl. Chem.*, vol. 91, no. 10, pp. 1631–1640, 2019, doi: 10.1515/pac-2018-1221.
- [10] N. Sriraksa, R. Kongsui, S. Thongrong, A. Duangjai, and T. Hawiset, "Effect of Azadirachta indica flower extract on functional recovery of sciatic nerve crush injury in rat models of DM," *Exp. Ther. Med.*, vol. 17, no. 1, pp. 541–550, 2019, doi: 10.3892/etm.2018.6931.
- [11] M. K. Gautam, M. Gangwar, S. K. Singh, and R. K. Goel, "Effects of Azadirachta indica on Vascular Endothelial Growth Factor and Cytokines in Diabetic Deep Wound," *Planta Med.*, vol. 81, no. 9, pp. 713–721, 2015, doi: 10.1055/s-0035-1545917.
- [12] A. Venthodika, N. Chhikara, S. Mann, M. K. Garg, S. A. Sofi, and A. Panghal, "Bioactive compounds of Aegle marmelos L., medicinal values and its food applications: A critical review," *Phyther. Res.*, no. March, pp. 1–21, 2020, doi: 10.1002/ptr.6934.

- [13] M. Fawzi Mahomoodally, A. Mollica, A. Stefanucci, M. Zakariyyah Aumeeruddy, R. Poorneeka, and G. Zengin, "Volatile components, pharmacological profile, and computational studies of essential oil from *Aegle marmelos* (Bael) leaves: A functional approach," *Ind. Crops Prod.*, vol. 126, no. August, pp. 13–21, 2018, doi: 10.1016/j.indcrop.2018.09.054.
- [14] S. N. Panaskar, M. M. Joglekar, S. S. Taklikar, V. S. Haldavnekar, and A. U. Arvindekar, "Aegle marmelos Correa leaf extract prevents secondary complications in streptozotocin-induced diabetic rats and demonstration of limonene as a potent antiglycating agent," *J. Pharm. Pharmacol.*, vol. 65, no. 6, pp. 884–894, 2013, doi: 10.1111/jphp.12044.
- [15] J. Younas and F. Hussain, "In vitro Antidiabetic Evaluation of *Allium sativum* L.," *Int. J. Chem. Biochem. Sci.*, vol. 5, pp. 22–25, 2014.
- [16] A. Ghorbani, "Phytotherapy for diabetic dyslipidemia: Evidence from clinical trials," *Clin. Lipidol.*, vol. 8, no. 3, pp. 311–319, 2013, doi: 10.2217/clp.13.26.
- [17] A. A. Maan *et al.*, "The therapeutic properties and applications of Aloe vera: A review," *J. Herb. Med.*, vol. 12, no. January, pp. 1–10, 2018, doi: 10.1016/j.hermed.2018.01.002.
- [18] M. Sánchez, E. González-Burgos, I. Iglesias, and M. P. Gómez-Serranillos, "Pharmacological update properties of aloe vera and its major active constituents," *Molecules*, vol. 25, no. 6, pp. 1–37, 2020, doi: 10.3390/molecules25061324.
- [19] M. S. Mokhele, D. Tswaledi, O. Aboyade, J. Shai, and D. Katerere, "Investigation of Aloe ferox leaf powder on anti-diabetes activity," *South African J. Bot.*, vol. 128, pp. 174–181, 2020, doi: 10.1016/j.sajb.2019.10.012.
- [20] D. T. Loots, M. Pieters, M. S. Islam, and L. Botes, "Antidiabetic effects of aloe ferox and aloe greatheadii var. Davyana leaf gel extracts in a low-Dose streptozotocin diabetes rat model," *S. Afr. J. Sci.*, vol. 107, no. 7–8, pp. 1–6, 2011, doi: 10.4102/sajs.v107i7/8.532.
- [21] P. Deepa, K. Sowndhararajan, S. Kim, and S. J. Park, "A role of Ficus species in the management of diabetes mellitus: A review," *J. Ethnopharmacol.*, vol. 215, pp. 210–232, 2018, doi: 10.1016/j.jep.2017.12.045.
- [22] N. Mehan *et al.*, "Phytopharmacology of *Ficus religiosa* L. and its significance as nanoparticulate carrier," *Ann. Phytomedicine An Int. J.*, vol. 8, no. 2, pp. 186–193, 2019, doi: 10.21276/ap.2019.8.2.24.
- [23] M. Fan, E. K. Kim, Y. J. Choi, Y. Tang, and S. H. Moon, "The role of momordica charantia in resisting obesity," *Int. J. Environ. Res. Public Health*, vol. 16, no. 18, 2019, doi: 10.3390/ijerph16183251.
- [24] L. Sun, X. Zhang, L. Dong, C. Zhang, P. Guo, and C. Wu, "The triterpenoids of the bitter gourd (*Momordica Charantia*) and their pharmacological activities: A review," *J. Food Compos. Anal.*, vol. 96, p. 103726, 2021, doi: 10.1016/j.jfca.2020.103726.
- [25] B. W. C. Roffey, A. S. Atwal, T. Johns, and S. Kubow, "Water extracts from *Momordica charantia* increase glucose uptake and adiponectin secretion in 3T3-L1 adipose cells," *J. Ethnopharmacol.*, vol. 112, no. 1, pp. 77–84, 2007, doi: 10.1016/j.jep.2007.02.003.
- [26] M. Bortolotti, D. Mercatelli, and L. Polito, "Momordica charantia, a nutraceutical approach for inflammatory related diseases," *Front. Pharmacol.*, vol. 10, no. MAY, pp. 1–9, 2019, doi: 10.3389/fphar.2019.00486.
- [27] S. Deshaware, S. Gupta, R. S. Singhal, M. Joshi, and P. S. Variyar, "Debittering of bitter gourd juice using β -cyclodextrin: Mechanism and effect on antidiabetic potential," *Food Chem.*, vol. 262, no. November 2017, pp. 78–85, 2018, doi: 10.1016/j.foodchem.2018.04.077.
- [28] S. Sharma, S. Kumar, C. Katare, and G. Prasad, "Plausible effect of bottle gourd

- (*Lagenaria siceraria*) pulp on glycemic status and lipid profile of the subjects with type II diabetes,” *World J. Pharm. Res. World J. Pharm. Res. SJIF Impact Factor Res. Artic. ISSN*, vol. 4045, no. 2, pp. 1426–1434, 2015.
- [29] U. A. Attar and S. G. Ghane, “In vitro antioxidant, antidiabetic, antiacetylcholine esterase, anticancer activities and RP-HPLC analysis of phenolics from the wild bottle gourd (*Lagenaria siceraria* (Molina) Standl.),” *South African J. Bot.*, vol. 125, pp. 360–370, 2019, doi: 10.1016/j.sajb.2019.08.004.
- [30] M. Fukushima, T. Matsuda, K. Yamagishi, and M. Nakano, “Comparative hypocholesterolemic effects of six dietary oils in cholesterol-fed rats after long-term feeding,” *Lipids*, vol. 32, no. 10, pp. 1069–1074, 1997, doi: 10.1007/s11745-997-0138-5.
- [31] C. Katare *et al.*, “Lipid-Lowering and Antioxidant Functions of Bottle Gourd (*Lagenaria siceraria*) Extract in Human Dyslipidemia,” *J. Evidence-Based Complement. Altern. Med.*, vol. 19, no. 2, pp. 112–118, 2014, doi: 10.1177/2156587214524229.
- [32] D. Ahmed and N. Ashiq, “In vitro analysis of anti-diabetic and anti-oxidative potential of pedicles of fruit-vegetable bottle gourd,” *Pak. J. Pharm. Sci.*, vol. 31, no. 6, pp. 2497–2501, 2018.
- [33] D. Singh and P. K. Chaudhuri, “A review on phytochemical and pharmacological properties of Holy basil (*Ocimum sanctum* L.),” *Ind. Crops Prod.*, vol. 118, no. March, pp. 367–382, 2018, doi: 10.1016/j.indcrop.2018.03.048.
- [34] S. Khairun Fadila, A. Chun Hui, K. Sook Mei, and C. Cheng Hock, “Chemical constituents and antioxidant capacity of *ocimum basilicum* and *ocimum sanctum*,” *Iran. J. Chem. Chem. Eng.*, vol. 38, no. 2, pp. 139–152, 2019.
- [35] P. Universiti, “Antidiabetic and In Vitro Enzyme Inhibition Studies of Methanol Extract of *Ocimum tenuiflorum* Linn Leaves and Its Fractions Authors : Leila Mousavi , Rabeta Mohd Salleh * and Vikneswaran Murugaiyah * Correspondence : rabetausm.my DOI : <https://doi.org/10>,” vol. 31, no. 1, 2020.
- [36] N. Jamshidi, C. Da Costa, and M. Cohen, “Holybasil (tulsi) lowers fasting glucose and improved lipid profile in adults with metabolic disease: A meta-analysis of randomized clinical trials,” *J. Funct. Foods*, vol. 45, no. March, pp. 47–57, 2018, doi: 10.1016/j.jff.2018.03.030.
- [37] M. S. Baliga, H. P. Bhat, B. R. V. Baliga, R. Wilson, and P. L. Palatty, “Phytochemistry, traditional uses and pharmacology of *Eugenia jambolana* Lam. (black plum): A review,” *Food Res. Int.*, vol. 44, no. 7, pp. 1776–1789, 2011, doi: 10.1016/j.foodres.2011.02.007.
- [38] K. Ravi, S. Rajasekaran, and S. Subramanian, “Antihyperlipidemic effect of *Eugenia jambolana* seed kernel on streptozotocin-induced diabetes in rats,” *Food Chem. Toxicol.*, vol. 43, no. 9, pp. 1433–1439, 2005, doi: 10.1016/j.fct.2005.04.004.
- [39] K. Ravi, S. Rajasekaran, and S. Subramanian, “Hypoglycemic effect of *Eugenia jambolana* seed kernels on streptozotocin-induced diabetes in rats,” *Pharm. Biol.*, vol. 41, no. 8, pp. 598–603, 2003, doi: 10.1080/13880200390501929.
- [40] K. Ravi, K. Sivagnanam, and S. Subramanian, “Anti-diabetic activity of *Eugenia jambolana* seed kernels on streptozotocin-induced diabetic rats,” *J. Med. Food*, vol. 7, no. 2, pp. 187–191, 2004, doi: 10.1089/1096620041224067.
- [41] M. Devgun, A. Nanda, and S. H. Ansari, “*Pterocarpus marsupium* Roxb. - A comprehensive review,” *Pharmacogn. Rev.*, vol. 3, no. 6, pp. 359–363, 2009.
- [42] A. Islam, L. Rebello, and S. Chepyala, “A Review of Anti-diabetic Activity of *Gymnema sylvestre* and *Pterocarpus marsupium* : Special Emphasis on its Combination in 4DM,” *Int. J. Nat. Life Sci.*, vol. 3, no. 2, pp. 40–51, 2019.

- [43] K. Begum, N. Begum, S. Reshma, W. Fatima, M. Ali, and A. Sultana, "Pterocarpus Marsupium Extraction and Evaluation for Diabetic Neuropathy," *Indo Am. J. Pharm. Sci.*, vol. 4, no. 07, pp. 1888–1897, 2017.
- [44] S. Chaudhary, P. S. Chaudhary, S. K. Chikara, M. C. Sharma, and M. Iriti, "Review on Fenugreek (*Trigonella foenum-graecum* L.) and its important secondary metabolite diosgenin," *Not. Bot. Horti Agrobot. Cluj-Napoca*, vol. 46, no. 1, pp. 22–31, 2018, doi: 10.15835/nbha46110996.
- [45] H. Zhang, J. Xu, M. Wang, X. Xia, R. Dai, and Y. Zhao, "Steroidal saponins and sapogenins from fenugreek and their inhibitory activity against α -glucosidase," *Steroids*, vol. 161, no. May, p. 108690, 2020, doi: 10.1016/j.steroids.2020.108690.
- [46] N. P. Lin and R. J. Chein, "Total synthesis and absolute structure of N55, a positive modulator of GLP-1 signaling," *Org. Biomol. Chem.*, vol. 18, no. 43, pp. 8899–8907, 2020, doi: 10.1039/d0ob01722a.
- [47] F. Khan *et al.*, "Comprehensive review on phytochemicals, pharmacological and clinical potentials of gymnema sylvestre," *Front. Pharmacol.*, vol. 10, no. OCT, 2019, doi: 10.3389/fphar.2019.01223.
- [48] Y. S. Oh, "Plant-derived compounds targeting pancreatic beta cells for the treatment of diabetes," *Evidence-based Complement. Altern. Med.*, vol. 2015, 2015, doi: 10.1155/2015/629863.
- [49] P. Kanetkar, R. Singhal, and M. Kamat, "Gymnema sylvestre: A memoir," *J. Clin. Biochem. Nutr.*, vol. 41, no. 2, pp. 77–81, 2007, doi: 10.3164/jcbrn.2007010.
- [50] R. S. Shenoy, K. V. H. Prashanth, and H. K. Manonmani, "In Vitro Antidiabetic Effects of Isolated Triterpene Glycoside Fraction from *Gymnema sylvestre*," *Evidence-based Complement. Altern. Med.*, vol. 2018, 2018, doi: 10.1155/2018/7154702.
- [51] I. E. Juárez-Rojop *et al.*, "Hypoglycemic effect of *Carica papaya* leaves in streptozotocin-induced diabetic rats," *BMC Complement. Altern. Med.*, vol. 12, 2012, doi: 10.1186/1472-6882-12-236.
- [52] I. E. Juárez-Rojop *et al.*, "Phytochemical screening and hypoglycemic activity of carica papaya leaf in streptozotocin-induced diabetic rats," *Brazilian J. Pharmacogn.*, vol. 24, no. 3, pp. 341–347, 2014, doi: 10.1016/j.bjp.2014.07.012.
- [53] K. Krishnan, L. E. Mathew, N. R. Vijayalakshmi, and A. Helen, "Anti-inflammatory potential of β -amyrin, a triterpenoid isolated from *Costus igneus*," *Inflammopharmacology*, vol. 22, no. 6, pp. 373–385, 2014, doi: 10.1007/s10787-014-0218-8.
- [54] I. S. Costa, A. F. Medeiros, G. Piuvezam, G. C. B. S. Medeiros, B. L. L. Maciel, and A. H. A. Morais, "Insulin-like proteins in plant sources: A systematic review," *Diabetes, Metab. Syndr. Obes. Targets Ther.*, vol. 13, pp. 3421–3431, 2020, doi: 10.2147/DMSO.S256883.
- [55] M. R. Hardikar, M. E. Varma, A. A. Kulkarni, P. P. Kulkarni, and B. N. Joshi, "Elucidation of hypoglycemic action and toxicity studies of insulin-like protein from *Costus igneus*," *Phytochemistry*, vol. 124, pp. 99–107, 2016, doi: 10.1016/j.phytochem.2016.02.001.
- [56] A. Aruna, "Synthesis and Characterization of Silver Nanoparticles of Insulin Plant (*Costus pictus* D. Don) leaves," *Asian J. Biomed. Pharm. Sci.*, vol. 4, no. 34, pp. 1–6, 2014, doi: 10.15272/ajbps.v4i34.523.
- [57] V. Vinotha *et al.*, "Synthesis of ZnO nanoparticles using insulin-rich leaf extract: Anti-diabetic, antibiofilm and anti-oxidant properties," *J. Photochem. Photobiol. B Biol.*, vol. 197, no. May, p. 111541, 2019, doi: 10.1016/j.jphotobiol.2019.111541.

- [58] M. Dorri, S. Hashemitabar, and H. Hosseinzadeh, "Cinnamon (*Cinnamomum zeylanicum*) as an antidote or a protective agent against natural or chemical toxicities: a review," *Drug Chem. Toxicol.*, vol. 41, no. 3, pp. 338–351, 2018, doi: 10.1080/01480545.2017.1417995.
- [59] W. A. M. Niroshani Wariyapperuma, S. Kannangara, Y. S. Wijayasinghe, S. Subramaniam, and B. Jayawardena, "In vitro anti-diabetic effects and phytochemical profiling of novel varieties of *Cinnamomum zeylanicum* (L.) extracts," *PeerJ*, vol. 8, 2020, doi: 10.7717/peerj.10070.
- [60] P. Ranasinghe, S. Pigera, G. S. Premakumara, P. Galappaththy, G. R. Constantine, and P. Katulanda, "Medicinal properties of 'true' cinnamon (*Cinnamomum zeylanicum*): A systematic review," *BMC Complement. Altern. Med.*, vol. 13, no. 1, p. 1, 2013, doi: 10.1186/1472-6882-13-275.
- [61] A. K. Ghosh, S. Banerjee, H. I. Mullick, and J. Banerjee, "Zingiber officinale: A natural gold," *Int. J. Pharma Bio Sci.*, vol. 2, no. 1, pp. 283–294, 2011.
- [62] G. Singh, I. P. S. Kapoor, P. Singh, C. S. de Heluani, M. P. de Lampasona, and C. A. N. Catalan, "Chemistry, antioxidant and antimicrobial investigations on essential oil and oleoresins of *Zingiber officinale*," *Food Chem. Toxicol.*, vol. 46, no. 10, pp. 3295–3302, 2008, doi: 10.1016/j.fct.2008.07.017.
- [63] P. Hasimun, I. K. Adnyana, R. Valentina, and E. Lisnasari, "Potential alpha-glucosidase inhibitor from selected zingiberaceae family," *Asian J. Pharm. Clin. Res.*, vol. 9, no. 1, pp. 141–144, 2016.
- [64] B. H. Ali, G. Blunden, M. O. Tanira, and A. Nemmar, "Some phytochemical, pharmacological and toxicological properties of ginger (*Zingiber officinale* Roscoe): A review of recent research," *Food Chem. Toxicol.*, vol. 46, no. 2, pp. 409–420, 2008, doi: 10.1016/j.fct.2007.09.085.
- [65] G. C. N. Carvalho, J. C. G. Lira-Neto, M. F. M. de Araújo, R. W. J. F. de Freitas, M. L. Zanetti, and M. M. C. Damasceno, "Effectiveness of ginger in reducing metabolic levels in people with diabetes: a randomized clinical trial," *Rev. Lat. Am. Enfermagem*, vol. 28, p. e3369, 2020, doi: 10.1590/1518-8345.3870.3369.
- [66] N. K. Saini and J. Biosys, "Available online through," no. May, 2014.
- [67] D. Potdar, R. R. Hirwani, and S. Dhulap, "Phyto-chemical and pharmacological applications of *Berberis aristata*," *Fitoterapia*, vol. 83, no. 5, pp. 817–830, 2012, doi: 10.1016/j.fitote.2012.04.012.
- [68] B. Roshanravan *et al.*, "The effects of *Berberis vulgaris* L. and *Berberis aristata* L. in metabolic syndrome patients: a systematic and meta-analysis study," *Arch. Physiol. Biochem.*, vol. 0, no. 0, pp. 1–12, 2020, doi: 10.1080/13813455.2020.1828482.
- [69] P. S. Monitor, "A GOLD ORNAMENTAL PLANT – RUBIA CORDIFOLIA :A REVIEW Komal A. Kale*, Sonia J. Patell, Ankita B. Chaudhary, Nehal K. Gohil," vol. 8, no. 2, pp. 157–164, 2017.
- [70] R. Baskar *et al.*, "Antihyperglycemic activity of aqueous root extract of *Rubia cordifolia* in streptozotocin-induced diabetic rats," *Pharm. Biol.*, vol. 44, no. 6, pp. 475–479, 2006, doi: 10.1080/13880200600798593.
- [71] P. Sciences, H. Science, and T. Nadu, "JASMINUM GRANDIFLORUM LINN. – AN UPDATE REVIEW P. Rajasri Bharathi, Shubashini K. Sripathi * and A. Naga Lakshmi Department of Chemistry, Avinashilingam Institute for Home Science and Higher Education for Women, Coimbatore - 641043, Tamil Nadu, India.," vol. 11, no. 5, pp. 1994–2010, 2020, doi: 10.13040/IJPSR.0975-8232.11(5).1994-10.
- [72] F. Ferreres, C. Grosso, A. Gil-Izquierdo, P. Valentão, and P. B. Andrade, "Assessing

- Jasminum grandiflorum L. authenticity by HPLC-DAD-ESI/MSn and effects on physiological enzymes and oxidative species,” *J. Pharm. Biomed. Anal.*, vol. 88, pp. 157–161, 2014, doi: 10.1016/j.jpba.2013.08.040.
- [73] H. Hirapara, V. Ghorl, A. Anovadiya, S. Baxi, and C. Tripathi, “Effects of ethanolic extract of *Jasminum grandiflorum* Linn. flowers on wound healing in diabetic Wistar albino rats.,” *Avicenna J. phytomedicine*, vol. 7, no. 5, pp. 401–408, 2017, doi: 10.22038/ajp.2017.14406.1580.
- [74] A. P. Chaturvedi and Y. B. Tripathi, “Methanolic extract of leaves of *Jasminum grandiflorum* Linn modulates oxidative stress and inflammatory mediators,” *Inflammopharmacology*, vol. 19, no. 5, pp. 273–281, 2011, doi: 10.1007/s10787-011-0087-3.
- [75] K. Shah, M. Patel, R. Patel, and P. Parmar, “*Mangifera Indica* (Mango),” *Pharmacogn. Rev.*, vol. 4, no. 7, pp. 42–48, 2010, doi: 10.4103/0973-7847.65325.
- [76] R. J. Nijveldt, E. van Nood, D. E. van Hoorn, P. G. Boelens, K. van Norren, and P. A. van Leeuwen, “Flavonoids a review of probable mechanisms of action,” *Am J Clin Nutr*, vol. 74, no. 4, pp. 418–425, 2012.
- [77] A. O. Aderibigbe, T. S. Emudianughe, and B. A. S. Lawal, “Antihyperglycaemic effect of *Mangifera indica* in rat,” *Phyther. Res.*, vol. 13, no. 6, pp. 504–507, 1999, doi: 10.1002/(SICI)1099-1573(199909)13:6<504::AID-PTR533>3.0.CO;2-9.
- [78] M. K. Ediriweera, K. H. Tennekoon, and S. R. Samarakoon, “A Review on Ethnopharmacological Applications, Pharmacological Activities, and Bioactive Compounds of *Mangifera indica* (Mango),” *Evidence-based Complement. Altern. Med.*, vol. 2017, 2017, doi: 10.1155/2017/6949835.
- [79] N. A. Khalaf, A. K. Shakya, A. Al-Othman, Z. El-Agbar, and H. Farah, “Antioxidant activity of some common plants,” *Turkish J. Biol.*, vol. 32, no. 1, pp. 51–55, 2008.