

Nutritional Agonists of (PPAR)- γ : An Immunomodulatory approach to control cytokine storm in covid19 patients

ABSTRACT

Recent examinations express that multi organ failure is seen in Corona virus infected patients with different pathway. It has been shown in contemplates that increased levels of cytokines like IL-1B and INF gamma were observed. It is called as cytokine storm with higher convergences of CCL2 and CXCL10. The cytokine storm is trailed by our immune system attacking own body which thus may cause numerous organ abnormalities and conclusive outcome being death. There is currently no specific treatment for viral illness, and this methodology is an optional path for focusing on specific qualities that may diminish cytokine storm. In such manner peroxisome proliferators activated receptors (PPARs) have a place with group of transcription factors which are known to manage the inflammatory mechanisms in body. This immunomodulatory approach is intended for focusing on the nutritional PPAR-gamma ligands that can be found in food, further Molecular docking studies were performed. The activation or increased expression levels of PPAR gamma because of chosen agonists may reduce the cytokine storm in the covid patients. Thus, this is one such fascinating way to deal with neutralization of the cytokines exorbitantly elevated by use of substances like pomegranate, lemon grass and so on to activate PPARs reliably.

Keywords: PPAR, Corona virus, Cytokine storm, Covid, Cytokines, Immune system.

Introduction

The higher rate of mortality due to the corona virus outbreak has become a major threat to public health and number of cases is expected to grow exponentially [1]. Mononuclear inflammatory cells accumulation, pneumocyte hyperplasia and alveolar damage are observed in patients infected severely with corona virus [2]. There are many mechanisms that are associated with the deterioration of the viral disease and one of it is the cytokine storm that causes threat to the patients

[3]. The cytokine storm is due to the sudden outburst of the pro-inflammatory components [4]. The modulation of the aggressive release of the cytokines is important to reduce the multi-organ dysfunction in corona virus infected patients. Though many of the infected patients show mild symptoms nearly 20% need hospitalization and 7% of patients needed intensive care [5]. When the virus replicates inside the cells, ARDS acute respiratory distress syndrome is developed [6]. Similarly other problems like organ failure, kidney injury and sepsis are prevailed. The detailed examinations of studies and reports showed that it is due to elevated levels of cytokines like interferon's and interleukin's [7]. When there is no known cure for corona virus infection, it is critical to investigate all other treatment options [8]. This article majorly focuses on the cytokine storm and its modulation by nutritional agonists. The remarkable role of PPAR-gamma increased expression also cooperates in reducing the cytokines risk [9].

Cytokine storm and its affects in covid patients

Cytokines are small secreted proteins that are discharged by cells with the end goal of intercellular communication and signaling [10]. Such cytokines are redundant in their activities, this means activities like autocrine, paracrine, endocrine movement and, through receptor binding, can bring out many responses, depending upon the target cell [11]. The control over cell expansion and differentiation are the major functions of cytokines. Prior to inspecting the effects of cytokine storms, it is significant to investigate the cytokines at the core of the cytokine storm. Severe lung diseases are considered best examples of severe lung diseases, in which nearby inflammation gushes out over into the circulatory system, delivering sepsis, as characterized by relentless hypotension, hyper-or hypothermia [12]. Viral, bacterial, and contagious aspiratory diseases all leads to the sepsis disorder and these etiological agents are hard to separate on clinical grounds [13]. Sometimes, cytokine storms in relation to tissue damage without microbial infection can also be persevered. Not with-standing lung infection, the cytokine storm is an outcome of extreme diseases in the gastrointestinal region, urinary tract and nervous system [14].

Patients with extreme sepsis show plasma cytokine profiles, which change after some time because of aspiratory or non-pulmonary infections. The intense reaction cytokines TNF and IL-1 β and the chemotactic cytokines IL-8 and MCP-1 show up in the early minutes to hours after infection, trailed by a more continued increment in IL-6 [15]. The calming cytokine IL-10 shows up to some degree later, as the body endeavors to control the acute inflammation [16]. Infectious

diseases remain as a severe threat all over the world accounting for increasing number of mortalities rates [17]. Against this scenery of antimicrobial resistance and the rise of new microbes, expanding interest has turned towards the advancement of medications that target on the immune reaction to infection and it is logical to target this response in order to cut down the self-inflicted damage initiated by host cells [18].

Role of PPAR- γ :

The activation and working of PPARs require the heterodimerization with another nuclear receptor, the retinoid X receptor (RXR). The RXR family has three diverse isoforms (RXR- α , - β , - γ) that are initiated by endogenous 9-cis retinoic acid. Upon ligand binding, the PPAR-RXR complex moves into the nucleus, where it binds to the PPRE elements of target DNA [19-23]. The relationship with the ligand initiates conformational changes, co-repressor separation and co-activator association, prompting ligand-initiated trans-activation [24].




Fig 1: PPAR-G and RXR signalling on binding of ligand

Nutritional agonists Sources of PPAR- γ

Turmeric is derived from the plant *Curcuma longa* [25] primarily used as a food ingredient, but in recent years it has also been used as a treatment for inflammatory-related problems [26]. Curcumin, the phytochemical component of turmeric, can control the transcription factor expression, cytokines and enzymes related to inflammation [27]. The anti-inflammatory property of curcumin mediates the PPAR gamma gene, which therefore inhibits TNF- α , a cytokine associated with inflammation. The mechanism by which curcumin controls PPAR- γ has not yet been explained [28]. It was known that, curcumin instigates an anti-inflammatory effect through the up-regulation of PPAR- γ , which leads to the inhibition of NF- κ B, a favorable inflammatory mediator [29]. Rosemary and sage are spices that are fundamentally found in the Mediterranean zone; they are utilized in the kitchen as fragrant plants, yet before, they have been additionally utilized for their antiseptic and anti-inflammatory properties [30]. They contain dipertenoids, like carnosic acid and carnosol, which have been proved in having anti-inflammatory effects. In reality, carnosic acid and carnosol involve with the key pathways of inflammation, hindering the activation of NF- κ B and the production of COX-2, eicosanoids, IL-1 β , and TNF- α . Moreover, carnosic acid and carnosol have been appeared to have agonistic impacts, consequently initiating PPAR- γ and its expression [31]. Lemongrass is a perennial herb utilized for the most part for the extraction of its oil, which is utilized for aroma in the kitchen and as a scent in beautifiers. Lemongrass has been utilized for quite a long time, particularly in Southeast Asia, for its pain relieving and anti-inflammatory properties [32]. Lemongrass oil contains citral, which gives it the particular smell and scent and is a combination of aldehydes that have the same molecular formula yet different structure, the trans isomer geranial and the cis isomer neral. Citral activates PPAR- γ , which represses the expression of cytokines, for example, IL-1 β and IL-6, furthermore, suppresses the action of the COX-2 promotor [33]. Hot pepper is a member from the Solanaceae family; it is the widest spread spice on the planet, also, it is commonly utilized in cooking [34]. It contains capsaicin, the most present compound in hot pepper, which, along with different capsaicinoids, is responsible for the spicy taste [35]. Capsaicinoids have antifungal and antibacterial properties and have been utilized to treat joint pains. Capsaicin has anti-inflammatory properties. An examination has demonstrated that it represses the lipopolysaccharide mediated production of TNF- α , a provocative cytokine, by initiating PPAR- γ [36]. Pomegranate, a fruit grown primarily in Southeast Asia, the Mediterranean region, and in the USA [37]. It has been utilized for many years as a restorative natural product for its possible

valuable properties; actually, the different parts of the organic product appear to contain antimicrobial, antifungal, cancer prevention agent, and anti-inflammatory properties [38]. Specifically, pomegranate seed oil contains puniceic acid, which is a conjugated alpha-linoleic acid with anti-inflammatory and immunomodulatory capability. Puniceic acid can inhibit expression of IL-6, IL-8, IL-12, and TNF- α , by modulating PPAR-gamma, which restrains the expression of the NF- κ B pathway [39]. Polyunsaturated unsaturated fats are identified with the regulation of inflammation; specifically, they are responsible for the reduction of the pro-inflammatory cytokines by activating PPAR- γ [40]. Docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA) are contained in ocean food and fish oil, and they are engaged with the inhibition of various aspects like leucocytes chemotaxis, eicosanoid production. Their cooperation with PPAR- γ prompts the hindrance of NF- κ B, a key factor of pro-inflammatory cytokines [41].




Fig 2: The activation of PPAR- γ by above mentioned agonists showing regulatory role on cytokine storm. The pro-inflammatory cytokines released by macrophages and lung endothelial cells induce immune responses in turn PPAR- γ prevents over production of cytokines becoming target for immunomodulatory therapies in viral infections.

MATERIALS AND METHODS

SwissDock is a docking web service platform in which the structure of the target protein, just as that of the ligand, can be automatically prepared for docking. Moreover, the bulky process of the docking system is hidden behind web interface giving sensible parameters as input records. All calculations are performed on the worker side, so that docking runs don't need any computational force from the client. The understanding of results and their interpretation into existing examination is encouraged by the consistent perception using UCSF Chimera. UCSF Chimera: It is a molecular visualization program that helps and coordinates for the analysis of molecular structures and data. By using this online service high quality animations can be generated. The development is supported by National Institute of Health. Molinspiration for ADME analysis and PROCHECK

Molecular Docking

Selection of target protein: In the target selection port within swiss dock PDB code or protein name is given as input. If the selected target is unavailable then the PDB structure file can be uploaded. Selection of ligand: Ligand name is given as input and it retrieves the structure from zinc library or input can be Zinc Accession Number. If the ligand is unavailable then ligand can be downloaded from DrugBank or pubchem.

The image shows a screenshot of the SwissDock web interface. It is divided into two main sections: 'Target selection' and 'Ligand selection'. Each section has a search input field and a 'Search' button. Below the search fields, there are instructions on what can be entered: 'ie. PDB code, protein name, sequence, or URL or upload file (max 5MB)' for targets, and 'ie. ZINC AC, ligand name or category (like scaffolds or sidechains), or URL or upload file (max 5MB)' for ligands. A large 'UNIVERSITY' watermark is visible across the page.

Target selection

Search for targets:

Search

ie. PDB code, protein name, sequence, or URL
or upload file (max 5MB)

Ligand selection

Search for ligands:

Search

ie. ZINC AC, ligand name or category (like scaffolds or sidechains), or URL
or upload file (max 5MB)

Fig. 3: Target and ligand selection in SwissDock




Fig. 4: Structure of PPAR-G complexed with ligand (Source PDB)

RESULTS AND DISCUSSION

Chart 1. Analysis of results using Chimera (Binding energy assessment)

Curcumin

Remark Fullfitness	-3050.3596
H Bonds	1
deltaG	-8.782826
deltaGvdw	-53.0341

Carnasol

Remark Fullfitness	-3111.4092
H Bonds	3
deltaG	-9.008968
deltaGvdw	-66.4635

Citral

Remark Fullfitness	-3077.2585
H Bonds	1
deltaG	-6.7350316
deltaGvdw	-25.7825

Capsaicin

Remark Fullfitness	-3083.7202
H Bonds	2
deltaG	-8.320375
deltaGvdw	-47.0803

Ecosapentanoic acid

Remark Fullfitness	-3030.5256
H Bonds	1
deltaG	-6.6861825
deltaGvdw	-27.4205




Fig. 5: Chimera based 3D image obtained after docking with curcumin




Fig. 6: Chimera based 3D image obtained after docking with carnosol




Fig. 7: Chimera based 3D image obtained after docking with citral




Fig. 8: Chimera based 3D image obtained after docking with capsaicin




Fig. 9: Chimera based 3D image obtained after docking with EPA

On obtaining results from five selected ligands the best deltaG value is recorded for carnosol (-9.008968) with 3 hydrogen bonds and vanderwaal forces energy was recorded around -66.4635. Among curcumin, carnosol, citral, capsaicin and EPA; carnosol is assumed to have best agonistic property for PPAR-gamma and EPA is recorded with lowest ability of agonism.

Table 1. Absorption, Distribution, Metabolism, and Excretion (ADME) Prediction by Molinspiration:

Name of ligand	Mol.Wt	nAtoms	N Violations	volume
Curcumin	368.38	27	0	332.18
Carnosol	302.12	20	0	331.12
Citral	152.24	11	0	169.74
Capsaicin	305.42	22	0	310.37
EPA	302.46	22	1	327.70

Since many drug related projects come up short during clinical preliminaries because of poor ADME properties, it is a needy practice to do ADME tests at the beginning phases. Different exploratory and computational techniques have been done to get ADME properties in an affordable way as far as time and cost. As in vitro and in vivo exploratory information on ADME have accumulated, the precision of in silico models in ADME increases and in this manner, numerous in silico models are currently broadly utilized in drug research. In light of the requests from drug discovery analysts, the advancement of in silico models in ADME has become more dynamic. In silico ADME properties of the selected molecules were predicted using Molinspiration. Lipinski's rule stating the properties and their ranges molecular weight < to be less than 500, number of hydrogen bond donors < 5, number of hydrogen bond acceptors < 10, violations and total volume was also assessed, which helped in distinguishing between drug-like or non-drug like profiles of the selected candidates. In this regard of given constraints and standard values, Citral is observed to have less molecular weight (152.24) and curcumin with very high molecular weight (368.38).

Ramachandran Plot and statistical values:




Fig. 10: Ramachandran plot obtained using PROCHECK

Table 2: The shaded region on the plot "core" regions represents the most favorable combinations of phi-psi values. Ideally, one would hope to have over 90% of the residues in these "core" regions. The percentage of residues in the "core" regions is one of the better guides to stereochemical quality. The percentage of residues in most favorable region is around 99.8% only 0.2% residues are present in generously allowed regions. 0% of disallowed regions is observed from graph.

Residues in most favorable regions	416	90.08%
Residues in additional allowed regions	41	9.0%
Residues in generously allowed regions	1	0.2%
Residues in disallowed regions	0	0.0%
Number of non-glycine and non-proline residues	458	100%
Number of end residues	12	
Number of glycine residues	22	
Number of proline residues	18	
Total	510	

CONCLUSION

Currently the newly invaded corona virus infection represents a worldwide health problem. This made the national clinical systems go through stress. Somehow recent studies also proved the relation between the cytokine storm and hospitalization of patients infected with the corona virus. In this regard finding nutritional agonists for PPAR-gamma could be effective and best therapeutic strategy to reduce the elevated levels of cytokines. Characteristic nutritional compounds are proved as rich hotspot for the discovering novel PPAR-gamma ligands and many of them are basically assorted agonists from medicinal plants and from food products. Interesting activation pathways and insilico approaches gives us the clear understanding of capabilities that nutritional ligands can have as agonists for increasing the expression of PPAR-gamma. The food products and compounds mentioned in previous sections would stand as the valid option for modulating the immune system mechanisms thereby reduce cytokine storm.

REFERENCES

1. Preysi C, Navreet K. S, Srinivasan G, Heena D. COVID-19 in India- Impact and Mitigation Strategies. *Biomed Pharmacol J* 2021;14(2).
2. Tian S., Hu W., Niu L., Liu H., Xu H., Xiao S.-Y. Pulmonary Pathology of Early-Phase 2019 Novel Coronavirus (COVID-19) Pneumonia in Two Patients With Lung Cancer. *J. Thorac. Oncol.* 2020 doi: 10.1016/j.jtho.2020.02.010.
3. Kumar VP, Mounica MS, Swetha TN, Lakshmi GV, Anusri T, Amulya PL, Sirisha A. Cytokine therapy: clinical and progressive status in cancer. *International Journal of Immunological Studies.* 2011 Jan 1;1(3):264-71.
4. Yiu HH, Graham AL, Stengel RF. Dynamics of a cytokine storm. *PloS one.* 2012 Oct 1;7(10):e45027.
5. Liu PP, Blet A, Smyth D, Li H. The science underlying COVID-19: implications for the cardiovascular system. *Circulation.* 2020 Apr 15.

6. Cockrell AS, Yount BL, Scobey T, Jensen K, Douglas M, Beall A, Tang XC, Marasco WA, Heise MT, Baric RS. A mouse model for MERS coronavirus-induced acute respiratory distress syndrome. *Nature microbiology*. 2016 Nov 28;2(2):1-1.
7. Sinha P, Matthay MA, Calfee CS. Is a “cytokine storm” relevant to COVID-19?. *JAMA internal medicine*. 2020 Sep 1;180(9):1152-4.
8. Ciavarella C, Motta I, Valente S, Pasquinelli G. Pharmacological (or Synthetic) and Nutritional Agonists of PPAR- γ as Candidates for Cytokine Storm Modulation in COVID-19 Disease. *Molecules*. 2020 Jan;25(9):2076.
9. Christodoulides C, Vidal-Puig A. PPARs and adipocyte function. *Molecular and cellular endocrinology*. 2010 Apr 29;318(1-2):61-8.
10. Stow JL, Murray RZ. Intracellular trafficking and secretion of inflammatory cytokines. *Cytokine & growth factor reviews*. 2013 Jun 1;24(3):227-39.
11. Altan-Bonnet G, Mukherjee R. Cytokine-mediated communication: a quantitative appraisal of immune complexity. *Nature reviews Immunology*. 2019 Apr;19(4):205-17.
12. Mangalmurti N, Hunter CA. Cytokine storms: understanding COVID-19. *Immunity*. 2020 Jun 28.
13. Baharoon S, Al-Jahdali H, Al Hashmi J, Memish ZA, Ahmed QA. Severe sepsis and septic shock at the Hajj: etiologies and outcomes. *Travel Medicine and Infectious Disease*. 2009 Jul 1;7(4):247-52.
14. Opoku A, Antwi JS, Danso JO, Mensah ON, Baffoe-Sarpong K, Yiadom AK, Owusu NA, Twene P, Mensah RS. Dimensions of Pathophysiology of COVID 19 on the Body Systems and Its Implication for Investigations, Treatment and Further Studies. *Journal of Advances in Medicine and Medical Research*. 2020 Jul 6:11-21.
15. Wong CK, Lam CW, Wu AK, Ip WK, Lee NL, Chan IH, Lit LC, Hui DS, Chan MH, Chung SS, Sung JJ. Plasma inflammatory cytokines and chemokines in severe acute respiratory syndrome. *Clinical & Experimental Immunology*. 2004 Apr;136(1):95-103.

16. Kao CC. HIV-1 Envelope Glycoprotein 120 Induced Neuroinflammation Does Not Result in Neurocognitive Impairment in An in vivo Rat Model (Doctoral dissertation).
17. Becker K, Hu Y, Biller-Andorno N. Infectious diseases—a global challenge. *International Journal of Medical Microbiology*. 2006 Aug 1;296(4-5):179-85.
18. Montanaro L, Campoccia D, Arciola CR. Advancements in molecular epidemiology of implant infections and future perspectives. *Biomaterials*. 2007 Dec 1;28(34):5155-68.
19. Subbarayan V, Sabichi AL, Kim J, Llansa N, Logothetis CJ, Lippman SM, Menter DG. Differential peroxisome proliferator-activated receptor- γ isoform expression and agonist effects in normal and malignant prostate cells. *Cancer Epidemiology and Prevention Biomarkers*. 2004 Nov 1;13(11):1710-6.
20. Shao D, Rangwala SM, Bailey ST, Krakow SL, Reginato MJ, Lazar MA. Interdomain communication regulating ligand binding by PPAR- γ . *Nature*. 1998 Nov;396(6709):377-80.
21. Perez A, Kastner P, Sethi S, Lutz Y, Reibel C, Chambon P. PMLRAR homodimers: distinct DNA binding properties and heteromeric interactions with RXR. *The EMBO journal*. 1993 Aug;12(8):3171-82.
22. Kumar VP, Chakrapani R, Prasad MK. Cancer Immune System: A focus on clinical trails of Cancer Vaccines. *Journal of Pure and Applied Microbiology*. 2011 Apr 1;5(1):441-4.
23. De Lera AR, Bourguet W, Altucci L, Gronemeyer H. Design of selective nuclear receptor modulators: RAR and RXR as a case study. *Nature reviews Drug discovery*. 2007 Oct;6(10):811-20.
24. Tian T, Li A, Lu H, Luo R, Zhang M, Li Z. TAZ promotes temozolomide resistance by upregulating MCL-1 in human glioma cells. *Biochemical and biophysical research communications*. 2015 Aug 7;463(4):638-43.
25. Kocaadam B, Şanlıer N. Curcumin, an active component of turmeric (*Curcuma longa*), and its effects on health. *Critical reviews in food science and nutrition*. 2017 Sep 2;57(13):2889-95.

26. Pastorino G, Cornara L, Soares S, Rodrigues F, Oliveira MB. Liquorice (Glycyrrhizaglabra): A phytochemical and pharmacological review. *Phytotherapy research*. 2018 Dec;32(12):2323-39.
27. Ghosh S, Banerjee S, Sil PC. The beneficial role of curcumin on inflammation, diabetes and neurodegenerative disease: A recent update. *Food and Chemical Toxicology*. 2015 Sep 1;83:111-24.
28. Ciavarella C, Motta I, Valente S, Pasquinelli G. Pharmacological (or Synthetic) and Nutritional Agonists of PPAR- γ as Candidates for Cytokine Storm Modulation in COVID-19 Disease. *Molecules*. 2020 Jan;25(9):2076.
29. Arora S, Dev K, Agarwal B, Das P, Syed MA. Macrophages: their role, activation and polarization in pulmonary diseases. *Immunobiology*. 2018 Apr 1;223(4-5):383-96.
30. Sharangi AB, Guha S. Wonders of leafy spices: Medicinal properties ensuring Human Health. *Science International*. 2013;1(9):312-7.
31. Ciavarella C, Motta I, Valente S, Pasquinelli G. Pharmacological (or Synthetic) and Nutritional Agonists of PPAR- γ as Candidates for Cytokine Storm Modulation in COVID-19 Disease. *Molecules*. 2020 Jan;25(9):2076.
32. Campos CA, Lima BS, Trindade GG, Souza EP, Mota DS, Heimfarth L, Quintans JS, Quintans-Júnior LJ, Sussuchi EM, Sarmiento VH, Carvalho FM. Anti-hyperalgesic and anti-inflammatory effects of citral with β -cyclodextrin and hydroxypropyl- β -cyclodextrin inclusion complexes in animal models. *Life sciences*. 2019 Jul 15;229:139-48.
33. Shen Y, Sun Z, Guo X. Citral inhibits lipopolysaccharide-induced acute lung injury by activating PPAR- γ . *European journal of pharmacology*. 2015 Jan 15;747:45-51.
34. Omolo MA, Wong ZZ, Mergen K, Hastings JC, Le NC, Reil HA, Case KA, Baumler DJ. Antimicrobial properties of chili peppers. *Journal of Infectious Diseases and Therapy*. 2014 Jun 6.
35. Reyes-Escogido MD, Gonzalez-Mondragon EG, Vazquez-Tzompantzi E. Chemical and pharmacological aspects of capsaicin. *Molecules*. 2011 Feb;16(2):1253-70.

36. Wu LC, Fan NC, Lin MH, Chu IR, Huang SJ, Hu CY, Han SY. Anti-inflammatory effect of spilanthol from *Spilanthescmella* on murine macrophage by down-regulating LPS-induced inflammatory mediators. *Journal of agricultural and food chemistry*. 2008 Apr 9;56(7):2341-9.
37. Stover ED, Mercure EW. The pomegranate: a new look at the fruit of paradise. *HortScience*. 2007 Aug 1;42(5):1088-92.
38. Boddakayala N, Bodiga S, Bodiga VL, Tummala PK, Vemuru S, Setti R, Manne AA, Veeravilli S, Vemuri PK. Conformational Epitope Prediction of Birch Betv 1 and Hazel Cor A1 Towards B-Cells. *Journal of Pharmaceutical Research International*. 2021 May 5:146-59.
39. Klepsch V, Moschen AR, Tilg H, Baier G, Hermann-Kleiter N. Nuclear receptors regulate intestinal inflammation in the context of IBD. *Frontiers in Immunology*. 2019 May 14;10:1070.
40. Daynes RA, Jones DC. Emerging roles of PPARs in inflammation and immunity. *Nature Reviews Immunology*. 2002 Oct;2(10):748-59.
41. Calder PC. Omega-3 polyunsaturated fatty acids and inflammatory processes: nutrition or pharmacology?. *British journal of clinical pharmacology*. 2013 Mar;75(3):645-62.