

A Review on Cardioprotective mechanism of Chemical Constituents of Medicinal Plant

Abstract:

Plants are major source of human living. Since the beginning of the era plants have been used for medicinal purposes. There is dire to explore the mechanism of chemical constituents in plants and particularly saponins and cardiac glycosides due to their mechanism to save damaged cells in cardiac muscle. Databases like Google scholar, Medline, PubMed, and the Directory of Open Access Journals were searched to find the articles describing the cardioprotective mechanism of medicinal plants. Saponin, flavonoids, glycoside, steroid, alkaloids, tannin, phenol, phlobatanin, terpenoids, and anthraquinone are chemical constituents in plants enhances cardio protection activity and decreases cardiac abnormalities. The current review article provides data on the use of medicinal plants, specifically against cardiac diseases, as well as an investigation of molecules/phytoconstituents as plant secondary metabolites for their cardioprotective potential.

Keywords: Cardioprotective, Medicinal Plants, Oxidative Stress, Saponins, Cardiac Glycosides

1. Introduction:

Traditional medicines are knowledge, skills and continuous practice which depends on theories, beliefs of different cultures in various countries used to maintain good health, prevention, diagnosis of both physical and mental health [1]. Use of Medicinal plants are being followed since ages because, therapeutic effects can be examined and evaluation of toxic doses can be done to maintain quality of formulation [2]. Medicinal plants are major resource base for the traditional medicine & herbal industry [3]. Plants have different components like alkaloids, glycoside, flavonoids, poly phenols etc, and each of them have different actions in preventing, curing or healing diseases.

Recently, WHO Estimated that 80% of world population is using traditional medicine. India is known as the “Emporium of Medicinal plants” due to availability of several thousands of medicinal plants in the different bioclimatic zones. India is one of the richest countries in the world in terms of biodiversity and documented systems of medicine like Ayurveda, Unani, Siddha & Homoeopathy (AYUSH System of Medicine) [4]. Pharmacopeias all over world contain at least 25% of herbal drugs or drugs derived from

plants. According to WHO, 21000 plant species are being used as the medicines. Usage of medicinal plants is increasing in both developed and developing nations, as they are non-narcotic, less adverse effects, low-cost and only source of treatment in poor countries. Best advantage of herbal medicine is age and sex of person are not considered in the treatment. Medicinal plants are widely used because they are easily available at low cost and less side effects.

Bioactive agents derived from natural sources have gained critical importance in modern medicine, lowering the risk of cardiac ailments by scavenging free radical formation [5]. Herbal medicines play an important role in the health of a large proportion of the world's population and have been recognised as a component of various tribe's cultural heritage. Polyphenols protect the heart by inhibiting the oxidation. The majority of pharmaceutically significant drugs are derived from plants. Plant derivatives as drugs play an important role in animal and human health-care systems all over the world [6,7]. Plants are an important source of traditional medicines that are used to treat a variety of ailments.

Chronic diseases like cancer, heart diseases can decrease effects on people by consumptions of fruits and vegetables which are in in bioactive compounds. Based on research so far, it is evident that phytochemicals having beneficial health effects. It is important to study the compounds from natural sources with effects against Myocardial Infarction and also prevents ischemic reperfusion injury, helps to reduce the damage caused by hypoxia and ischemia condition, reduces infarcts size [8]. Cardioprotective plants contain a variety of bioactive compounds, such as diosgenin, isoflavones, sulforaphane,

carotized, catechin, and quercetin, which have been shown to improve cardio protection and thus reduce the risk of cardiac abnormalities [9].

The coronary artery disease (CAD) is leading disease worldwide. The most common CAD is Myocardial infarction. Nearly 15-20% of mortality rate is reported each year. The incidence of MI in USA is about 525 billion people based on American heart association data and according to national survey of UK, people suffered from MI was reported as 915 billion [10] Heart attacks, also known as myocardial infarctions (MI), and their complications are the leading causes of death worldwide. Herbal antioxidants are increasingly being used as protective agents against a variety of cardiovascular abnormalities [11].

Approximately every 40 seconds, an American will have a myocardial infarction. The average age of MI occurrence is 66 years for men and 72 years for women [12]. In contrast to developed countries, developing countries like Sri Lanka, India, South Africa are also report high incidence of MI mostly in younger individuals aged between 45 to 65 years.

MI is an occlusion in the artery which blocks oxygenated blood supply to heart and depletes the oxygen to heart muscle which leads to oxidative stress and increased production of free radicals and causes cell death [13]. After onset of Myocardial ischemia, cell death is not immediate it can be identified imaging. Ischemic zone increases the demand of oxygen and nutrients which increases pressure and risk of cell death [14]. Earlier WHO defined MI from symptoms like ECG abnormalities, and enzymes. But Due to development of specific serological biomarkers like glutamine-oxaloacetic transaminase

(GOT), lactate dehydrogenase (LDH), creatine kinase (CK), MB fraction of CK, i.e., CKMB activity and CKMB etc, can be detected in case of myocardial necrosis and improved technology helps the accuracy of detecting MI with precise imaging [15].

Myocardial is a heart muscle and Infarction is a condition occurred due to increased oxygen demand or no oxygen supply to the heart. Myocardial infarction is divided into 2 types. Type 1 MI is caused because plaque rupture and Type 2 is caused due to reduction of oxygen supply. Two conditions are having different treatments and their pathogenesis is also poles apart. During ischemia or Hypoxia condition in the heart leads to a decrease in ATP production, which further Increases Cytosolic Calcium and Mitochondrial calcium [16].

The present article provides knowledge on mechanism of different plant components and how they act in repairing the MI tissue damage. World health Organization has started traditional medicine strategy which mainly encourages the use of traditional medicine as they play vital role in people's health and traditional medicines are low cost and effective when compared to other treatments [17].

2. Pathology of Myocardial Infarction:

During Myocardial Infarction, Hypoxia condition due to decreased oxygen in blood i.e., ischemia which results in reduction of ATP production, Ion pump function unbalance because of which increase in Na^+ and Ca^{2+} , which further increases cytosolic calcium and accumulation of calcium in Mitochondria [18]. Mitochondria is powerhouse of the cell and plays vital in pathogenesis of Myocardial infarction as they 40-50% of cardiomyocyte

cytoplasmic volume and important in cardiac balance since energy supply for cardiomyocytes is mostly derived from mitochondrial oxidative phosphorylation as they are special target for cellular damage. Besides this, activation of Phospholipids, protease, increases in loss of cell membrane phospholipids which further deteriorates the function of cell membrane and death of cytosol skeleton. Due to damage to cell membrane leakage of cell occurs and leads to cell death (fig. 1). Activation of free radicals leads to upregulation of a superoxide-generating enzyme as a result, production of Reactive Oxygen Species (ROS) starts [19-21].

Molecular and cellular mechanism during ischemic reperfusion injury is complex and damage occur in diverse pathways. Mitochondria are Major source for production ROS and major target for damage caused by ROS. Mitochondrial damage in cardiomyocytes leads to decreased energy production and loss of myocyte contractility and cell death [22]. During Ischemic reperfusion, ROS induces cell dysfunction and death by activating metalloproteinases and mitochondrial permeability transition pore (MPTP) which leads to cell swelling and death. ROS interacts with NO and increase in production of NO-producing enzyme which further damages cell by reaction of NO with superoxide and leads to cytotoxicity [23-26].

3. Ischemic reperfusion injury:

Ischemia causes an increase in intracellular sodium, hydrogen, and calcium ions, resulting in tissue acidosis. Reperfusion, in turn, causes rapid changes in ion flux, which leads to increased cytotoxicity [27,28]. Sodium-dependent pH regulatory mechanisms, such

as the $\text{Na}^+\text{-H}^+$ exchanger and the $\text{Na}^+\text{-HCO}_3^-$ transporter, are activated, resulting in intracellular sodium accumulation. Increases in sarcoplasmic reticular Ca^{2+} are caused by high sodium concentrations via the $\text{Na}^+\text{-Ca}^{2+}$ exchange [29] and Ca^{2+} overload is exacerbated by increased Ca^{2+} entry via sarcolemmal L-type Ca^{2+} channels [30,31] and decreased import of cytosolic Ca^{2+} into the sarcoplasmic reticulum by the SERCA $\text{Ca}^{2+}\text{ATPase}$ [32,33]. As a result, ATP depletion, ultrastructural mitochondrial damage, and myocardial stunning occur. Cardiac myocytes consume large quantities of energy. Normally, generation of ATP occurs by oxidative process to form water. Four electrons are required for oxygen to form water one at each step. Generation of free radicals causes cell damage and release of toxins into blood. Cell membrane damage also leads to leak of cytosol enzymes due breakdown of cell membrane phospholipids. Subsequent inflammation occurs due to activated neutrophils which utilises oxygen and releases lot of oxygen free radicals [34-36].

4. Cardioprotective Mechanism of Medicinal Plants:

Medicinal Plants have therapeutic effects which includes inhibiting, regulating the expression of various proteins such as contractile, and regulating the calcium levels and improvement in the functioning of mitochondria. Medicinal plants have evidence of decreasing the damage in heart muscles, and cardiomyocytes by regulating K_{ATP} channel (fig.2), and avoiding damage through oxidative stress and apoptosis. Inflammation inhibition, oxidative stress and apoptosis, endothelial nitric oxide synthase-nitric oxide (NOS-NO) signalling pathway activation, angiogenesis induction, and endothelial permeability suppression have all been shown to be beneficial effects of

medicinal plants/herbal products [37]. Types of chemical constituents present in plants are Saponin, flavonoids, glycoside, steroid, alkaloids, tannin, phenol, phlobatanin, terpenoids, and anthraquinone

5. Saponins and Cardio protection:

Saponins contains amphiphilic compounds with high molecular weight and having triterpenoid and steroid as lipophilic moiety and sugars act as hydrophilic moiety [38-40]. Saponins have a rigid skeleton with at least four hydrocarbon rings with attached one or two sugar groups. Traditionally, they are subdivided into triterpenoid and steroid glycosides. Steroidal saponins contains 27 carbon atoms forming the core structures like spirostan and furostan [41]. Saponins are 11 main classes: dammaranes, tirucallanes, lupanes, hopanes, oleananes, taraxasteranes, ursanes, cycloartanes, lanostanes, cucurbitanes, and steroids. The oleanane skeleton is the most common [42,43].

Pharmacological effects of saponins include stimulation of immune responses and also have ability to inhibit cell proliferation, to counteract angiogenesis and stimulates apoptosis (44,45). Besides that, Saponins also exhibit effects like antioxidant, anti-hypoxic, anoxia/reoxygenation, Ca²⁺ ion regulation or calcium antagonist, cardiocyte apoptosis, vasodilatory effect, and angiogenesis [46].

Ginseng has been most popular and common product, known for its cardio protection; which contains saponins as active constituent [47,48]. Overall, structural similarities with cardioactive phytosterols, as well as intriguing pharmacological effects such as hemolytic or permeabilization of cell membrane, antilipemic, serum cholesterol

lowering, and anticoagulant activity, prompted researchers to investigate the role of saponins in cardio protection [49,50].

Saponins cause erythrocytes to expand and rupture, releasing haemoglobin. Saponin's ability to cause erythrocyte death or haemolysis may limit the therapeutic use of the substances. Saponins have been proposed for the treatment of a variety of diseases, including diabetes, obesity and osteoporosis.

Increased Reactive oxygen species (ROS) causes intracellular oxidative, which is major reason of CVDs. ROS toxicity during reperfusion causes myocardial ischemia/reperfusion (I/R) injury through xanthine decomposition in mitochondria, an increase in cellular accumulation of lipid peroxides, a depletion of endogenous antioxidants, and Ca²⁺ ion overloading [51].

The cardioprotective activity of *Allium chinensis* is more when compared to a clinically approved calcium channel blocker nimodipine; which increases the malondialdehyde (MDA) formation and nitric oxide (NO) [52]. Other examples of Saponins viz., glycyrrhizic acid, asperosaponin VI, elatoside C, tribulosin, platycodin D, astragaloside IV, protodioscin, and trillin will enhance the super oxide dismutase (SOD), which work at cellular defense against ROS induced cardiac damage [53-56].

Saponins exhibit hypocholesterolaemic activity which involves forming insoluble complexes with cholesterol, affect micelle formation, interfere with bile acid metabolism, inhibit lipase activity and regulate cholesterol homeostasis [57,58]. This activity has been demonstrated in both animal and human trials. The best example of saponins currently being used as cardiac drug is Digoxin which is a procured from medicinal plant *Digitalis purpurea* and saikosaponin from *Bupleurumfalcatum* which is used as herbal supplement.

Saponins rich Food sources like *Saponaria*, soya, chick pea, *Yucca*, alfalfa, fenugreek, *Quillaja*, *Gypsophila*, and garlic resulted in reductions of cholesterol concentrations [59].

Total saponins and degradation production of total saponins of *Panacis majoris* Rhizoma protects the cardiomyocytes from injury induced by MIR, and also decreases the injury caused by oxidative stress, while maintaining the levels of LDH, CK-MB. But no significant change in infarct size is observed [60]. Other effects are shown in (Fig.3)[61].

6. Cardiac Glycosides and Cardio protection:

Cardiac glycosides contain steroidal framework in its core structure and their moiety is responsible for the activity. This steroid structure contains unsaturated lactone ring which characterizes the subgroup of glycosides [62]. Usually, cardiac glycosides are identified as secondary metabolites in plants and most of them belong to angiosperms. Recently cardiac glycosides are being identified from animal origin also [63]. Cardiac glycosides contain two classes of compounds cardenolids and bufadienolides, both differ by structure of Aglycone [64]. Plants produce both cardenolides and bufadienolides.

According to pathology of cell injury, Na-K-ATPase (sodium-potassium) pump is located in cell membrane that transports K ions in the cell and Na ions out the cell. This pump maintains balance between cytoplasm and extracellular fluid by using ATP [65]. Cardiac glycosides inhibit these pumps on Cardiac myocytes and prevents Na and calcium exist. Higher levels of cytoplasmic calcium improve the calcium intake into sarcoplasmic reticulum which enhances contractility of myocytes, together the ion build up in cell

increases cardiac output by accelerating force of contraction but intensity is unpredictable [66,67].

Convallaria Majalis is a Highly poisonous plant exhibits cardioprotective effects due to presence of 38 types of cardiac glycosides and active constituent of convallarin. This is more preferred than digitalis as it is safer. *Convallaria majalis* is used in the treatment of Arrhythmia and tachycardia [68]. Digoxin and digitoxin are two most widely used digitalis inotropes which belong to family of Scrophulariaceae [69]. The suffix “genin” refers to aglycone, without sugar, and digitoxin refers to digitoxigenin which contains aglycone and 3 sugar moieties. Aglycone portion in Cardiac glycosides is important for potential effect.

Normal, $\text{Na}^+\text{-K}^+$ pumps in the cardiac myocytes infuse potassium ions inside and diffuse sodium ions out of cell. Cardiac Glycosides inhibit the $\text{Na}^+\text{-K}^+$ pump, which stops sodium exit leads to increase intracellular sodium. In the same way, $\text{Na}^+\text{-Ca}^+$ pump allows calcium ions out of cell and sodium ions inside the cell and further raises intracellular sodium levels, which inhibit this pump; thus, calcium ions are not extruded and begins to build up inside the cell. Increased cytoplasmic calcium concentrations increase calcium uptake into the sarcoplasmic reticulum (SR) via the $\text{Ca}^{2+}\text{-ATPase}$ transporter [70]. For faster and stronger contraction, increased calcium in Sarcoplasmic reticulum, releases calcium on stimulation.

Due to Na-K-ATPase and cardiac glycoside binding is slow, it increases intracellular calcium gradually and action of is delayed [71]. Besides this, increased extracellular potassium also decreases binding of cardiac glycoside to Na-K-ATPase . Moreover, increases toxicity of cardiac glycosides to more presence in body and also

increases toxicity in presence of hypokalemia. When higher glycoside doses are administered, rhythm is lost and ventricular tachycardia develops, which is followed by fibrillation [72].

7. Conclusion:

The current review concluded that the therapeutic and prophylactic potential of plant phytoconstituents for the management of cardiovascular disorders has been investigated in several ways in, though the exact molecular mechanisms remain unknown. Phytoconstituents appear to have cardioprotective properties by Decreasing various factors, inhibiting key enzymes, and scavenging oxygen-free radicals. This review describes how phytochemicals like saponins, cardiac glycosides have diverse cardioprotective functions. Several plant-derived cardioprotective agents have been evaluated in clinical trials over the last few decades, with only a few being successful. Due to limited access to research articles and our search strategy, we were unable to include all of the studies describing the cardioprotective effects of medicinal plants or herbal agents in this review. However, the evidence presented in this review strongly suggests that medicinal plants/herbal products are a source of emerging medicines for the prevention and treatment of cardiovascular diseases. Developing more effective and safe agents from natural herbs is thus a promising approach to preventing and treating cardiovascular abnormalities. Documentation of clinical study criteria, on the other hand, is critical for standardizing the evaluation of medicinal plants/herbal agents.

Declaration of interest

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of this article.

NOTE:

The study highlights the efficacy of "Ayurved" which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

References:

1. World Health Organization. Traditional medicine: growing needs and potential. World Health Organization; 2002.
2. Zhang H. Bioactive Natural Products: Detection, Isolation, and Structural Determination. *Phytomedicine: International Journal of Phytotherapy & Phytopharmacology*. 2011 Jul 15;18(10):902-3.

3. <https://www.nmpb.nic.in/content/introduction>
4. https://www.nhp.gov.in/introduction-and-importance-of-medicinal-plants-and-herbs_mtl - Introduction and Importance of Medicinal Plants and Herbs
5. Wang CZ, Mehendale SR, Yuan CS. Commonly used antioxidant botanicals: active constituents and their potential role in cardiovascular illness. *The American journal of Chinese medicine*. 2007;35(04):543-58.
6. Zern TL, Fernandez ML. Cardioprotective effects of dietary polyphenols. *The Journal of nutrition*. 2005 Sep;135(10):2291-4.
7. Murad W, Azizullah A, Adnan M, Tariq A, Khan KU, Waheed S, Ahmad A. Ethnobotanical assessment of plant resources of Banda Daud Shah, district Karak, Pakistan. *Journal of ethnobiology and ethnomedicine*. 2013 Dec;9(1):1-0.
8. Hjartaker A, Knudsen MD, Tretli S, Weiderpass E. Consumption of berries, fruits and vegetables and mortality among 10,000 Norwegian men followed for four decades. *European journal of nutrition*. 2015 Jun 1;54(4):599-608.
9. Stoner GD. Foodstuffs for preventing cancer: the preclinical and clinical development of berries. *Cancer Prevention Research*. 2009 Mar 1;2(3):187-94.
10. Roger VL. Epidemiology of myocardial infarction. *Medical Clinics of North America*. 2007 Jul 1;91(4):537-52.
11. Biradar DP. Medicinal plants and phytomedicines. *Ann. Phytomed*. 2015 Jan 1;4(1):1-5.
12. Jayaraj JC, Davatyan K, Subramanian SS, Priya J. Epidemiology of myocardial infarction. In *Myocardial Infarction 2018* Nov 5. IntechOpen.
13. Benjamin EJ, Muntner P, Alonso A, Bittencourt MS, Callaway CW, Carson AP, Chamberlain AM, Chang AR, Cheng S, Das SR, Delling FN. Heart disease and stroke Statistics-2019 update a report from the American Heart Association. *Circulation*. 2019.
14. Pasotti M, Prati F, Arbustini E The pathology of myocardial infarction in the pre- and post-interventional era *Heart* 2006;92:1552-1556.

15. Thygesen K, Alpert JS, White HD, Joint ESC/ACCF/AHA/WHF Task Force for the Redefinition of Myocardial Infarction. Universal definition of myocardial infarction. *Journal of the American College of Cardiology*. 2007 Nov 27;50(22):2173-95.
16. Reddy K, Khaliq A, Henning RJ. Recent advances in the diagnosis and treatment of acute myocardial infarction. *World journal of cardiology*. 2015 May 26;7(5):243.
17. https://apps.who.int/iris/bitstream/handle/10665/92455/9789241506090_eng.pdf;jsessionid=3E68702BF27C01A0149E6F017F6316CE?sequence=1
18. Laguens RP, GÓMEZ-DUMM CL. Fine structure of myocardial mitochondria in rats after exercise for one-half to two hours. *Circulation research*. 1967 Sep;21(3):271-80.
19. Kane JJ, Murphy ML, Bisset JK, deSoyza N, Doherty JE, Straub KD. Mitochondrial function, oxygen extraction, epicardial ST segment changes and tritiated digoxin distribution after reperfusion of ischemic myocardium. *The American journal of cardiology*. 1975 Aug 1;36(2):218-24.
20. Capetanaki Y. Desmin cytoskeleton: a potential regulator of muscle mitochondrial behavior and function. *Trends in cardiovascular medicine*. 2002 Nov 1;12(8):339-48.
21. Andreyev AY, Kushnareva YE, Starkov AA. Mitochondrial metabolism of reactive oxygen species. *Biochemistry (Moscow)*. 2005 Feb;70(2):200-14.
22. Jennings RB, Ganote CE. Mitochondrial structure and function in acute myocardial ischemic injury. *Circulation research*. 1976 May 1;38(5 Suppl 1):I80-91.
23. Inoue M, Sato EF, Nishikawa M, Park AM, Kira Y, Imada I, Utsumi K. Mitochondrial generation of reactive oxygen species and its role in aerobic life. *Current medicinal chemistry*. 2003 Dec 1;10(23):2495-505.
24. Murphy MP. How mitochondria produce reactive oxygen species. *Biochemical journal*. 2009 Jan 1;417(1):1-3.
25. Starkov AA. The role of mitochondria in reactive oxygen species metabolism and signaling. *Annals of the New York Academy of Sciences*. 2008 Dec; 1147:37.

26. Tsutsui H, Kinugawa S, Matsushima S. Mitochondrial oxidative stress and dysfunction in myocardial remodelling. *Cardiovascular research*. 2009 Feb 15;81(3):449-56.
27. Bond JM, Herman B, Lemasters JJ. Protection by acidotic pH against anoxia/reoxygenation injury to rat neonatal cardiac myocytes. *Biochem Biophys Res Commun*. 1991;179:798–803.
28. Lemasters JJ, Bond JM, Chacon E, Harper IS, Kaplan SH, Ohata H, Trollinger DR, Herman B, Cascio WE. The pH paradox in ischemia-reperfusion injury to cardiac myocytes. *EXS*. 1996;76:99–114.
29. Tani M, Neely JR. Role of intracellular Na⁺ in Ca²⁺ overload and depressed recovery of ventricular function of reperfused ischemic rat hearts. Possible involvement of H⁺-Na⁺ and Na⁺-Ca²⁺ exchange. *Circ Res*. 1989;65:1045–1056.
30. Bush LR, Romson JL, Ash JL, Lucchesi BR. Effect of diltiazem on extent of ultimate myocardial injury resulting from temporary coronary artery occlusion in dogs. *J Cardiovasc Pharmacol*. 1982;4:285–296.
31. du Toit EF, Opie LH. Modulation of severity of reperfusion stunning in the isolated rat heart by agents altering calcium flux at onset of reperfusion. *Circ Res*. 1992;70:960–967.
32. Krause S, Hess ML. Characterization of cardiac sarcoplasmic reticulum dysfunction during short-term, normothermic, global ischemia. *Circ Res*. 1984;55:176–184.
33. Kaplan P, Hendrikx M, Mattheussen M, Mubagwa K, Flameng W. Effect of ischemia and reperfusion on sarcoplasmic reticulum calcium uptake. *Circ Res*. 1992;71:1123–1130.
34. Kusuoka H, Porterfield JK, Weisman HF, Weisfeldt ML, Marban E. Pathophysiology and pathogenesis of stunned myocardium. Depressed Ca²⁺ activation of contraction as a consequence of reperfusion-induced cellular calcium overload in ferret hearts. *J Clin Invest*. 1987;79:950–961.
35. Nayler WG. The role of calcium in the ischemic myocardium. *Am J Pathol*. 1981;102:262–270.

36. Zimmerman ANE, Daems W, Hulsmann WC, Snijder J, Wisse E, Durrer D. Morphological changes of heart muscle caused by successive perfusion with calcium-free and calcium-containing solutions (calcium paradox) *Cardiovasc Res.* 1967;1:201–209.
37. Liu C, Huang Y. Chinese herbal medicine on cardiovascular diseases and the mechanisms of action. *Frontiers in pharmacology.* 2016 Dec 1;7:469.
38. Sparg SG, Light ME, Van Staden J. Biological activities and distribution of plant saponins. *Journal of ethnopharmacology.* 2004 Oct 1;94(2-3):219-43.
39. Vincken JP, Heng L, de Groot A, Gruppen H. Saponins, classification and occurrence in the plant kingdom. *Phytochemistry.* 2007 Feb 1;68(3):275-97.
40. Killeen GF, Madigan CA, Connolly CR, Walsh GA, Clark C, Hynes MJ, Timmins BF, James P, Headon DR, Power RF. Antimicrobial saponins of *Yucca schidigera* and the implications of their in vitro properties for their in vitro impact. *Journal of Agricultural and Food Chemistry.* 1998; 46:3178- 3186. doi:10.1021/jf970928j
41. Goud MJP, Komraiah A, Rao KN, Ragan A, Raju VS, Charya MAS. Antibacterial activity of some folklore medicinal plants from South India. *African Journal of Traditional, Complementary and Alternative Medicines.* 2008;5(4):421- 426. doi:10.4314/ajtcam.v5i4.446
42. Vincken JP, Heng L, de Groot A, Gruppen H. Saponins, classification and occurrence in the plant kingdom. *Phytochemistry.* 2007;68(3):275- 297. doi: 10.1016/j.phytochem. 2006.10.008
43. Bissinger R, Modicano P, Alzoubi K, Honisch S, Faggio C, Abed M, Lang F. Effect of saponin on erythrocytes. *International Journal of Hematology.* 2014;100(1):51- 59. doi:10.1007/s12185- 014- 1605
44. Du J- R, Long F- Y, Chen C. Research progress on natural triterpenoid saponins in the chemoprevention and chemotherapy of cancer. In: Bathaie SZ, Tamanoi F editors.
45. *The Enzymes, Vol 36. Natural Products and Cancer Signaling: Isoprenoids, Polyphenols and Flavonoids.* 2014; Elsevier Inc. London, UK; pp. 9- 130. doi:10.1016/ B978- 0- 12- 802215- 3.00006- 9

46. Elekofehinti OO. Saponins: Anti- diabetic principles from medicinal plants—A review *Pathophysiology*. 2015;**22**:95- 103. doi:10.1016/j.pathophys.2015.02.001
47. Karakida F, Ikeya Y, Tsunakawa M, Yamaguchi T, Ikarashi Y, Takeda S, *et al*. Cerebral protective and cognition-improving effects of sinapic acid in rodents. *Biol Pharm Bull* 2007;**30**:514-9.
48. Shi Y, Han B, Yu X, Qu S, Sui D. Ginsenoside Rb3 ameliorates myocardial ischemia-reperfusion injury in rats. *Pharmaceutical biology*. 2011 Sep 1;**49**(9):900-6.
49. Scott GI, Colligan PB, Ren BH, Ren J. Ginsenosides Rb1 and Re decrease cardiac contraction in adult rat ventricular myocytes: role of nitric oxide. *British journal of pharmacology*. 2001 Nov;**134**(6):1159-65.
50. Dini I, Tenore GC, Dini A. Saponins in *Ipomoea batatas* tubers: Isolation, characterization, quantification and antioxidant properties. *Food Chemistry*. 2009 Mar 15;**113**(2):411-9. Marrelli M, Conforti F, Araniti F, Statti GA. Effects of saponins on lipid metabolism: A review of potential health benefits in the treatment of obesity. *Molecules*. 2016;**21**: 1404. doi:10.3390/molecules21101404
51. Misra MK, Sarwat M, Bhakuni P, Tuteja R, Tuteja N. Oxidative stress and ischemic myocardial syndromes. *Med Sci Monit*. 2009 Sep 30;**15**(10):209-19.
52. Ren G, Qiao HX, Yang J, Zhou CX. Protective effects of steroids from *Allium chinense* against H₂O₂- induced oxidative stress in rat cardiac H9C2 cells. *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives*. 2010 Mar;**24**(3):404-9.
53. Haleagrahara N, Varkkey J, Chakravarthi S. Cardioprotective effects of glycyrrhizic acid against isoproterenol-induced myocardial ischemia in rats. *International journal of molecular sciences*. 2011 Oct;**12**(10):7100-13.
54. Li C, Gao Y, Tian J, Xing Y, Zhu H, Shen J. Long-term oral Asperosaponin VI attenuates cardiac dysfunction, myocardial fibrosis in a rat model of chronic myocardial infarction. *Food and chemical toxicology*. 2012 May 1;**50**(5):1432-8.
55. Wang M, Meng XB, Yu YL, Sun GB, Xu XD, Zhang XP, Dong X, Ye JX, Xu HB, Sun YF, Sun XB. Elatoside C protects against hypoxia/reoxygenation-induced

- apoptosis in H9c2 cardiomyocytes through the reduction of endoplasmic reticulum stress partially depending on STAT3 activation. *Apoptosis*. 2014 Dec;19(12):1727-35.
56. Zhang S, Li H, Yang SJ. Tribulosin protects rat hearts from ischemia/reperfusion injury. *Acta Pharmacologica Sinica*. 2010 Jun;31(6):671-8.
57. Basu N, Rastogi RP. Triterpenoid saponins and saponins. *Phytochemistry*. 1967 Sep 1;6(9):1249-70
58. Zhao D. Challenges associated with elucidating the mechanisms of the hypocholesterolaemic activity of saponins. *Journal of Functional Foods*. 2016;23:52- 65. doi:10.1016/j.jff.2016.02.023
59. Güçlü- Ustündağ O, Mazza G. Saponins: properties, applications and processing. *Critical Reviews in Food Science and Nutrition*. 2007;47(3):231- 258. doi:10.1080/10408390600698197
60. Li M, Li X, Zhou L, Jin Y. Effects of total saponins from *Panacis majoris* Rhizoma and its degradation products on myocardial ischemia-reperfusion injury in rats. *Biomedicine & Pharmacotherapy*. 2020 Oct 1; 130:110538.
61. Nikitha G, SANDUR R. Cardioprotective potential of plants and plant-derived principles—a review. *Asian J. Pharmaceut. Clin. Res*. 2019; 12:46-56.
62. Steyn PS, van Heerden FR. Bufadienolides of plant and animal origin. *Natural product reports*. 1998;15(4):397-413.
63. Na SM. K-ATPase: Ubiquitous multifunctional transmembrane protein and its relevance to various pathophysiological conditions. *J Clin Med Res*. 2010;2(1):17.
64. Pettit GR, Green B, Das Gupta AK, Whitehouse PA, Yardley JP. Steroids and related natural products. I. Bufadienolides. 3. Synthetic route to isocardenolides. *The Journal of organic chemistry*. 1970 May;35(5):1381-4.
65. M.D. Forrest, the sodium-potassium pump is an information processing element in brain computation, *Front. Physiol.* 5 (472) (2014), doi: <http://dx.doi.org/10.3389/fphys.2014.00472>.
66. P. Bravený, Heart, calcium and time, *Exp. Clin. Cardiol.* 7 (2002) 3–6.

67. Tokhtaeva E, Sachs G, Vagin O. Assembly with the Na, K-ATPase α 1 subunit is required for export of β 1 and β 2 subunits from the endoplasmic reticulum. *Biochemistry*. 2009 Dec 8;48(48):11421-31.
68. Morris ET. *Fragrance: The story of perfume from Cleopatra to Chanel*. Scribner Book Company; 1984.
69. Melero CP, Medarde M, San Feliciano A. A short review on cardiotonic steroids and their aminoguanidine analogues. *Molecules*. 2000 Jan;5(1):51-81.
70. Fameli N, van Breemen C, Kuo KH. A quantitative model for linking Na⁺/Ca²⁺ exchanger to SERCA during refilling of the sarcoplasmic reticulum to sustain [Ca²⁺] oscillations in vascular smooth muscle. *Cell calcium*. 2007 Dec 1;42(6):565-75.
71. Radzyukevich TL, Lingrel JB, Heiny JA. The cardiac glycoside binding site on the Na, K-ATPase α 2 isoform plays a role in the dynamic regulation of active transport in skeletal muscle. *Proceedings of the National Academy of Sciences*. 2009 Feb 24;106(8):2565-70.
72. Kumar A, De T, Mishra A, Mishra AK. Oleandrin: A cardiac glycosides with potent cytotoxicity. *Pharmacognosy reviews*. 2013 Jul;7(14):131.

List of Figures:

Figure 1: Pathology of Myocardial Cell Damage during Ischemia and Hypoxia Condition

Fig 2: Cardioprotective mechanism of medicinal plants

Fig 3: Cardioprotective effect of Saponins

Figures:




Fig 1:




Fig 2:




Fig 3:

UNDER PEER REVIEW