

**QUALITATIVE AND QUANTITATIVE STUDY OF PHYTO-CONSTITUENTS AND
ANTIOXIDANT POTENTIAL OF RHIZOMES OF *KAEMPFERIA GALANGA*,
KAEMPFERIA PARVIFLORA AND *KAEMPFERIA PULCHRA***

ABSTRACT

Medicinal plants are the potent source of biologically active compounds and have always been of field of interest for the effective chemotherapeutic agents and offering a broad spectrum of activity with greater emphasis on preventive action. The objective of this study was to screen the phytochemicals, estimate the content of phenolic, flavonoids and alkaloids compounds and determines the antioxidant capacity of the rhizomes of *Kaempferia galanga* Linn, *Kaempferia parviflora* and *Kaempferia pulchra* (*K. galanga*, *K. parviflora* and *K. pulchra*, Zingiberaceae). Qualitative analysis of various phytochemical constituents and quantitative analysis of total phenol, flavonoids and alkaloids were determined by the well-known test protocol available in the literature. The ethanolic extract of rhizomes of *K. galanga*, *K. parviflora* and *K. pulchra* was studied for antioxidant activity on different *in vitro* models namely 1,1-diphenyl, 2-picryl hydrazyl (DPPH) assay and Nitric oxide (NO) radical scavenging method. Phytochemical analysis of ethanolic extract of all three plants revealed the presence of flavonoids, alkaloids, saponins, phenolics, carbohydrate, and tannin. The total phenolic, flavonoids and alkaloids content of ethanolic extract of *K. galanga*, *K. parviflora* and *K. pulchra* rhizomes were 0.813, 1.146; 1.047, 1.237, 0.285; 0.755, 0.822, 0.975/100mg respectively. All extracts showed dose dependent free radical scavenging property in the tested models, which was comparable to that of ascorbic acid. The present study concluded that the crude extract of *K. galanga*, *K. parviflora* and *K. pulchra* rhizomes is a rich source of secondary phytoconstituents which impart significant antioxidant potential. It is expected that the important phytochemical properties recognized by our study in the indigenous medicinal plants will be very useful in the curing of various diseases when taken along with our food.

Keywords: *Kaempferia galanga*, *Kaempferia parviflora*, *Kaempferia pulchra*, Zingiberaceae

INTRODUCTION

Herbal medicines have become more popular in the treatment of any diseases due to the popular belief that green medicine is safe, easily available and with fewer side effects. Many plants are cheaper and more accessible to most people especially in the developing countries than orthodox medicine, and there is a lower incidence of adverse effects after use. These reasons might account for their worldwide attention and use [1]. The medicinal properties of some plants have been documented by some researchers [2-4]. Medicinal plant constitutes the

main source of new pharmaceuticals and healthcare products [5]. Extraction and characterization of several phyto compounds of these green factories have given birth to some high activity profile drugs [6]. Indeed, the market and public demand has been so great that there is a great risk that many medicinal plants today face either extinction or loss of genetic diversity [7]. Knowledge of the chemical constituents of the plant is desirable because such information will be valuable for the synthesis of complex chemical substances. Reactive oxygen species (ROS) or oxygen free radicals can cause damage to cells and tissues during infections and various degenerative disorders such as cardiovascular diseases, aging and neurodegenerative diseases, like Alzheimer's disease, mutations and cancer [8,9]. The most widely used synthetic antioxidants, butylated hydroxyanisole (BHA) and butylated hydroxytoluene (BHT) have been restricted because of serious concerns about their carcinogenic potential [9, 10]. Natural antioxidants, especially phenolics and flavonoids, are safe; they protect the human body from free radicals and retard the progress of many chronic diseases as well as lipid oxidative rancidity in foods [11]. Numerous studies were carried out on plants with antioxidant properties [10-12]. However, there is still great interest in finding new antioxidants from natural sources. *K. galanga* is an important medicinal plant and has been traditionally used in tropics and subtropics of Asia including Bangladesh, India, China, Japan and Indo-China [13]. *Kaempferia* is a genus of about 70 species of rhizomatous, aromatic perennial herbs found in Africa and South East Asia [14]. In India, it is cultivated mainly in the states of Tamil Nadu, Kerala, Karnataka and West Bengal [15]. The rhizomes of this plant are widely used in East Asia for a wide range of traditional and modern medicinal applications. It promotes digestion and cures rheumatism, asthma, headaches, cough, toothaches, skin disease, piles, fever, epilepsy, splenic disorders and wounds. The rhizomes contain alkaloid, starch, gum, fatty matter with a fragrant essential oil [16]. Pharmacological properties such as anti-inflammatory and analgesic, antidiarrhoeal, nematocidal, mosquito repellent and larvicidal, vasorelaxant, sedative, antineoplastic, antimicrobial, anti-oxidant and cytotoxic activity has been reported [17]. *K. parviflora* or Krachaidam is originally found in the North and Northeast of Thailand. The rhizomes of *K. parviflora*, also known as black ginger are popular as health-promoting herbs and traditionally used as a folk medicine for managing a variety of diseases, including inflammation, ulcers, gout, colic disorder, abscesses, allergy, and osteoarthritis [18, 19]. A number of pharmacological researches on *K. parviflora* have claimed the valuable benefits for a variety of diseases. *K. parviflora* has recently been reported to possess antimycobacterial, antiplasmodial, anti-peptic ulcer and anti-viral protease effects as well as

modulators of multidrug resistance in cancer cells [20]. *K. pulchra* a perennial herb is cultivated in some tropical countries including Myanmar, Indonesia, Malaysia and Thailand. It is commonly known as Shan-pan-oot and has been extensively used for cough, blood stimulation, quenching heat, urinary tract infection and mellitus diabetes and as a carminative, deodorant, and diuretic. It has been reported to possess antiinflammatory and antitumor activities. The rhizomes have been used locally for self-medication by cancer and AIDS patients. Previous studies reported the presence of sandar-acopimaradiene diterpenoids and ethyl 4-methoxy-trans-cinnamate [21-24]. Despite the immense ethno-medicinal properties attributed to *K. galanga*, *K. parviflora* and *K. pulchra*, the reported phyto-pharmacological study on variety levels of these plants is relatively infrequent to the best of our knowledge. Therefore, the present study was aimed to evaluate and compare antioxidant activity of the ethanolic extracts of *K. galanga*, *K. parviflora* and *K. pulchra*, by using classical in-vitro assays for the purpose of validating its ethno medicinal use.

MATERIAL AND METHOD

Plant material

Rhizomes of *K. galanga*, *K. parviflora* and *K. pulchra* were collected from rural area of Bhopal in the month of Dec. 2019. Plant material (rhizomes part) selected for the study were washed thoroughly under running tap water and then were rinsed in distilled water; they were allowed to dry for some time at room temperature. Then the plant material was shade dried without any contamination for about 3 to 4 weeks. Dried plant material was grinded using electronic grinder. Powdered plant material was observed for their colour, odour, taste and texture. Dried plant material was packed in air tight container and stored for phytochemical and biological studies.

Chemical reagents

All the chemicals used in this study were obtained from Hi Media Laboratories Pvt. Ltd. (Mumbai, India), Sigma-Aldrich Chemical Co. (Milwaukee, WI, USA), SD Fine-Chem. Ltd. (Mumbai, India) and SRL Pvt. Ltd. (Mumbai, India). All the chemicals and solvent used in this study were of analytical grade.

Defatting of plant material

The shade dried plant material was subjected to extraction with petroleum ether using maceration method. The extraction was continued till the defatting of the material had taken place.

Successive extraction with different solvents

45.6 gm dried powdered of *K. galanga*, 52.4 gm dried powdered of *K. parviflora* and 40 gm dried powdered of *K. pulchra* have been extracted with chloroform, ethyl acetate, ethanol and aqueous using maceration process for 48 hrs, filtered and dried using vacuum evaporator at 40°C. The dried crude concentrated extract was weighed to calculate the extractive yield then transferred to glass vials (6 × 2 cm) and stored in a refrigerator (4°C), till used for analysis [25].

Phytochemical screening

Phytochemical screening to detect the presence of bioactive agents was performed by standard procedures [26, 27]. After the addition of specific reagents to the solution, the tests were detected by visual observation of color change or by precipitate formation.

Total phenol determination

The total phenolic content was determined using the method of Olufunmiso *et al* [28]. A volume of 2ml of each extracts or standard was mixed with 1 ml of Folin Ciocalteu reagent (previously diluted with distilled water 1:10 v/v) and 1 ml (7.5g/l) of sodium carbonate. The mixture was vortexed for 15s and allowed to stand for 10min for colour development. The absorbance was measured at 765 nm using a UV/visible spectrophotometer. The total phenolic content was calculated from the standard graph of gallic acid and the results were expressed as gallic acid equivalent (mg/100mg).

Total flavonoids determination

The total flavonoid content was determined using the method of Olufunmiso *et al* [28]. 1ml of 2% AlCl₃ solution was added to 3 ml of extract or standard and allowed to stand for 15 min at room temperature; the absorbance of the reaction mixture was measured at 420 nm using UV/visible spectrophotometer. The content of flavonoids was calculated using standard graph of quercetin and the results were expressed as quercetin equivalent (mg/100mg).

Total alkaloids determination

The plant extract (1mg) was dissolved in methanol, added 1ml of 2 N HCl and filtered [29]. This solution was transferred to a separating funnel, 5 ml of bromocresol green solution and 5 ml of phosphate buffer were added. The mixture was shaken with 1, 2, 3 and 4 ml chloroform by vigorous shaking and collected in a 10-ml volumetric flask and diluted to the volume with chloroform. A set of reference standard solutions of atropine (40, 60, 80, 100 and 120 µg/ml) were prepared in the same manner as described earlier. The absorbance for test and standard solutions were determined against the reagent blank at 470 nm with an UV/Visible spectrophotometer. The total alkaloid content was expressed as mg of AE/100mg of extract.

Antioxidant activity

DPPH radical scavenging assay

DPPH scavenging activity was measured by modified method of Olufunmiso *et al.*, 2011 [28]. DPPH scavenging activity was measured by the spectrophotometer. Stock solution (6 mg in 100ml methanol) was prepared such that 1.5 ml of it in 1.5 ml of methanol gave an initial absorbance. Decrease in the absorbance in presence of sample extract at different concentration (10-100 µg/ml) was noted after 15 minutes. 1.5 ml of DPPH solution was taken and volume made till 3 ml with methanol, absorbance was taken immediately at 517 nm for control reading. 1.5 ml of DPPH and 1.5 ml of the test sample of different concentration were put in a series of volumetric flasks and final volume was adjusted to 3 ml with methanol. Three test samples were taken and each processed similarly. Finally the mean was taken. Absorbance at zero time was taken for each concentration. Final decrease in absorbance was noted of DPPH with the sample at different concentration after 15 minutes at 517 nm. The percentage inhibition of free radical DPPH was calculated from the following equation: % inhibition = [(absorbance of control - absorbance of sample)/absorbance of control] × 100%. Though the activity is expressed as 50% inhibitory concentration (IC50), IC50 was calculated based on the percentage of DPPH radicals scavenged. The lower the IC50 value, the higher is the antioxidant activity.

Nitric oxide (NO^o) radical scavenging assay

The determination of NO^o radical scavenging ability of the extracts is based on the inhibition of NO^o radical generated from sodium nitroprusside in phosphate buffer saline solution by Griess reagent (1% sulfanilamide, 2% orthophosphoric acid and 0.1% naphthyl ethylenediamine dihydrochloride). Scavengers of nitric oxide act against oxygen, prompting to lessened production of nitrite ions which can be monitored at 546 nm [30]. Briefly, sodium nitroprusside (0.6 ml, 5 mM) solution was mixed with and without varying the concentration of the extracts or Ascorbic acid (2 ml, 10-200µg/ml) and incubated at 25 ± 2°C for 5 h. Incubated solution (2 ml) was mixed with equal volume of Griess reagent and absorbance of the purple colored azo dye chromophore was measured at λ_{max} 546 nm using UV-Vis spectrophotometer. The NO^o radical scavenging ability was calculated using following

$$\text{formula: Scavenging activity (\%)} = \frac{(\text{Abs control} - \text{Abs sample})}{\text{Abs control}} \times 100$$

RESULTS AND DISCUSSION

The crude extracts so obtained after each of the successive maceration extraction process were concentrated on water bath by evaporation the solvents completely to obtain the actual yield of extraction. The yield of extracts obtained from the rhizomes of *K. galanga*, *K.*

parviflora and *K. pulchra* using chloroform, ethyl acetate, ethanol and water as solvents are depicted in the Table 1. The results of qualitative phytochemical analysis of the crude powder rhizomes of *K. galanga*, *K. parviflora* and *K. pulchra* are shown in Table 2-4. Ethanolic extract of all three plants revealed the presence of flavonoids, alkaloids, saponins, phenolics, carbohydrate, and tannin. Total phenolic compounds (TPC) was expressed as mg/100mg of gallic acid equivalent of dry extract sample using the equation obtained from the calibration curve: $Y = 0.011X + 0.011$, $R^2 = 0.998$, where X is the gallic acid equivalent (GAE) and Y is the absorbance. Total flavonoids content was calculated as quercetin equivalent (mg/100mg) using the equation based on the calibration curve: $Y = 0.032X + 0.018$, $R^2 = 0.998$, where X is the quercetin equivalent (QE) and Y is the absorbance. Total alkaloid content was calculated as atropine equivalent mg/100mg using the equation based on the calibration curve: $Y = 0.007X + 0.024$, $R^2 = 0.995$, where X is the Atropine equivalent (AE) and Y is the absorbance. The total phenolic, flavonoids and alkaloids content of ethanolic extract of *K. galanga*, *K. parviflora* and *K. pulchra* rhizomes were 0.813, 1.146; 1.047, 1.237, 0.285; 0.755, 0.822, 0.975/100mg respectively Table 5-7. DPPH radical scavenging assay measured hydrogen donating nature of extracts [31]. Under DPPH radical scavenging activity the inhibitory concentration 50% (IC₅₀) value of *K. galanga*, *K. parviflora* and *K. pulchra* ethanolic rhizomes extract was found to be 58.79, 34.69 and 83.01 µg/ml as compared to that of ascorbic acid (17.68 µg/ml). A dose dependent activity with respect to concentration was observed Table 8. Extracts showed NO^o scavenging effects by competing with oxygen to react with NO directly hence inhibited the nitrite ion formation [32]. *K. galanga*, *K. parviflora* and *K. pulchra* ethanolic rhizomes extract showed nitric oxide (NO^o) radical scavenging activity with IC₅₀ value of 90.36, 65.65 and 106.50 µg/ml as compared to that of ascorbic acid (IC₅₀ 24.63 µg/ml) Table 9.

Table 1 Results of percentage yield of *K. galangal*, *K. parviflora* and *K. pulchra*

S. No.	Extracts	(%) yield <i>K. galangal</i>	(%) yield <i>K. parviflora</i>	(%) yield <i>K. pulchra</i>
1.	Chloroform	2.36	2.88	3.65
2.	Ethyl acetate	5.42	5.74	4.47
3.	Ethanol	7.69	9.68	7.16
4.	Aqueous	8.44	11.27	6.53

Table 2 Result of phytochemical screening of extracts of *K. galangal*

S. No.	Constituents	Chloroform extract	Ethyl acetate extract	Ethanol extract	Aqueous extract
1.	Alkaloids A) Hager's Test:	-Ve	-Ve	-Ve	-Ve
2.	Glycosides				

	A) Legal's Test:	-Ve	-Ve	-Ve	-Ve
3.	Flavonoids A) Lead acetate Test: B) Alkaline Reagent Test:	-Ve -Ve	-Ve -Ve	+Ve +Ve	+Ve -Ve
4.	Saponins A) Froth Test:	-Ve	+Ve	+Ve	+Ve
5.	Phenolics A) Ferric Chloride Test:	-Ve	-Ve	+Ve	+Ve
6.	Proteins A) Xanthoproteic Test:	-Ve	-Ve	-Ve	-Ve
7.	Carbohydrate A) Fehling's Test:	+Ve	-Ve	+Ve	+Ve
8.	Tannin A) Gelatin test:	+Ve	+Ve	+Ve	+Ve

Table 3 Result of phytochemical screening of extracts of *K. parviflora*

S. No.	Constituents	Chloroform extract	Ethyl acetate extract	Ethanol extract	Aqueous extract
1.	Alkaloids A) Hager's Test:	-Ve	-Ve	+Ve	-Ve
2.	Glycosides A) Legal's Test:	-Ve	-Ve	+Ve	+Ve
3.	Flavonoids A) Lead acetate Test: B) Alkaline Reagent Test:	-Ve -Ve	+Ve -Ve	-Ve +Ve	+Ve -Ve
4.	Saponins A) Froth Test:	-Ve	-Ve	+Ve	+Ve
5.	Phenolics A) Ferric Chloride Test:	-Ve	+Ve	+Ve	+Ve
6.	Proteins A) Xanthoproteic Test:	-Ve	-Ve	-Ve	-Ve
7.	Carbohydrate A) Fehling's Test:	+Ve	-Ve	+Ve	-Ve
8.	Tannin B) Gelatin test:	-Ve	-Ve	+Ve	+Ve

Table 4 Result of phytochemical screening of extracts of *K. pulchra*

S. No.	Constituents	Chloroform extract	Ethyl acetate extract	Ethanol extract	Aqueous extract
1.	Alkaloids A) Hager's Test:	-Ve	-Ve	+Ve	+Ve
2.	Glycosides A) Legal's Test:	-Ve	-Ve	-Ve	-Ve
3.	Flavonoids A) Lead acetate Test: B) Alkaline Reagent	+Ve -Ve	+Ve -Ve	+Ve +Ve	+Ve +Ve

	Test:				
4.	Saponins A) Froth Test:	-Ve	-Ve	+Ve	+Ve
5.	Phenolics A) Ferric Chloride Test:	-Ve	-Ve	+Ve	-Ve
6.	Proteins A) Xanthoproteic Test:	-Ve	-Ve	-Ve	-Ve
7.	Carbohydrate A) Fehling's Test:	-Ve	-Ve	-Ve	-Ve
8.	Tannin A) Gelatin test:	++Ve	-Ve	+Ve	-Ve

Table 5 Estimation of total phenolic, flavonoids and alkaloid content of *K. galangal*

S. No.	Extracts	Total phenolic content (mg/100mg of dried extract)	Total flavonoids content (mg/ 100 mg of dried extract)	Total alkaloid content (mg/ 100 mg of dried extract)
1	Chloroform	-	-	-
2	Ethyl acetate	-	-	-
3	Ethanol	0.813	1.146	-
4	Aqueous	0.572	0.490	-

Table 6 Estimation of total phenolic, flavonoids and alkaloid content of *K. parviflora*

S. No.	Extracts	Total phenolic content (mg/100mg of dried extract)	Total flavonoids content (mg/ 100 mg of dried extract)	Total alkaloid content (mg/ 100 mg of dried extract)
1	Chloroform	-	-	-
2	Ethyl acetate	0.559	0.981	-
3	Ethanol	1.047	1.237	0.285
4	Aqueous	0.823	0.563	-

Table 7 Estimation of total phenolic, flavonoids and alkaloid content of *K. pulchra*

S. No.	Extracts	Total phenolic content (mg/100mg of dried extract)	Total flavonoids content (mg/ 100 mg of dried extract)	Total alkaloid content (mg/ 100 mg of dried extract)
1	Chloroform	-	0.134	-
2	Ethyl acetate	-	0.465	-
3	Ethanol	0.755	0.822	0.975
4	Aqueous	-	0.357	0.681

Table 8 % Inhibition of ascorbic acid and ethanolic extract of *K. galangal*, *K. parviflora* and *K. pulchra* using DPPH method

S. No.	Concentration (µg/ml)	% Inhibition			
		Ascorbic acid	<i>K. galangal</i>	<i>K. parviflora</i>	<i>K. pulchra</i>
1	10	44.65	30.11	35.41	28.66
2	20	48.62	39.57	45.89	30.87
3	40	65.34	42.98	57.22	39.24
4	60	69.65	49.77	60.55	43.51
5	80	77.41	57.32	68.77	46.55

6	100	84.13	65.24	69.32	56.44
IC 50		17.68	58.79	34.69	83.01

Table 9 Inhibition of ascorbic acid and ethanolic extract of *K. galangal*, *K. parviflora* and *K. pulchra* using NO method

S. No.	Concentration (µg/ml)	% Inhibition			
		Ascorbic acid	<i>K. galangal</i>	<i>K. parviflora</i>	<i>K. pulchra</i>
1	20	47.70	20.47	31.22	15.64
2	40	52.92	27.65	39.78	21.75
3	60	67.43	34.27	46.49	29.54
4	80	68.89	48.54	57.84	37.49
5	100	74.42	53.32	63.24	49.36
IC 50		24.63	90.36	65.65	106.50

CONCLUSION

Due to the presence of a good number of flavonoid, alkaloid and phenolics content and appreciable quantities of secondary metabolites in rhizomes parts of the plant studied here, the plant can be seen as a potential source of useful drugs. The presence of phytoconstituents in a considerable amount might serve to recognize the potential pharmacological importance of this plant in disease control. The medicinal value of plants lies in some chemical substances that have a definite physiological action on human body. It also justifies the folklore medicinal uses and claims about the therapeutic values of this plant as curative agent. We therefore suggest further the isolation, purification and characterization of the bioactive compounds from rhizomes of *K. galanga*, *K. parviflora* and *K. pulchra* with a view to obtain useful chemotherapeutic agents.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

1. Sofowora E. Medicinal plants and medicine in Africa 2nd eds. In.: John Wiley and Sons.; hibueze U, Akubugwo E: Nutritive values and phytochemical contents of some

- leafy vegetables grown with different fertilizers. *Agricultural Biology Journal of North America* 2011; 2:1437-1444.
2. Ukpabi Chibueze, E. Akubugwo. Nutritive values and phytochemical contents of some leafy vegetables grown with different fertilizers. *Agriculture and Biology Journal of North America* 2011; 2(12):1437-1444.
 3. Idu M, Obaruyi G, Erhabor J. Ethnobotanical uses of plants among the binis in the treatment of ophthalmic and ENT (ear, nose and throat) ailments. *Ethnobotanical Leaflets* 2009; (4):9
 4. Banso A, Adeyemo S. Evaluation of antibacterial properties of tannins isolated from *Dichrostachys cinerea*. *Afr J Biotechnol* 2007; 6(15):1785-1787
 5. Ivanova D, Gerova D, Chervenkov T, Yankova T. Polyphenols and antioxidant capacity of Bulgarian medicinal plants. *J Ethnopharmacology* 2005; 96(1-2):145-150
 6. Mandal V, Mohan Y, Hemalatha S. Microwave assisted extraction-an innovative and promising extraction tool for medicinal plant research. *Pharmacogn Rev* 2007; 1(1):7-18
 7. Misra A. Studies on biochemical and physiological aspects in relation to phyto-medicinal qualities and efficacy of the active ingredients during the handling, cultivation and harvesting of the medicinal plants. *J Med Plants Res* 2009; 3(13):1140-1146.
 8. Lee SE, Hwang HJ, Ha JS, Jeog HS, Kim JH. Screening of medicinal plant extracts for antioxidant activity, *Life Sci.* 2003; 73: 167-179;
 9. Tadhani MB, Patel VH, Subhash R. *In vitro* antioxidant activities of *Stevia rebaudiana* leaves, *J. Food Comp. Anal.* 2007; 20: 323-329;
 10. Baydar NG, Ozkan G, Yasar S. Evaluation of the antiradical and antioxidant potential of grape extracts, *Food Cont.* 2007; 18: 1131-1136;
 11. Kumaran A, Karunakaran J. *In vitro* antioxidant activities of methanol extracts of five *Phyllanthus* species from India, *LWT – Food Sci. Tech.* 2007; 40: 344-352;
 12. Prakash D, Suri S, Upadhyay M, Singh BN. Total phenol, antioxidant and free radical scavenging activities of some medicinal plants, *Int. J. Food Sci. Nutr.* 2007; 58: 18-28;
 13. Aydin H, Hanefi O, Idris T, Oner CA. Antinociceptive activity of alpha-pinene and fenchone. *Pharmacologyonline*, 2008; 3: 363–369 (2008).

14. Granger RE, Campbell EL, Johnston GAR. (+)- and (-)- borneol: efficacious positive modulators of GABA action at human recombinant $\alpha 1\beta 2\gamma 2$ LGABAA receptors. *Biochemical Pharmacology*, 2005; 69(7), 1101–1111.
15. Preetha TS, Hemanthakumar AS, Krishnan PN. A comprehensive review of *Kaempferia galanga* L. (Zingiberaceae): A high sought medicinal plant in Tropical Asia. *Journal of Medicinal Plants Studies*, 2013; 4: 270–76.
16. Kim NJ, Byun SG, Cho JE, Cheng K, Ahn YJ. Larvicidal activity of *Kaempferia galanga* rhizome phenyl propanoids towards three mosquito species. *Pest Management science*, 2008; 64, 857–862.
17. Mohammad Shawkat Ali, Pritesh Ranjan Dash and Mahmuda Nasrin. Study of sedative activity of different extracts of *Kaempferia galanga* in Swiss albino mice *BMC Complementary and Alternative Medicine* 2015; 15:158.
18. Saokaew S, Wilairat P, Raktanyakan P. et al., Clinical Effects of *Krachaidum* (*Kaempferia parviflora*): A Systematic Review, *Evidence-Based Complementary and Alternative Medicine*, 2017; 22(3), 413-428.
19. Toda K, Hitoie S, Takeda S, Shimoda H. Black ginger extract increases physical fitness performance and muscular endurance by improving inflammation and energy metabolism, *Heliyon*, 2016; 2(5) Article ID e00115.
20. Supinya Tewtrakul, Sanan Subhadhirasakul, Sopa Kummee. Anti-allergic activity of compounds from *Kaempferia parviflora*. *Journal of Ethnopharmacology* 116 (2008) 191–193
21. The Traditional Medicine Formulations Used in Myanmar Traditional Medicine; Department of Traditional Medicine, Ministry of Health: Myanmar, 1990.
22. Tuchinda P, Udchachon J, Reutrakul V, Santisuk T, Skelton BW, White AH, Taylor WC. *Phytochemistry* 1994, 36, 731–734.
23. Prasad S, Yadav VR, Sundaram C, Reuter S, Hema PS, Nair MS, Chaturvedi MM, Aggarwal BBJ. *Biol. Chem.* 2010, 285, 26987–26997.
24. Prawat U, Tuntiwachwuttikul P, Taylor WC, Engelhardt LM, Skelton BW, White AH. *Phytochemistry* 1993, 32, 991–997.
25. Mukherjee PK. Quality control of herbal drugs. 2nd Ed. Business Horizons; 2007.
26. Khandelwal KR. Practical pharmacognosy technique and experiments. 23rd Ed. Nirali Prakashan; 2005.
27. Kokate CK. Practical pharmacognosy. 4th Ed. Vallabh Prakashan; 1994.

28. Olufunmiso OO, Afolayan AJ, Phenolic content and antioxidant property of the bark extract of *Ziziphus mucronata* willd. Subsp. *mucronata* willd, BMC Complement Alter Med. 2011; 11:130.
29. Fazel Shamsa, Hamidreza Monsef, Rouhollah Ghamooshi, Mohammadreza Verdian-rizi. Spectrophotometric determination of total alkaloids in some Iranian medicinal plants. Thai J Pharm Sci. 2008; 32: 17-20.
30. Shirwaikarkar A, Somashekar AP. Anti-inflammatory and free radical scavenging studies of *Aristyolochia bracteolate* Lam. India J Pharm Sci 2003;65:67-9.
31. Hudson BJ. Food antioxidants. In: Gordon MH, editor. The Mechanism of Antioxidant Action *in Vitro*. London: Elsevier Applied Science; 1990.
32. Sunil C, Ignacimuthu S. *In vitro* and *in vivo* antioxidant activity of *Symplocos cochinchinensis* S. Moore leaves containing phenolic compounds. Food Chem Toxicol 2011;49:1604-9.