

## PRODUCTION OF PROTEASE ENZYME AND ANTIMICROBIAL ACTIVITIES FROM MARINE ACTINOBACTERIUM OF STREPTOMYCES SP.

**Running title:** PROTEASE ENZYME AND ANTIMICROBIAL ACTIVITIES FROM MARINE STREPTOMYCES SP.

### ABSTRACT

#### Introduction:

Actinobacteria are Gram-positive bacteria that have traits of both bacteria and fungus. They are found in both terrestrial and aquatic habitats and aid in the breakdown of deceased species' organic matter. Marine actinobacteria are unusual in that they have antibacterial, anticancer, antiviral, insecticidal and enzyme inhibitory activities. The biggest genus of Actinobacteria is Streptomyces. They create more than two-thirds of all natural antibiotics that are therapeutically relevant (*e.g.* Neomycin, Cypemycin, Grisemycin, Bottromycins and Chloramphenicol). Alkaline proteases are generated by a variety of microorganisms, including yeast, bacteria, actinomycetes, fungus and plants. The process of destroying or suppressing disease-causing bacteria is referred to as antimicrobial activity. This is accomplished using a variety of antimicrobial drugs as well as antiviral properties.

**Aim:** To find out the antimicrobial activity of protease enzymes produced by the *Streptomyces* sp of marine actinobacterium.

**Materials and methods:** A sediment sample was collected and marine Actinobacteria were isolated in Kuster's Agar Medium (KUA) and were identified based on their chemotaxonomical characteristics.

**Results:** Screening for protease was carried out to assess enzyme production and enzyme assay was carried out to assess the enzymatic activity. The effect and pH on enzyme production was observed. Then the antimicrobial potential of protease enzymes was tested against few bacterial strains at optimum temperature and pH.

**Conclusion:** From the above study, we conclude that the protease enzyme obtained from streptomyces shows a potential antimicrobial (antibacterial) effect.

**Keywords:** Protease Enzyme, Effect of pH, Antibacterial potential, Novel drug

## **INTRODUCTION:**

Gram-positive filamentous bacteria with a fungal morphology are known as actinomycetes. They are members of the phylum Actinobacteria and have a complex life cycle (Ozcan et al. 2012). They are found in terrestrial ecosystems, especially in soil, where they play an important role in the recycling of industrial wastes and biomaterials by decomposing complex polymeric structures in dead plants, animals, and flora (S, Ayswarya, and S 2020). Actinobacteria, especially *Streptomyces* sp., are known for producing a wide range of biologically active compounds that are used as antibacterials, antifungals, antivirals, antithrombotics, immunomodulators, anti-tumor drugs, and enzyme inhibitors in a variety of fields, including medicine (S, Ayswarya, and S 2020; Sivasankar et al. 2020). Due to the emergence of multi-resistant microorganisms to almost all available antibiotics, many researchers are now concentrating their efforts on discovering novel antimicrobials from a variety of natural resources, including those developed by actinomycetes, especially those isolated from previously unknown or understudied environments. Many metabolic pathways, primarily organised by polyketide synthases (PKS) and non-ribosomal peptide synthetases, generate antimicrobials (NRPS) (Handayani et al. 2018). Actinobacteria have a high prevalence and presence of these biosynthetic genes (Handayani et al. 2018; Maiti and Mandal 2021) (Vignesh et al. 2019) (Pranati et al. 2019).

*Streptomyces* are bacteria that form mycelium and have a complex developmental life cycle that involves sporulation and programmed cell death (Undabarrena et al. 2017a). Their unparalleled versatility in secondary metabolism pathways has made them important suppliers of bioactive molecules, accounting for two-thirds of all antibiotics recognised (Undabarrena et al. 2017a; Park and Kwon 2018). Genome mining has become a powerful method for revealing *Streptomyces* species' biotechnological potential, allowing researchers to identify biosynthetic gene clusters (BGCs) and even predict the chemical core structure of molecules (Undabarrena et al. 2017a; Park and Kwon 2018; “Antimicrobial Metabolites from *Streptomyces* Sp. SN0280,”

n.d.). *Streptomyces* have linear chromosomes, unlike other bacteria, and their genomes are among the largest in the bacterial world (Undabarrena et al. 2017a; Park and Kwon 2018; “Antimicrobial Metabolites from *Streptomyces* Sp. SN0280,” n.d.; Enany 2018). Because of their exceptional position as antibiotic producers, soil microorganisms from the *Streptomyces* genus have sparked a lot of interest, but their marine equivalent has gotten less attention (Chen et al. 2020). Extreme abiotic selection pressures and enormous biological diversity characterise the marine environment. Despite all of the isolation studies on marine actinobacteria, little is known about the molecular mechanisms that enable bacteria to adapt to their surroundings (Chen et al. 2020; Park and Kwon 2018). It is thought that marine actinobacteria have adapted by developing specific biological traits, leading to the speculation that novel organisms from previously unknown environments may contain unique bioactive cofactors. Chile has a long, largely unexplored coastline in the South Pacific. In Valparaiso Central Bay and the Comau fjord in Northern Patagonia, bioprospecting of actinobacteria for the discovery of novel marine-derived natural products, specifically antibiotics, was carried out (Undabarrena et al. 2017b). Both sites yielded a plethora of novel actinobacteria organisms with antimicrobial properties. The genome of a selected antimicrobial-producer marine *Streptomyces* strain from the marine sediment sample was sequenced in this context (Sivaperumal, Kamala, and Rajaram 2015).

Proteases are present in prokaryotes, fungi, plants, and animals and are essential for their survival. Serine, cysteine, and metalloproteases are commonly found in many pathogenic parasites, where they play important roles in immune evasion, nutrient acquisition for growth and proliferation, dissemination facilitation, and tissue damage during infection (Karthik et al. 2014). Proteases, as a result, play a crucial role. Several clinical trials have shown their benefits in oncology, inflammatory disorders, blood rheology management, and immune regulation, so their use in medicine is gaining traction (D et al. 2019). Proteases are widely employed in leather, textiles, medicines, and detergents, accounting for more than 60% of the global enzyme market (Mourão and Schwartz 2013). Alkaline proteases are generated by a variety of microorganisms, including yeast, bacteria, actinomycetes, fungus, and plants (Karuppiah, Sun, and Li 2016). The process of destroying or suppressing disease-causing bacteria is referred to as antimicrobial activity. This is accomplished using a variety of antibacterial drugs (Kasanah and Triyanto 2019) (Mohapatra et al. 2020). Many plants are known for their antibacterial, antifungal, and antiviral properties (Aafreen et al. 2019) (Das et al. 2019) (Das et al. 2019; A et

al. 2019). Further, our team has extensive knowledge and research experience that has translated into high quality publications (Rajeshkumar et al. 2018; Nandhini, Rajeshkumar, and Mythili 2019; Vairavel, Devaraj, and Shanmugam 2020; M. Gomathi et al. 2020; Rajasekaran et al. 2020), (Santhoshkumar et al. 2019), (Raj R, D, and S 2020), (Saravanan et al. 2018), (Gheena and Ezhilarasan 2019), (Ezhilarasan, Sokal, and Najimi 2018), (Ezhilarasan 2018), ((Vairavel, Devaraj, and Shanmugam 2020; A. C. Gomathi et al. 2020; Dua et al. 2019), (Ramesh et al. 2018; Arumugam, George, and Jayaseelan 2021; Joseph and Prasanth 2021; Ezhilarasan, Apoorva, and Ashok Vardhan 2019; Duraisamy et al. 2019), (Gnanavel, Roopan, and Rajeshkumar 2019), (Markov et al. 2021). The aim of this study was to find out the antimicrobial activity of protease enzymes produced by the *Streptomyces* sp of marine actinobacterium.

## **MATERIALS AND METHODS :**

### Sample collection

The sediment sample was collected from the Parangipettai area, Tamilnadu. The collected sample was sun dried for 48 hrs and turned into fine powder by mortar and pestle.

### Isolation of actinobacteria

Isolation and enumeration of actinobacteria were carried out in Kuster's agar medium (KUA) supplemented with 0.5% (W/v) NaCl. To minimize the fungal and bacterial contamination, KUA medium was supplemented with cycloheximide (10 µg/ml) and nalidixic acid (10 µg/ml) respectively (Kathiresan et al., 2005). Collected sediment samples were serially diluted and inoculated on KUA medium and incubated at 36°C for 7 days. The colonies were counted and the population density has been expressed as colony forming units per gram (CFU/g) of sediments. Morphologically distinct colonies were selected and pure cultures were obtained.

### Identification of marine actinobacteria

**Aerial mass colour:** The colour of the mature sporulating aerial mycelium was recorded in naked eye. When the aerial mass colour fell between two colours series, both the colours were recorded. If the aerial mass colour of a strain to be studied showed intermediate tints, then also, both the colour series were noted. The media used were Yeast Extract-Malt Extract Agar and Inorganic-Salt Starch Agar.

**Melanoid pigments:** The grouping was made on the production of melanoid pigments (i.e. greenish brown, brownish black or distinct brown, pigment modified by other colours) on the medium. The strains were grouped as melanoid pigment produced (+) and not produced (-). In a few cases, the production of melanoid pigments was delayed or weak, and therefore, it was not distinguishable. This is indicated as variable (V). This test was carried out on the media ISP-1 and ISP-7 (Appendix I), as recommended by the International Streptomyces Project (Shirling and Gottlieb, 1966).

**Reverse side pigments:** Reverse side pigment production of the isolate was determined on ISP7 medium. The pigment production was noted as distinctive (+) and not distinctive or none (-). In

case, a colour with low chroma such as pale yellow, olive or yellowish brown occurred, it was included in the latter group (-).

**Soluble pigments:** Soluble pigment production of isolate was observed on ISP7 medium. The diffusible pigment production other than melanin was considered positive (+) and not produced (-). The colour was recorded (red, orange, green, yellow, blue and violet).

**Spore chain morphology:** Spore morphological characters of the strains were studied by inoculating a loopful of one week old cultures into solidified agar medium contained sterile glass slide. The cultures were incubated at 28±20 C and examined periodically for the formation of aerial mycelium, sporophore structure and spore morphology.

**Chemotaxonomical characteristics**

### Hydrolysis

Hydrolysis was done for releasing amino acids. Harvested cells of each strain weighing 20 mg (fresh) were placed in an ampo bottle and 1 ml of 6 N HCl was added and sealed with an alcohol blast burner. The samples were kept at 1210 C for 20 h in a sand bath. The bottles were cooled by keeping them at a room temperature of 28±20C. Hydrolysis was also done for releasing sugars. Harvested cells of each strain weighing 50 mg (fresh) were placed in an ampo bottle and 1 ml of 0.5N HCl was added and sealed with alcohol blast burner. The samples were kept at 1100 C for 2 h. The bottles were then cooled by keeping them at a room temperature of 28±20 C.

**Thin Layer Chromatography (TLC):** Spotting of the whole cell hydrolysates was made carefully on TLC plate using a microliter pipette. Spots were 5-10 mm in diameter. This was done by multiple applications on the same spot of very small portions of the sample, which were dried by a hand dryer.

**Amino acids:** Each sample (3 µl) was applied on the baselines of the TLC plate (20 cm x 20 cm). Adjacent to this, 1µl of DL-diaminopimelic acid (an authentic material mixture of DAP isomers) and 1µl of amino acetic acid (glycine) were spotted as standards. The TLC plate was developed with the solvent system containing methanol: pyridine: glacial acetic acid: H<sub>2</sub>O (5: 0.5: 0.125: 2.5 v/v). It took approximately more than 4 h for development. The spots were visualized by spraying with 0.4% ninhydrin solution in water-saturated n-butanol, followed by heating at 1000 C for 5 min. Spots of amino acids ran faster than DAP. The sample spots were immediately compared with the spots of the standards since spots gradually disappeared in a few hours.

**Whole-Cell sugars:** On a cellulose TLC plate (20 cm x 20 cm), 5µl of samples was spotted along with 3 µl of sugar solutions as standards on the same plates. Galactose, arabinose, xylose and madurose were the sugars, which were used as standards. The TLC plate was developed with the solvent mixture containing ethyl acetate: pyridine: acetic acid: distilled water (8: 5: 1: 1.5 v/v). The development time was more than 4 h. Spots were visualized by spraying with aniline phthalate reagent (3.25 g of phthalic acid dissolved in 2 ml of aniline and made upto 100 ml with

water saturated n-butanol). The sprayed plate was heated at 100 C for 4 min. Hexoses appeared as yellowish brown spots and pentoses, as maroon coloured spots.

Assimilation of carbon source: The ability of the actinobacterial strain in utilizing various carbon compounds as source of energy was studied, following the method recommended by International Streptomyces Project (Shirling and Gottlieb, 1966). Chemically pure carbon source certified to be free of admixture with other carbohydrates and contaminating materials were used for this purpose. Carbon sources for this test were Arabinose, Xylose, Inositol, Mannitol, Fructose, Rhamnose, Sucrose and Raffinose. These carbon sources were sterilized by ether sterilization without heating. The media and plates were prepared and inoculated according to the convention of ISP project (Shirling and Gottlieb, 1966). For each of the carbon sources, utilization is expressed as positive (+), negative (-), or doubtful ( $\pm$ ). In the 'doubtful' strains, only a trace of growth slightly greater than that of the control was noticed.

#### Screening of protease Production

Protease activity of the strains was screened qualitatively in skim milk agar. After inoculation, the plates were incubated at 37°C for 3 days. The diameters of clear zone around colonies on skim milk agar were measured.

#### Determination of Enzyme activity

The medium was inoculated with 1 ml of spore suspension of a 7 days old culture and incubated in a rotary shaker (150rpm) at ambient temperature for three days. The cell free supernatant was collected by centrifugation at 12,000rpm for 15 min. The supernatant was the enzyme source. The substrate 2% of casein was prepared with a 50mM phosphate buffer (pH 7). 1ml of crude enzyme was added with 1ml of casein (2%) solution which incubated for 60 min at (50°C) desired temperature. After incubation 2ml of Trichloroacetic acid (TCA) solution were added and boiled for 15 min in a boiling water bath. Before cooling 0.5ml of folin phenol reagent was added and the color was measured at 660 nm. One unit of protease activity was defined as the amount of enzyme that releases 1 mg of reducing sugar as tyrosine per ml per min under the assay conditions.

Bacterial Suspension: The fungal pathogens *Aspergillus flavus* and *Candida albicans* and bacterial pathogen *Pseudomonas aeruginosa* was collected from Department of Microbiology, Saveetha medical college and hospital, Tamilnadu. The bacterial pathogens were cultured in Muller –Hinton Broth for 24 hr at room temperature. From this bacterial suspension was prepared with saline and the optical density was measured at 600 nm. The concentration of microbial suspension was fixed as 10<sup>6</sup> CFU/ml. 1ml of suspension was spread over on Muller Hinton agar plate and incubated for 24hrs at ambient temperature.

Antibacterial activity: The antimicrobial activity of the enzyme protease was performed with disc diffusion method. Whatman filter paper discs (5mm) were impregnated with various concentrations (50, 100, 150, 200, 250 µg/ml) of enzyme sample with oral antibiotic tetracycline and DMSO as a negative control. The inoculated plates were incubated for 24hr at room temperature and the inhibition zones around the discs were measured. All the results were expressed from an average of three with standard deviation.

Minimum Inhibitory Concentration:

Minimal Inhibition Concentration of the enzyme protease was determined in 5 concentrations (50, 100, 150, 200, 250 µg/ml) with tetracycline (Standard) and DMSO (negative control) on Mueller Hinton broth). The inoculated bacteria in test tubes are incubated for 24hr in ambient temperature then the optical density was observed.

**RESULTS AND DISCUSSION:**

The rise of widespread antibiotic-resistant bacteria heightened the need to discover new antimicrobial agents. Actinomycetes, especially *Streptomyces* sp., have attracted a lot of attention because they produce a lot of useful bioactive metabolites. Isolating these species from less-explored environments may improve the chances of discovering new microbial species.

Fig-1,2 antibiotic-resistant bacteria



Table-1 :- Depicts the antimicrobial activity in different concentrations

## Antimicrobial activity

$\mu\text{g/ml}$	<i>S. mutans</i>	<i>Klebsiella pneumoniae</i>	<i>Pseudomonas aeruginosa</i>
0	0	0	0
50	4 +/- 2.2	5 +/- 1.3	6 +/- 2.2
100	13 +/- 3.4	11 +/- 2.6	13 +/- 2.6
150	17 +/- 2.5	17 +/- 2.1	21 +/- 2.4
200	21 +/- 2.1	22 +/- 2.4	26 +/- 2.1
250	26 +/- 2.7	27 +/- 2.9	35 +/- 2.7

Table -2 :- Depicts the minimum inhibitory concentration of protease enzyme against the bacteria.

MIC	0	10	20	30	40	50	MIC
<i>S. mutans</i>	+	+	+	+	-	-	30 $\mu\text{g/ml}$
Tetracycli	+	+	+	-	-	-	20 $\mu\text{g/ml}$
<i>Klebsiella pneumoni</i>	+	+	+	-	-	-	20 $\mu\text{g/ml}$
Tetracycli	+	+	+	-	-	-	20 $\mu\text{g/ml}$
<i>Pseudomonas</i>	+	+	+	+	-	-	30 $\mu\text{g/ml}$
Tetracycli	+	+	-	-	-	-	10 $\mu\text{g/ml}$

This study isolated marine actinobacteria from a sediment sample and identified streptomycetes genus from the isolate using specific characteristics of the bacteria. Then the production of enzyme was confirmed by screening for protease and enzyme assay was carried out. Enzyme assay revealed that the total activity of the enzyme was 96.24 IU/mg in a media of 1L. The effect of the temperature and pH on enzyme production was studied on Muller hinton broth. It was found that the optimum pH and temperature for maximum enzymatic activity was and respectively. It was also observed that as the temperature increased, the enzymatic activity increased. However at higher temperature, the rate decreased again which might probably be due to enzyme degeneration. Finally, antibacterial testing was done by incorporating the enzyme in different concentrations into the media plates containing each of the bacteria and the zone of inhibition was measured. It was observed that at the higher concentrations, *S. mutans*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa* exhibited higher susceptibility towards the antibiotic when compared to *Staphylococcus* (Ezhilarasan, Apoorva, and Ashok Vardhan 2019) (Danda, Krishna, et al. 2010) (Ramadurai et al. 2019) (Sathivel et al. 2008) (Panda et al. 2016) (P. Neelakantan et al. 2012) (Govindaraju, Neelakantan, and Gutmann 2017) (Sekhar, Narayanan, and Baig 2001) (DeSouza et al. 2014) (Nasim et al. 2010) (Danda, Muthusekhar, et al. 2010) ("Molecular Structure and Vibrational Spectra of 2,6-Bis(benzylidene)cyclohexanone: A Density Functional Theoretical Study" 2011) (Putchala et al. 2013) (Prasanna Neelakantan, Grotra, and Sharma 2013) (Suresh et al. 2014). However, the minimum inhibitory concentration of protease

enzyme that prevented the growth of the bacterial strains was very close to that of the standard tetracycline.

## CONCLUSION :

Actinomycetes, especially streptomycetes, are still a major source of bioactive compounds used to treat infections, cancer, and a variety of other ailments. From the above study, we conclude that the protease enzyme obtained from streptomyces shows a potential antimicrobial effect (antibacterial).

## REFERENCES:

- Aafreen, M. Maajida, M. Maajida Aafreen, Roy Anitha, Rachel Christina Preethi, S. Rajeshkumar, and T. Lakshmi. 2019. "Anti-Inflammatory Activity of Silver Nanoparticles Prepared from Ginger Oil—An Invitro Approach." *Indian Journal of Public Health Research & Development*.  
<https://doi.org/10.5958/0976-5506.2019.01552.3>.
- A, Mohamed Thamemul Ansari K., Mohamed Thamemul Ansari K, Anitha Roy, and S. Rajeshkumar. 2019. "Anti-Inflammatory Activity of Cinnamon Oil Mediated Silver Nanoparticles -An in Vitro Study." *International Journal of Research in Pharmaceutical Sciences*.  
<https://doi.org/10.26452/ijrps.v10i4.1579>.
- "Antimicrobial Metabolites from Streptomyces Sp. SN0280." n.d.  
<https://doi.org/10.1021/acs.jnatprod.6b01016.s001>.
- Arumugam, Paramasivam, Rinku George, and Vijayashree Priyadharsini Jayaseelan. 2021. "Aberrations of m6A Regulators Are Associated with Tumorigenesis and Metastasis in Head and Neck Squamous Cell Carcinoma." *Archives of Oral Biology* 122 (February): 105030.
- Chen, Chao, Shuai Chen, Wanxing Zhang, Fenghao Yuan, Jicheng Yu, and Qiu Liu. 2020. "Streptomyces Sp. S501, a Marine Petroleum-Degrading Actinobacterium Isolated from Sediment of Yalujiang Estuary, Northern China, and Its Genome Annotation." *Current Microbiology* 77 (11): 3643–50.
- Danda, A. K., Tatiparthi M. Krishna, V. Narayanan, and A. Siddareddi. 2010. "Influence of Primary and Secondary Closure of Surgical Wound after Impacted Mandibular Third Molar Removal on Postoperative Pain and Swelling--a Comparative and Split Mouth Study." *Journal of Oral and Maxillofacial Surgery: Official Journal of the American Association of Oral and Maxillofacial*

- Surgeons* 68 (2). <https://doi.org/10.1016/j.joms.2009.04.060>.
- Danda, A. K., M. R. Muthusekhar, V. Narayanan, M. F. Baig, and A. Siddareddi. 2010. "Open versus Closed Treatment of Unilateral Subcondylar and Condylar Neck Fractures: A Prospective, Randomized Clinical Study." *Journal of Oral and Maxillofacial Surgery: Official Journal of the American Association of Oral and Maxillofacial Surgeons* 68 (6). <https://doi.org/10.1016/j.joms.2009.09.042>.
- Das, Anubhav, Anitha Roy, S. Rajeshkumar, and T. Lakshmi. 2019. "Anti-Inflammatory Activity of Turmeric Oil Mediated Silver Nanoparticles." *Research Journal of Pharmacy and Technology*. <https://doi.org/10.5958/0974-360x.2019.00596.1>.
- D, Dhamodharan, Jemimah Naine S, Merlyn Keziah S, and Subathra Devi C. 2019. "Novel Fibrinolytic Protease Producing VITSD8 from Marine Sponges." *Marine Drugs* 17 (3). <https://doi.org/10.3390/md17030164>.
- DeSouza, S. I., M. R. Rashmi, A. P. Vasanthi, S. M. Joseph, and R. Rodrigues. 2014. "Mobile Phones: The next Step towards Healthcare Delivery in Rural India?" *PloS One* 9 (8). <https://doi.org/10.1371/journal.pone.0104895>.
- Dua, Kamal, Ridhima Wadhwa, Gautam Singhvi, Vamshikrishna Rapalli, Shakti Dhar Shukla, Madhur D. Shastri, Gaurav Gupta, et al. 2019. "The Potential of siRNA Based Drug Delivery in Respiratory Disorders: Recent Advances and Progress." *Drug Development Research* 80 (6): 714–30.
- Duraisamy, Revathi, Chitra Shankar Krishnan, Hariharan Ramasubramanian, Jayakrishnakumar Sampathkumar, Saravanakumar Mariappan, and Azhagarasan Navarasampatti Sivaprakasam. 2019. "Compatibility of Nonoriginal Abutments With Implants: Evaluation of Microgap at the Implant-Abutment Interface, With Original and Nonoriginal Abutments." *Implant Dentistry* 28 (3): 289–95.
- Enany, Shymaa. 2018. *Basic Biology and Applications of Actinobacteria*. BoD – Books on Demand.
- Ezhilarasan, Devaraj. 2018. "Oxidative Stress Is Bane in Chronic Liver Diseases: Clinical and Experimental Perspective." *Arab Journal of Gastroenterology: The Official Publication of the Pan-Arab Association of Gastroenterology* 19 (2): 56–64.
- Ezhilarasan, Devaraj, Velluru S. Apoorva, and Nandhigam Ashok Vardhan. 2019. "Syzygium Cumini Extract Induced Reactive Oxygen Species-Mediated Apoptosis in Human Oral Squamous Carcinoma Cells." *Journal of Oral Pathology & Medicine: Official Publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology* 48 (2): 115–21.
- Ezhilarasan, Devaraj, Etienne Sokal, and Mustapha Najimi. 2018. "Hepatic Fibrosis: It Is Time to Go with Hepatic Stellate Cell-Specific Therapeutic Targets." *Hepatobiliary & Pancreatic Diseases International: HBPDI* 17 (3): 192–97.
- Gheena, S., and D. Ezhilarasan. 2019. "Synergic Acid Triggers Reactive Oxygen Species-mediated Cytotoxicity in HepG2 Cells." *Human & Experimental Toxicology* 38 (6): 694–702.
- Gnanavel, V., Selvaraj Mohana Roopan, and S. Rajeshkumar. 2019. "Aquaculture: An Overview of Chemical Ecology of Seaweeds (food Species) in Natural Products." *Aquaculture* 507 (May): 1–6.
- Gomathi, A. C., S. R. Xavier Rajarathinam, A. Mohammed Sadiq, and S. Rajeshkumar. 2020. "Anticancer Activity of Silver Nanoparticles Synthesized Using Aqueous Fruit Shell Extract of Tamarindus Indica on MCF-7 Human Breast Cancer Cell Line." *Journal of Drug Delivery Science and Technology* 55 (February): 101376.
- Gomathi, M., A. Prakasam, P. V. Rajkumar, S. Rajeshkumar, R. Chandrasekaran, and P. M. Anbarasan. 2020. "Green Synthesis of Silver Nanoparticles Using Gymnema Sylvestre Leaf Extract and Evaluation of Its Antibacterial Activity." *South African Journal of Chemical Engineering*. <https://doi.org/10.1016/j.sajce.2019.11.005>.
- Govindaraju, L., P. Neelakantan, and J. L. Gutmann. 2017. "Effect of Root Canal Irrigating Solutions on the Compressive Strength of Tricalcium Silicate Cements." *Clinical Oral Investigations* 21 (2). <https://doi.org/10.1007/s00784-016-1922-0>.
- Handayani, Ira, Shanti Ratnakomala, Puspita Lisdiyanti, Fahrurrozi, Mohammad Alanjary, Wolfgang Wohlleben, and Yvonne Mast. 2018. "Complete Genome Sequence of Streptomyces Sp. Strain BSE7F, a Bali Mangrove Sediment Actinobacterium with Antimicrobial Activities." *Genome*

- Announcements*. <https://doi.org/10.1128/genomea.00618-18>.
- Joseph, Betsy, and C. S. Prasanth. 2021. "Is Photodynamic Therapy a Viable Antiviral Weapon against COVID-19 in Dentistry?" *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology*.
- Karthik, L., Gaurav Kumar, Tarun Keswani, Arindam Bhattacharyya, S. Sarath Chandar, and K. V. Bhaskara Rao. 2014. "Protease Inhibitors from Marine Actinobacteria as a Potential Source for Antimalarial Compound." *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0090972>.
- Karuppiah, V., W. Sun, and Z. Li. 2016. "Natural Products of Actinobacteria Derived from Marine Organisms." *Studies in Natural Products Chemistry*. <https://doi.org/10.1016/b978-0-444-63602-7.00013-8>.
- Kasanah, Noer, and Triyanto Triyanto. 2019. "Bioactivities of Halometabolites from Marine Actinobacteria." *Biomolecules*. <https://doi.org/10.3390/biom9060225>.
- Maiti, Pulak Kumar, and Sukhendu Mandal. 2021. "Streptomyces Cupreus Sp. Nov., an Antimicrobial Producing Actinobacterium Isolated from Himalayan Soil." *Archives of Microbiology*. <https://doi.org/10.1007/s00203-020-02160-y>.
- Markov, Alexander, Lakshmi Thangavelu, Surendar Aravindhan, Angelina Olegovna Zekiy, Mostafa Jarahian, Max Stanley Chartrand, Yashwant Pathak, Farogh Marofi, Somayeh Shamlou, and Ali Hassanzadeh. 2021. "Mesenchymal Stem/stromal Cells as a Valuable Source for the Treatment of Immune-Mediated Disorders." *Stem Cell Research & Therapy* 12 (1): 192.
- Mohapatra, Subhashree, L. Leelavathi, S. Rajeshkumar, Sri Sakthi D., and Jayashri P. 2020. "Assessment of Cytotoxicity, Anti-Inflammatory and Antioxidant Activity of Zinc Oxide Nanoparticles Synthesized Using Clove and Cinnamon Formulation - An In-Vitro Study." *Journal of Evolution of Medical and Dental Sciences*. <https://doi.org/10.14260/jemds/2020/405>.
- "Molecular Structure and Vibrational Spectra of 2,6-Bis(benzylidene)cyclohexanone: A Density Functional Theoretical Study." 2011. *Spectrochimica Acta. Part A, Molecular and Biomolecular Spectroscopy* 78 (1): 113–21.
- Mourão, Caroline, and Elisabeth Schwartz. 2013. "Protease Inhibitors from Marine Venomous Animals and Their Counterparts in Terrestrial Venomous Animals." *Marine Drugs*. <https://doi.org/10.3390/md11062069>.
- Nandhini, N. T., S. Rajeshkumar, and S. Mythili. 2019. "The Possible Mechanism of Eco-Friendly Synthesized Nanoparticles on Hazardous Dyes Degradation." *Biocatalysis and Agricultural Biotechnology* 19 (May): 101138.
- Nasim, I., P. Neelakantan, R. Sujeer, and C. V. Subbarao. 2010. "Color Stability of Microfilled, Microhybrid and Nanocomposite Resins--an in Vitro Study." *Journal of Dentistry* 38 Suppl 2. <https://doi.org/10.1016/j.jdent.2010.05.020>.
- Neelakantan, Prasanna, Deeksha Grotra, and Subash Sharma. 2013. "Retreatability of 2 Mineral Trioxide Aggregate-Based Root Canal Sealers: A Cone-Beam Computed Tomography Analysis." *Journal of Endodontia* 39 (7): 893–96.
- Neelakantan, P., A. A. Varughese, S. Sharma, C. V. Subbarao, M. Zehnder, and G. De-Deus. 2012. "Continuous Chelation Irrigation Improves the Adhesion of Epoxy Resin-Based Root Canal Sealer to Root Dentine." *International Endodontic Journal* 45 (12). <https://doi.org/10.1111/j.1365-2591.2012.02073.x>.
- Ozcan, Kadriye, Ataç Uzel, E. Esin Hames-Kocabas, and Erdal Bedir. 2012. "Isolation of Secondary Metabolites from a Marine Derived Streptomyces Sp. 9CM17 and Investigation of Their Antimicrobial Activities." *New Biotechnology*. <https://doi.org/10.1016/j.nbt.2012.08.634>.
- Panda, S., J. Doraiswamy, S. Malaiappan, S. S. Varghese, and M. Del Fabbro. 2016. "Additive Effect of Autologous Platelet Concentrates in Treatment of Intrabony Defects: A Systematic Review and Meta-Analysis." *Journal of Investigative and Clinical Dentistry* 7 (1). <https://doi.org/10.1111/jicd.12117>.
- Park, Jin-Soo, and Hak Kwon. 2018. "New Naphthoquinone Terpenoids from Marine Actinobacterium, Streptomyces Sp. CNQ-509." *Marine Drugs*. <https://doi.org/10.3390/md16030090>.
- Pranati, T., Roy Anitha, S. Rajeshkumar, and Thangavelu Lakshmi. 2019. "Preparation of Silver

- Nanoparticles Using Nutmeg Oleoresin and Its Antimicrobial Activity against Oral Pathogens.” *Research Journal of Pharmacy and Technology*. <https://doi.org/10.5958/0974-360x.2019.00471.2>.
- Putchala, Manisha Chandini, Pratibha Ramani, Herald J. Sherlin, Priya Premkumar, and Anuja Natesan. 2013. “Ascorbic Acid and Its pro-Oxidant Activity as a Therapy for Tumours of Oral Cavity – A Systematic Review.” *Archives of Oral Biology*. <https://doi.org/10.1016/j.archoralbio.2013.01.016>.
- Rajasekaran, S., D. Damodharan, K. Gopal, B. Rajesh Kumar, and Melvin Victor De Pours. 2020. “Collective Influence of 1-Decanol Addition, Injection Pressure and EGR on Diesel Engine Characteristics Fueled with diesel/LDPE Oil Blends.” *Fuel* 277 (October): 118166.
- Rajeshkumar, S., S. Venkat Kumar, Arunachalam Ramaiah, Happy Agarwal, T. Lakshmi, and Selvaraj Mohana Roopan. 2018. “Biosynthesis of Zinc Oxide Nanoparticles using *Mangifera Indica* Leaves and Evaluation of Their Antioxidant and Cytotoxic Properties in Lung Cancer (A549) Cells.” *Enzyme and Microbial Technology* 117 (October): 91–95.
- Raj R, Kathiswar, Ezhilarasan D, and Rajeshkumar S. 2020. “ $\beta$ -Sitosterol-Assisted Silver Nanoparticles Activates Nrf2 and Triggers Mitochondrial Apoptosis via Oxidative Stress in Human Hepatocellular Cancer Cell Line.” *Journal of Biomedical Materials Research. Part A* 108 (9): 1899–1908.
- Ramadurai, N., D. Gurunathan, A. V. Samuel, E. Subramanian, and S. J. L. Rodrigues. 2019. “Effectiveness of 2% Articaine as an Anesthetic Agent in Children: Randomized Controlled Trial.” *Clinical Oral Investigations* 23 (9). <https://doi.org/10.1007/s00784-018-2775-5>.
- Ramesh, Asha, Sheeja Varghese, Nadathur D. Jayakumar, and Sankari Malaiappan. 2018. “Comparative Estimation of Sulfiredoxin Levels between Chronic Periodontitis and Healthy Patients - A Case-Control Study.” *Journal of Periodontology* 89 (10): 1241–48.
- Santhoshkumar, J., B. Sowmya, S. Venkat Kumar, and S. Rajeshkumar. 2019. “Toxicology Evaluation and Antidermatophytic Activity of Silver Nanoparticles Synthesized Using Leaf Extract of *Passiflora Caerulea*.” *South African Journal of Chemical Engineering* 29 (July): 17–23.
- Saravanan, Muthupandian, Selvaraj Arokiyaraj, Thangavelu Lakshmi, and Arivalagan Pugazhendhi. 2018. “Synthesis of Silver Nanoparticles from *Phenerochaete Chryso sporium* (MTCC-787) and Their Antibacterial Activity against Human Pathogenic Bacteria.” *Microbial Pathogenesis* 117 (April): 68–72.
- Sathivel, A., H. R. Raghavendran, P. Srinivasan, and T. Devaki. 2008. “Anti-Peroxidative and Anti-Hyperlipidemic Nature of *Ulva Lactuca* Crude Polysaccharide on D-Galactosamine Induced Hepatitis in Rats.” *Food and Chemical Toxicology: An International Journal Published for the British Industrial Biological Research Association* 46 (10). <https://doi.org/10.1016/j.fct.2008.07.016>.
- S, Ayswarya, Ayswarya, and S. 2020. “Antimicrobial and Antitubercular Activity of Endophytic Actinobacterium and *Streptomyces* Sp. SACC 4 Isolated from the Mangrove Plant, *Rhizophora Apiculate*.” *Bioscience Biotechnology Research Communications*. <https://doi.org/10.21786/bbrc/13.3/9>.
- Sekhar, C. H., V. Narayanan, and M. F. Baig. 2001. “Role of Antimicrobials in Third Molar Surgery: Prospective, Double Blind, randomized, Placebo-Controlled Clinical Study.” *The British Journal of Oral & Maxillofacial Surgery* 39 (2). <https://doi.org/10.1054/bjom.2000.0557>.
- Sivaperumal, P., K. Kamala, and R. Rajaram. 2015. “Bioactive DOPA Melanin Isolated and Characterised from a Marine Actinobacterium *Streptomyces* Sp. MVCS6 from Versova Coast.” *Natural Product Research* 29 (22): 2117–21.
- Sivasankar, Palaniappan, Subramaniam Poongodi, Anderson O. Lobo, and Arivalagan Pugazhendhi. 2020. “Characterization of a Novel Polymeric Biofloculant from Marine Actinobacterium *Streptomyces* Sp. and Its Application in Recovery of Microalgae.” *International Biodeterioration & Biodegradation*. <https://doi.org/10.1016/j.ibiod.2020.104883>.
- Suresh, P., K. Marimuthu, S. Ranganathan, and T. Rajmohan. 2014. “Optimization of Machining Parameters in Turning of Al-SiC-Gr Hybrid Metal Matrix Composites Using Grey-Fuzzy Algorithm.” *Transactions of Nonferrous Metals Society of China*. [https://doi.org/10.1016/s1003-6326\(14\)63412-9](https://doi.org/10.1016/s1003-6326(14)63412-9).
- Undabarrena, Agustina, Juan Antonio Ugalde, Eduardo Castro-Nallar, Michael Seeger, and Beatriz

- Cámara. 2017a. "Genome Sequence of Streptomyces Sp. H-KF8, a Marine Actinobacterium Isolated from a Northern Chilean Patagonian Fjord." *Genome Announcements*.  
<https://doi.org/10.1128/genomea.01645-16>.
- . 2017b. "Genome Sequence of Sp. H-KF8, a Marine Actinobacterium Isolated from a Northern Chilean Patagonian Fjord." *Genome Announcements* 5 (6). <https://doi.org/10.1128/genomeA.01645-16>.
- Vairavel, Mathivadani, Ezhilarasan Devaraj, and Rajeshkumar Shanmugam. 2020. "An Eco-Friendly Synthesis of Enterococcus Sp.–mediated Gold Nanoparticle Induces Cytotoxicity in Human Colorectal Cancer Cells." *Environmental Science and Pollution Research* 27 (8): 8166–75.
- Vignesh, S., Roy Anitha, S. Rajesh Kumar, and T. Lakshmi. 2019. "Evaluation of the Antimicrobial Activity of Cumin Oil Mediated Silver Nanoparticles on Oral Microbes." *Research Journal of Pharmacy and Technology*. <https://doi.org/10.5958/0974-360x.2019.00634.6>.
- Hui Tian, Jamil Shafi, Mingshan Ji, Yuhui Bi, and Zhiguo Yu, 2017. "Antimicrobial Metabolites from Streptomyces Sp. SN0280." *Journal of Natural Products* 2017 80 (4), 1015-1019 DOI: 10.1021/acs.jnatprod.6b01016.