

Cytotoxicity Activity Of Green Tea and Mint Formulation

Running Title: Cytotoxicity activity of green tea and mint formulation

ABSTRACT:

Background:

Cytotoxicity refers to a substance or process which results in cell damage or cell death. The prefix 'cyto' refers to a cell and 'toxic' to poison. The term is often used to describe chemotherapy drugs that kill cancer cells, but it may also be used to describe toxins, such as venom. Within our own immune system, we have cells that are considered cytotoxic, such as the T cell that kills bacteria, viruses, and cancer cells.

Materials and Methods:

100ml of distilled water, 1 gram of green tea leaf and 1 gram of powdered mint is added and the mixture is heated for 15 to 20 minutes and then filtered with the help of filtered paper. The mixture was again heated and concentrated from 70ml to 20ml. 2 grams of iodine free salt was weighed and dissolved in 200ml of distilled water. 6 well ELISA plates were taken and 10-12ml of saline water was filled. To that 10 nauplii were added to each well(5 μ l, 10 μ l, 20 μ l, 40 μ l, 80 μ l). Then the nanoparticles were added according to the concentration level. The plates were incubated for 24 hours. After 24 hours, the elisa plates were observed and noted for the number of live nauplii's present.

Results and Discussion:

The cytotoxicity activity of green tea with mint formulation shows that all the introduced shrimps were alive in the control whereas it is shows that the lethality of 5 μ l is 60%, 10 μ l is 70%, 20 μ l is 70%, 40 μ l is 70% and 80 μ l is 100%.

Conclusion:

Based on the result of the study it is finally concluded that cytotoxic activity of green tea and mint formulation has better effect.

Keyword: cytotoxicity, mint, green tea, antioxidant, green synthesis.

Introduction:

Cytotoxicity (1) refers to a substance or process which results in cell damage or cell death. The prefix 'cyto' refers to a cell and 'toxic' to poison. The term is often used to describe chemotherapy drugs that kill cancer (2)(3)(2,4) cells, but it may also be used to describe toxins, such as venom. Within our own immune system, we have cells that are considered cytotoxic, such as the T cell that kills bacteria, viruses, and cancer cells. And it also kills any cell harboring such pathogens (5) by recognizing foreign peptides that are transported to the cell surface bound to MHC class I molecules. Oxygen radicals, organic hydroperoxides, and other types of free radicals are produced by certain chemical carcinogens and appear to play a role in the carcinogenic process(6)(7). We now control many tumors with antitumor agents. However, the need for new antitumor agents is pressing, especially against solid tumors, especially since antitumor agents need to be used for long durations. It seems that edible plants, food spices, and daily beverages that have been continuously consumed by many people for many years may be a superior source of antitumor agent(8).

Many aromatic plants used in medicine, food, and pharmaceutical industries belong to the lamiaceae family. In this family, mentha is a well known genus that includes 25-30 species that are generally grown in temperate areas around the world(9). Mentha also acts as a good expectorant. The most cited activities of the plants are its antiviral, antibacterial(10)(11), antifungal and cytotoxicity(12). The widespread use of *M.piperita* in traditional medicines has inspired us to explore its potential biological activities, knowing that there are few previous studies reporting the cytotoxicity. 120 edible plants for antitumor promoting activities against the non-12-O-tetradecanoylphorbol-13-acetate (TPA)-type promoter, okadaic acid (OA), which promotes tumor formation by inhibiting protein phosphatase-2A. Peppermint was one of only eight plants that showed strong activity (86–100%) in suppressing the effect of OA.

Menthol derived from *M. piperita* appears to affect cytosolic arylamine N- acetyltransferase (NAT) activity in the human liver tumor cell line J5 differentially dependent on dose (Lin et al., 2001); higher doses (32 and 3.2 mM) inhibited NAT, a more moderate dose (0.32 mM) had no effect, and lower doses (0.032 and 0.0032 mM) promoted NAT relative to controls(13). Herbal(14) teas are not true teas, true teas like green tea, black tea, oolong tea. Green tea has been considered a medicine and a healthful beverage since ancient times. Traditional Chinese medicine has recommended this plant for headaches, body aches and pains, digestion, depression, detoxification, as an energizer and, in general, to prolong life. Green tea leaves contain three main components which act upon human health: xanthic bases (caf- feine and theophylline), essential oils and especially, polyphe- nolic compounds. Caffeine acts mainly upon the central ner- vous system, stimulating wakefulness, facilitating ideas association and decreasing the sensation of fatigue(15).

Green tea polyphenols- namely catechin, epicatechin, epigallocatechin, and epigallocatechin-3-gallate are important constituents of green tea. They are known to possess chemopreventive and therapeutic properties against various diseases, including cancer(16)(16,17). Several studies using cancer cell lines(18) and animal models of cariogenesis have shown that green tea polyphenol anti cancer and apoptosis(19) inducing properties. An important, well noted attribute of the chemopreventive action of green tea polyphenols is their differential activity in selectively targeting cancer cells and sparing normal cells(20).

Our team has extensive knowledge and research experience that has translate into high quality publications(21–25),(26),(27),(28),(29),(30),(31),(23,32,33),(34–38) ,(39),(40)

The aim of the study is to evaluate the cytotoxic effect of green tea and mint formulation.

Materials and Methods:

Preparation of Herbal Formulation:

100ml of distilled water, 1 gram of green tea leaf and 1 gram of powdered mint is added and the mixture is heated for 15 to 20 minutes and then filtered with the help of filtered paper. The mixture was again heated and concentrated from 70ml to 20ml.





Figure 1: Shows the preparation of herbal formulation



Brine shrimp lethality assay:

Figure 2: Shows the Brine shrimp lethality assay.

Determination of cytotoxicity effect: 2 grams of iodine free salt was weighed and dissolved in 200ml of distilled water. 6 well ELISA plates were taken and 10-12ml of saline water was filled. To that 10 nauplii were added to each well(5 μ l, 10 μ l, 20 μ l, 40 μ l, 80 μ l). Then the nanoparticles were added according to the concentration level. The plates were incubated for 24 hours. After 24 hours, the elisa plates were observed and noted for the number of live nauplii's present.

Results and Discussion:




Figure 3: Shows the count of alive shrimp after 24 hours in Brine shrimp assay

The cytotoxicity activity of green tea with mint formulation shows that all the introduced shrimps were alive in the control whereas it shows that the lethality of 5 μl is 60%, 10 μl is 70%, 20 μl is 70%, 40 μl is 70% and 80 μl is 100% (Figure 3). The cytotoxicity of the volatile may not be potent enough to use them as leading compounds for drug design, but they are worthy of further investigation as natural products isolated from daily beverages. Importantly, these volatile compounds are identified in many edible plants, food spices and beverages are frequently used for fragrance and flavor. There is no doubt that these volatile compounds have long been consumed by many people(41). For example, the most potent cytotoxic volatile green tea flavour, nerolidol, is also a common component of many essential oils.

Cell death and the inhibition of cellular communication may occur in vivo following exposure to free radical- generating tumor promoters and may be mechanisms of tumor promotion by such compounds. Such promoters might include TPA which enhances chemiluminescence in epidermal cells and phagocytes(42)(42,43). On the other side, when assessing cytotoxicity, factors including size, shape, physicochemical surface property, concentration, exposure time, and cell type should be considered.

In addition to that, cytotoxic mechanism including disruption of the cell membrane, oxidative stress, destruction of the cytoskeleton and loss of mitochondrial function is important(44)(44,45) Similarly, in previous study, the protective effect of green tea extract and its

constituent polyphenols on the nephrotoxicity induced by the immunosuppressant FK506 in porcine renal proximal tubular cell line, LLC-PK1 cells, was evaluated. A significant increase in apoptotic cells but the addition of green tea extract, and particularly its major polyphenolic component epigallocatechin- gallate, suppress the cell death(46).

Conclusion:

The allopath based anticancer drugs are expensive and also show adverse effects via alteration in molecular pathways(37) (47) (48) (49) (50) (51) (52) (53) (54) (55) (56) (57) (58) (59) (60). Green tea has been widely used in traditional medicine since ancient times due to being inexpensive, efficacy, and fewer side effect properties. Based on the result of the study, it is finally concluded that cytotoxic activity of green tea and mint formulation has better lethality effects in 80µl is 100%.

Reference:

1. Shree MK, Kavya Shree M, Arivarasu L, Rajeshkumar S. Cytotoxicity and Antimicrobial Activity of Chromium Picolinate Mediated Zinc Oxide Nanoparticle [Internet]. Journal of Pharmaceutical Research International. 2020. p. 28–32. Available from: <http://dx.doi.org/10.9734/jpri/2020/v32i2030726>

2. Ashwini S, Ezhilarasan D, Anitha R. Cytotoxic Effect of *Caralluma fimbriata* Against Human Colon Cancer Cells [Internet]. Vol. 9, Pharmacognosy Journal. 2017. p. 204–7. Available from: <http://dx.doi.org/10.5530/pj.2017.2.34>
3. Roy A, Department of Pharmacology, Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Sciences, Chennai, Nadu- T, et al. Molecular docking analysis of compounds from *Andrographis paniculata* with EGFR [Internet]. Vol. 17, Bioinformation. 2021. p. 23–8. Available from: <http://dx.doi.org/10.6026/97320630017023>
4. Oncogenes Associated with Oral Cancer - An Update [Internet]. Indian Journal of Forensic Medicine & Toxicology. 2020. Available from: <http://dx.doi.org/10.37506/ijfmt.v14i4.12404>
5. Paul RP, Roy A, Maajida AM, Shanmugam R. Antibacterial Activity of White Pepper Oleoresin Mediated Silver Nanoparticles against Oral Pathogens [Internet]. Vol. 9, Journal of Evolution of Medical and Dental Sciences. 2020. p. 2352–5. Available from: <http://dx.doi.org/10.14260/jemds/2020/510>
6. Cerutti P. Prooxidant states and tumor promotion [Internet]. Vol. 227, Science. 1985. p. 375–81. Available from: <http://dx.doi.org/10.1126/science.2981433>
7. Kensler TW, Trush MA. Role of oxygen radicals in tumor promotion [Internet]. Vol. 6, Environmental Mutagenesis. 1984. p. 593–616. Available from: <http://dx.doi.org/10.1002/em.2860060412>
8. Kubo I, Ochi M, Vieira PC, Komatsu S. Antitumor agents from the cashew (*Anacardium occidentale*) apple juice [Internet]. Vol. 41, Journal of Agricultural and Food Chemistry. 1993. p. 1012–5. Available from: <http://dx.doi.org/10.1021/jf00030a035>
9. Pickersgill B. Sesame: The Genus *Sesamum*. Medicinal and Aromatic Plants – Industrial Profiles. Edited by D. Bedigian. Boca Raton, FL, USA: CRC Press (2011), pp. xxiii 532, £82.00. ISBN 978-0-8493-3538-9 [Internet]. Vol. 47, Experimental Agriculture. 2011. p. 733–4. Available from: <http://dx.doi.org/10.1017/s0014479711000627>
10. Ashwini K, Rajeshkumar S, Roy A, Lakshmi T. *Symplocos racemosa* bark assisted copper nanoparticles and its antibacterial activity against *Staphylococcus aureus* and *Lactobacilli* species [Internet]. Vol. 14, RESEARCH JOURNAL OF PHARMACY AND TECHNOLOGY. 2021. p. 300–2. Available from: <http://dx.doi.org/10.5958/0974-360x.2021.00054.8>
11. Sunar S, Rajeshkumar S, Roy A, Lakshmi T. Preparation of herbal formulation and its application on nanoparticles synthesis and antibacterial activity [Internet]. Vol. 10, International Journal of Research in Pharmaceutical Sciences. 2019. Available from: <http://dx.doi.org/10.26452/ijrps.v10i3.1447>
12. Sakkas H, Papadopoulou C. Antimicrobial Activity of Basil, Oregano, and Thyme Essential Oils [Internet]. Vol. 27, Journal of Microbiology and Biotechnology. 2017. p. 429–38. Available from: <http://dx.doi.org/10.4014/jmb.1608.08024>

13. Lazutka JR, Mierauskienė J, Slapšytė G, Dedonytė V. Genotoxicity of dill (*Anethum graveolens* L.), peppermint (*Mentha piperita* L.) and pine (*Pinus sylvestris* L.) essential oils in human lymphocytes and *Drosophila melanogaster* [Internet]. Vol. 39, Food and Chemical Toxicology. 2001. p. 485–92. Available from: [http://dx.doi.org/10.1016/s0278-6915\(00\)00157-5](http://dx.doi.org/10.1016/s0278-6915(00)00157-5)
14. Herbal Sources Used by The Public Against Infections [Internet]. Vol. 12, International Journal of Pharmaceutical Research. 2020. Available from: <http://dx.doi.org/10.31838/ijpr/2020.sp1.015>
15. Varnam A, Sutherland JM. Beverages: Technology, Chemistry and Microbiology. Springer Science & Business Media; 1994. 464 p.
16. Flores-González M, Talavera-Rojas M, Soriano-Vargas E, Rodríguez-González V. Practical mediated-assembly synthesis of silver nanowires using commercial *Camellia sinensis* extracts and their antibacterial properties [Internet]. Vol. 42, New Journal of Chemistry. 2018. p. 2133–9. Available from: <http://dx.doi.org/10.1039/c7nj03812g>
17. Moulton MC, Braydich-Stolle LK, Nadagouda MN, Kunzelman S, Hussain SM, Varma RS. Synthesis, characterization and biocompatibility of “green” synthesized silver nanoparticles using tea polyphenols [Internet]. Vol. 2, Nanoscale. 2010. p. 763. Available from: <http://dx.doi.org/10.1039/c0nr00046a>
18. Suhasini SJ, Jennifer Suhasini S, Roy A, Sosa G, Lakshmi T. The Cytotoxic effect of *Caralluma fimbriata* on KB cell lines [Internet]. Vol. 12, Research Journal of Pharmacy and Technology. 2019. p. 4995. Available from: <http://dx.doi.org/10.5958/0974-360x.2019.00865.5>
19. Roy A, Rasheed A, Sreeba AV, Rajagopal P. Molecular docking analysis of capsaicin with apoptotic proteins. *Bioinformation*. 2020 Jul 31;16(7):555–60.
20. Matsuzaki T, Hara Y. Antioxidative activity of tea leaf catechins [Internet]. Vol. 59, Journal of the agricultural chemical society of Japan. 1985. p. 129–34. Available from: <http://dx.doi.org/10.1271/nogeikagaku1924.59.129>
21. Rajeshkumar S, Kumar SV, Ramaiah A, Agarwal H, Lakshmi T, Roopan SM. Biosynthesis of zinc oxide nanoparticles using *Mangifera indica* leaves and evaluation of their antioxidant and cytotoxic properties in lung cancer (A549) cells. *Enzyme Microb Technol*. 2018 Oct;117:91–5.
22. Nandhini NT, Rajeshkumar S, Mythili S. The possible mechanism of eco-friendly synthesized nanoparticles on hazardous dyes degradation. *Biocatal Agric Biotechnol*. 2019 May 1;19:101138.
23. Vairavel M, Devaraj E, Shanmugam R. An eco-friendly synthesis of *Enterococcus* sp.–mediated gold nanoparticle induces cytotoxicity in human colorectal cancer cells. *Environ Sci Pollut Res*. 2020 Mar 1;27(8):8166–75.

24. Gomathi M, Prakasam A, Rajkumar PV, Rajeshkumar S, Chandrasekaran R, Anbarasan PM. Green synthesis of silver nanoparticles using *Gymnema sylvestre* leaf extract and evaluation of its antibacterial activity [Internet]. Vol. 32, South African Journal of Chemical Engineering. 2020. p. 1–4. Available from: <http://dx.doi.org/10.1016/j.sajce.2019.11.005>
25. Rajasekaran S, Damodharan D, Gopal K, Rajesh Kumar B, De Poures MV. Collective influence of 1-decanol addition, injection pressure and EGR on diesel engine characteristics fueled with diesel/LDPE oil blends. *Fuel*. 2020 Oct 1;277:118166.
26. Santhoshkumar J, Sowmya B, Venkat Kumar S, Rajeshkumar S. Toxicology evaluation and antidermatophytic activity of silver nanoparticles synthesized using leaf extract of *Passiflora caerulea*. *S Afr J Chem Eng*. 2019 Jul;29:17–23.
27. Raj R K, D E, S R. β -Sitosterol-assisted silver nanoparticles activates Nrf2 and triggers mitochondrial apoptosis via oxidative stress in human hepatocellular cancer cell line. *J Biomed Mater Res A*. 2020 Sep;108(9):1899–908.
28. Saravanan M, Arokiyaraj S, Lakshmi T, Pugazhendhi A. Synthesis of silver nanoparticles from *Phenerochaete chrysosporium* (MTCC-787) and their antibacterial activity against human pathogenic bacteria. *Microb Pathog*. 2018 Apr;117:68–72.
29. Gheena S, Ezhilarasan D. Syringic acid triggers reactive oxygen species–mediated cytotoxicity in HepG2 cells. *Hum Exp Toxicol*. 2019 Jun 1;38(6):694–702.
30. Ezhilarasan D, Sokal E, Najimi M. Hepatic fibrosis: It is time to go with hepatic stellate cell-specific therapeutic targets. *Hepatobiliary Pancreat Dis Int*. 2018 Jun;17(3):192–7.
31. Ezhilarasan D. Oxidative stress is bane in chronic liver diseases: Clinical and experimental perspective. *Arab J Gastroenterol*. 2018 Jun;19(2):56–64.
32. Gomathi AC, Xavier Rajarathinam SR, Mohammed Sadiq A, Rajeshkumar S. Anticancer activity of silver nanoparticles synthesized using aqueous fruit shell extract of *Tamarindus indica* on MCF-7 human breast cancer cell line. *J Drug Deliv Sci Technol*. 2020 Feb 1;55:101376.
33. Dua K, Wadhwa R, Singhvi G, Rapalli V, Shukla SD, Shastri MD, et al. The potential of siRNA based drug delivery in respiratory disorders: Recent advances and progress. *Drug Dev Res*. 2019 Sep;80(6):714–30.
34. Ramesh A, Varghese S, Jayakumar ND, Malaiappan S. Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study. *J Periodontol*. 2018 Oct;89(10):1241–8.
35. Arumugam P, George R, Jayaseelan VP. Aberrations of m6A regulators are associated with tumorigenesis and metastasis in head and neck squamous cell carcinoma. *Arch Oral Biol*. 2021 Feb;122:105030.
36. Joseph B, Prasanth CS. Is photodynamic therapy a viable antiviral weapon against COVID-

19 in dentistry? *Oral Surg Oral Med Oral Pathol Oral Radiol*. 2021 Jul;132(1):118–9.

37. Ezhilarasan D, Apoorva VS, Ashok Vardhan N. Syzygium cumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells. *J Oral Pathol Med*. 2019 Feb;48(2):115–21.
38. Duraisamy R, Krishnan CS, Ramasubramanian H, Sampathkumar J, Mariappan S, Navarasampatti Sivaprakasam A. Compatibility of Nonoriginal Abutments With Implants: Evaluation of Microgap at the Implant-Abutment Interface, With Original and Nonoriginal Abutments. *Implant Dent*. 2019 Jun;28(3):289–95.
39. Gnanavel V, Roopan SM, Rajeshkumar S. Aquaculture: An overview of chemical ecology of seaweeds (food species) in natural products. *Aquaculture*. 2019 May 30;507:1–6.
40. Markov A, Thangavelu L, Aravindhan S, Zekiy AO, Jarahian M, Chartrand MS, et al. Mesenchymal stem/stromal cells as a valuable source for the treatment of immune-mediated disorders. *Stem Cell Res Ther*. 2021 Mar 18;12(1):192.
41. Burdon RH. Carcinogenesis and Free Radicals [Internet]. *Free Radicals: from Basic Science to Medicine*. 1993. p. 187–98. Available from: http://dx.doi.org/10.1007/978-3-0348-9116-5_16
42. Fischer SM, Adams LM. Inhibitors of Arachidonic Acid Metabolism Inhibit Tumor-Promoter-Stimulated Chemiluminescence in Murine Keratinocytes [Internet]. *Prostaglandins, Leukotrienes, and Lipoxins*. 1985. p. 609–18. Available from: http://dx.doi.org/10.1007/978-1-4684-4946-4_58
43. Rao SM, Reddy JK. Peroxisome proliferation and hepatocarcinogenesis [Internet]. Vol. 8, *Carcinogenesis*. 1987. p. 631–6. Available from: <http://dx.doi.org/10.1093/carcin/8.5.631>
44. Chuang S-M, Lee Y-H, Liang R-Y, Roam G-D, Zeng Z-M, Tu H-F, et al. Extensive evaluations of the cytotoxic effects of gold nanoparticles [Internet]. Vol. 1830, *Biochimica et Biophysica Acta (BBA) - General Subjects*. 2013. p. 4960–73. Available from: <http://dx.doi.org/10.1016/j.bbagen.2013.06.025>
45. Tarantola M, Pietuch A, Schneider D, Rother J, Sunnick E, Rosman C, et al. Toxicity of gold-nanoparticles: Synergistic effects of shape and surface functionalization on micromotility of epithelial cells [Internet]. Vol. 5, *Nanotoxicology*. 2011. p. 254–68. Available from: <http://dx.doi.org/10.3109/17435390.2010.528847>
46. Schreiber SL, Crabtree GR. The mechanism of action of cyclosporin A and FK506 [Internet]. Vol. 13, *Immunology Today*. 1992. p. 136–42. Available from: [http://dx.doi.org/10.1016/0167-5699\(92\)90111-j](http://dx.doi.org/10.1016/0167-5699(92)90111-j)
47. Danda AK, Krishna TM, Narayanan V, Siddareddi A. Influence of primary and secondary closure of surgical wound after impacted mandibular third molar removal on postoperative pain and swelling--a comparative and split mouth study. *J Oral Maxillofac Surg* [Internet]. 2010 Feb [cited 2021 Sep 15];68(2). Available from:

<https://pubmed.ncbi.nlm.nih.gov/20116700/>

48. Ramadurai N, Gurunathan D, Samuel AV, Subramanian E, Rodrigues SJL. Effectiveness of 2% Articaine as an anesthetic agent in children: randomized controlled trial. *Clin Oral Investig* [Internet]. 2019 Sep [cited 2021 Sep 15];23(9). Available from: <https://pubmed.ncbi.nlm.nih.gov/30552590/>
49. Sathivel A, Raghavendran HR, Srinivasan P, Devaki T. Anti-peroxidative and anti-hyperlipidemic nature of *Ulva lactuca* crude polysaccharide on D-galactosamine induced hepatitis in rats. *Food Chem Toxicol* [Internet]. 2008 Oct [cited 2021 Sep 15];46(10). Available from: <https://pubmed.ncbi.nlm.nih.gov/18706469/>
50. Panda S, Doraiswamy J, Malaiappan S, Varghese SS, Del Fabbro M. Additive effect of autologous platelet concentrates in treatment of intrabony defects: a systematic review and meta-analysis. *J Investig Clin Dent* [Internet]. 2016 Feb [cited 2021 Sep 15];7(1). Available from: <https://pubmed.ncbi.nlm.nih.gov/25048153/>
51. Neelakantan P, Varughese AA, Sharma S, Subbarao CV, Zehnder M, De-Deus G. Continuous chelation irrigation improves the adhesion of epoxy resin-based root canal sealer to root dentine. *Int Endod J* [Internet]. 2012 Dec [cited 2021 Sep 15];45(12). Available from: <https://pubmed.ncbi.nlm.nih.gov/22612994/>
52. Govindaraju L, Neelakantan P, Gutmann JL. Effect of root canal irrigating solutions on the compressive strength of tricalcium silicate cements. *Clin Oral Investig* [Internet]. 2017 Mar [cited 2021 Sep 15];21(2). Available from: <https://pubmed.ncbi.nlm.nih.gov/27469101/>
53. Sekhar CH, Narayanan V, Baig MF. Role of antimicrobials in third molar surgery: prospective, double blind, randomized, placebo-controlled clinical study. *Br J Oral Maxillofac Surg* [Internet]. 2001 Apr [cited 2021 Sep 15];39(2). Available from: <https://pubmed.ncbi.nlm.nih.gov/11286448/>
54. DeSouza SI, Rashmi MR, Vasanthi AP, Joseph SM, Rodrigues R. Mobile phones: the next step towards healthcare delivery in rural India? *PLoS One* [Internet]. 2014 Aug 18 [cited 2021 Sep 15];9(8). Available from: <https://pubmed.ncbi.nlm.nih.gov/25133610/>
55. Nasim I, Neelakantan P, Sujeer R, Subbarao CV. Color stability of microfilled, microhybrid and nanocomposite resins--an in vitro study. *J Dent* [Internet]. 2010 [cited 2021 Sep 15];38 Suppl 2. Available from: <https://pubmed.ncbi.nlm.nih.gov/20553993/>
56. Danda AK, Muthusekhar MR, Narayanan V, Baig MF, Siddareddi A. Open versus closed treatment of unilateral subcondylar and condylar neck fractures: a prospective, randomized clinical study. *J Oral Maxillofac Surg* [Internet]. 2010 Jun [cited 2021 Sep 15];68(6). Available from: <https://pubmed.ncbi.nlm.nih.gov/20303209/>
57. Molecular structure and vibrational spectra of 2,6-bis(benzylidene)cyclohexanone: A density functional theoretical study. *Spectrochim Acta A Mol Biomol Spectrosc*. 2011 Jan 1;78(1):113–21.

58. Putchala MC, Ramani P, Herald J. Sherlin, Premkumar P, Natesan A. Ascorbic acid and its pro-oxidant activity as a therapy for tumours of oral cavity – A systematic review [Internet]. Vol. 58, Archives of Oral Biology. 2013. p. 563–74. Available from: <http://dx.doi.org/10.1016/j.archoralbio.2013.01.016>
59. Neelakantan P, Grotra D, Sharma S. Retreatability of 2 mineral trioxide aggregate-based root canal sealers: a cone-beam computed tomography analysis. J Endod. 2013 Jul;39(7):893–6.
60. Suresh P, Marimuthu K, Ranganathan S, Rajmohan T. Optimization of machining parameters in turning of Al-SiC-Gr hybrid metal matrix composites using grey-fuzzy algorithm [Internet]. Vol. 24, Transactions of Nonferrous Metals Society of China. 2014. p. 2805–14. Available from: [http://dx.doi.org/10.1016/s1003-6326\(14\)63412-9](http://dx.doi.org/10.1016/s1003-6326(14)63412-9)

UNDER PEER REVIEW