

Antiproliferative efficacy of Centella asiatica aqueous leaf extract on human lung cancer cell line.

ABSTRACT :

AIM: To assess the anti proliferative efficacy of Centella asiatica aqueous leaf extract on human lung cancer cell line

INTRODUCTION : Centella asiatica (family Apiaceae), commonly known as Indian Pennywort, is an ethnomedical plant that is widely used in India for treating skin problems and for revitalizing the brain and nervous system . It has been reported to have various pharmacological activities, including antioxidant, anti-inflammatory, anticancer, neuroprotective, cardioprotective, skin protective, radioprotective, immunomodulatory, memory-enhancing, and wound-healing properties. Asiatic acid and its derivatives possess a broad spectrum of pharmacological activities, such as anticancer, wound healing, anti-inflammatory, antidiabetic ,hepatoprotective and neuroprotective activities. The aim of the study is to assess the antiproliferative efficacy of Centella asiatica aqueous leaf extract on human lung cancer cell line.

MATERIALS AND METHODS : Human lung adenocarcinoma-A549 cell line was procured from National Centre for Cell Science (NCCS, Pune), India.

The effect of Centella asiatica on cell viability was measured by MTT assay following the method by Mosmann [2]. Briefly, the cells (1×10^5 cells per ml) were seeded in a 96 well micro titre plate (100 μ l per well) with replications. Treatment was conducted for 24 with different concentrations (50,100,150,200,250 and 300 μ M) of Centella asiatica .After incubation, 20 μ l of 5 mg/ml MTT stock solution was added to each well and incubated for 4 h at 37 $^{\circ}$ C. The obtained formazan crystals were solubilized with DMSO and the absorbance was measured at 570 nm using a microplate reader (SpectraMax M5, Molecular Devices, USA). Cell viability (%) has been shown as a ratio of absorbance (A570) in treated cells to absorbance in control cells (0.1 % DMSO) (A570). The IC50 was calculated as the concentration of sample needed to reduce 50 % of the absorbance in comparison to the DMSO-treated control. Percent cell viability was calculated following the equation: Cell viability (%) = $\{A570_{od} \text{ of (sample)}/A570_{od} \text{ of (control)}\} \times 100$

RESULTS AND DISCUSSION: Results show that centella asiatica has an antiproliferative effect on human lung cancer cell line. The characterisation of morphological changes in human lung cancer cells treated with Centella asiatica leaf extract compared to their respective controls were observed under phase contrast microscope. Visible destruction of the cancer cells were observed at the initial concentration of 25 microgram/ml. Further concentrations however showed increased antiproliferative activity proving it to be a potent anticancer drug.

CONCLUSION : This research concludes that centella asiatica leaf extract has a potent antiproliferative effect on human lung cancer cell lines. However further research is needed to understand the mechanisms of cytotoxicity of the plants.

INTRODUCTION:

Lung cancer is the most common type of cancer and is associated with a high rate of cancer-related mortality worldwide. Lung cancer is generally divided into two major subtypes: non-small-cell lung cancer (NSCLC) and small-cell lung cancer (1). Non-small cell lung cancer (NSCLC) types, including adenocarcinoma, squamous cell carcinoma and large cell carcinoma, account for more than 80% of total pulmonary malignancies (2). Although drugs targeting epidermal growth factor receptor (EGFR) mutations, such as gefitinib (Iressa®) and erlotinib (Tarceva®), have been approved for treatment of advanced NSCLC, their efficacy is limited, and more than 50% of these patients are not suitable for erlotinib and gefitinib treatment (3). Furthermore, almost all patients initially sensitive to gefitinib develop resistance to the drug (4). Therefore, it is essential to identify novel anticancer drugs or reagents against NSCLC. Centella asiatica (family Apiaceae), commonly known as Indian Pennywort, is an ethnomedical plant that is widely used in India for treating skin problems and for revitalizing the brain and nervous system (5). It has been reported to have various pharmacological activities, including antioxidant, anti-inflammatory (6), anticancer (7), neuroprotective, cardioprotective, skin, radioprotective, immunomodulatory, memory-enhancing, and wound-healing properties (8)(9). Studies on the chemical constituents of *C. asiatica* showed the presence of pentacyclic triterpenoids known as centelloids. These compounds are composed of terpene acids along with glycosides, including asiatic acid, asiaticoside, madecassic acid, madecassoside, brahmnic acid, brahmoside, brahminoside, thankinaside, isothankuniside, madasiatic acid, centic acid, centelloside, and cenellic acid (10).

Asiatic acid and its derivatives possess a broad spectrum of pharmacological activities, such as anticancer, wound healing, anti-inflammatory (11), antidiabetic (12), hepatoprotective and neuroprotective activities. Asiatic acid (AA) is a triterpene extracted from *Centella asiatica* (L.) Urban (Umbelliferae) that has a long history of successful use in both traditional Chinese and Indian Ayurvedic medicine. Previous studies have demonstrated that AA exhibits a variety of pharmacological effects not only as an antioxidant (13), antiinflammatory and neuroprotective agent but also against cancer (14). Medicinal herbs are a rich source of various bioactive compounds, which have long been used in the treatment of many diseases, including cancer. Recently, compounds isolated from medicinal herbs (15) have been identified as potent radiotherapy enhancers, such as radiosensitizers. Asiatic acid, asiaticoside, and madecassic acid have been reported as the major components of *C. asiatica* and these compounds have exhibited anticancer activity in various cancer cell lines. Asiatic acid has showed an antiproliferative effect by regulating apoptosis in a variety of human cancer cells, such as breast cancer, lung cancer, and melanoma cells (16). Our team has extensive knowledge and research experience that has translate into high quality publications(17–21),(22),(23),(24)(25),(26),(27),(19,28,29),(30–34).,(35),(36)

MATERIALS AND METHODS

CELL PROLIFERATION ASSAY

The effect of *Centella asiatica* on cell viability was measured by MTT assay following the method by Mosmann (37)

Principle

The assay is based on the reduction of soluble yellow tetrazolium salt to insoluble purple formazan crystals by metabolically active cells. Only live cells are able to take up the tetrazolium salt. The enzyme (mitochondrial dehydrogenase) present in the mitochondria of the live cells is able to convert internalized tetrazolium salt to formazan crystals, which are purple in colour. Then the cells are lysed using 20% SDS solution, which release the formazan crystal. These crystals are solubilized by DMF present in the solubilizer. The colour developed is then determined in an ELISA reader at 620 nm.

Reagents

1. MTT [3-(4,5-dimethylthiazol-2-yl) 2,5-diphenyl tetrazolium bromide]: 0.5 mg MTT/ml of serum-free DMEM.
2. Solubilization solution: 20% w/v SDS in 50% of Dimethyl formamide.
3. Phosphate Buffered Saline (PBS; pH 7.4).

Procedure

Briefly, the cells (1×10^5 cells per ml) were seeded in a 96 well micro titre plate (100 μ l per well) with replications. Treatment was conducted for 24 with different concentrations (50,100,150,200,250 and 300 μ M) of *Centella asiatica*. After incubation, 20 μ l of 5 mg/ml MTT stock solution was added to each well and incubated for 4 h at 37 °C. The obtained formazan crystals were solubilized with DMSO and the absorbance was measured at 570 nm using a microplate reader (SpectraMax M5, Molecular Devices, USA). Cell viability (%) has been shown as a ratio of absorbance (A₅₇₀) in treated cells to absorbance in control cells (0.1 % DMSO) (A₅₇₀). The IC₅₀ was calculated as the concentration of sample needed to reduce 50 % of the absorbance in comparison to the DMSO-treated control. Percent cell viability was

calculated following the equation: Cell viability (%) = {A570od of (sample)/A570 od of (control)} x 100

Spectrophotometrical absorbance of the purple blue formazan dye was measured in microplate reader at 620 nm. The OD of each sample was then compared with the control OD and the graph was plotted.

STATISTICAL ANALYSIS

All data obtained were analyzed by Students-t-test using MS-Excel, represented as mean \pm SD for six animals in each group. The results were computed statistically (SPSS/10 Software Package; SPSS Inc., Chicago, IL, USA) using one-way ANOVA. Post-hoc testing was performed for inter comparisons using the LSD. In all tests, the level of statistical significance was set at $p < 0.05$.

RESULTS AND DISCUSSION :

Non-small cell lung cancer (NSCLC) is the leading cause of cancer-related deaths worldwide. Although a number of drugs targeting EGFR mutations have been developed to date, most advanced cases are still incurable. Medicinal herbs are a rich source of various bioactive compounds, which have long been used in the treatment of many diseases, including cancer . Asiatic acid , asiaticoside and madecassic acid have been reported as the major components of *C. asiatica* (38) and these compounds have exhibited anticancer activity in various cancer cell lines. Asiatic acid has showed an antiproliferative effect by regulating apoptosis in a variety of human cancer cells, such as breast cancer, lung cancer, and melanoma cells(33) (39) (40) (41) (42) (43) (44) (45) (46) (47) (48) (49) (50) (51) (52). The characterisation of morphological changes in human lung cancer cells treated with *Centella asiatica* leaf extract compared to their respective

controls were observed under phase contrast microscope. Visible destruction of the cancer cells were observed at the initial concentration of 25 microgram/ml. Fig.1 shows the antiproliferative effect of centella asiatica as there is clear difference between the control cells and the destruction of A549 Cells that are treated with C.asiatica extract. The cells were treated with different concentrations such as 50, 100,150, 200, 250 and 300 $\mu\text{g/ml}$. As the concentration increased, the % of cell viability decreased gradually (fig.2) and showed increased antiproliferative activity proving it to be a potent anticancer drug.




Fig.1 shows the phase contrast microscopy of control cells and experimental A549 cells.

Antiproliferative efficacy of *Centella asiatica* aqueous leaf extract on human lung cancer cell line




Fig.2 Graph shows the antiproliferative activity of *Centella asiatica* where X axis represents concentration of *C. asiatica* extract and Y axis represents % of cell visibility.

CONCLUSION :

This research concludes that *centella asiatica* leaf extract has a potent antiproliferative effect on human lung cancer cell lines. However further research is needed to understand the mechanisms of cytotoxicity of the plants and fluorescent microscopy can be used distinguish between viable and non viable cells more clearly.

NOTE:

The study highlights the efficacy of "AYURVEDA" which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES:

1. James JT, Dubery IA. Pentacyclic triterpenoids from the medicinal herb, *Centella asiatica* (L.) Urban. *Molecules*. 2009 Oct 9;14(10):3922–41.
2. Bharadvaja N. *Centella Asiatica: A Pharmaceutically Important Medicinal Plant* [Internet]. Vol. 5, *Current Trends in Biomedical Engineering & Biosciences*. 2017. Available from: <http://dx.doi.org/10.19080/ctbeb.2017.05.555661>
3. Zalcman G, Bergot E, Lechapt E. Update on nonsmall cell lung cancer [Internet]. Vol. 19, *European Respiratory Review*. 2010. p. 173–85. Available from: <http://dx.doi.org/10.1183/09059180.00006610>
4. Ciardiello F, Tortora G. EGFR antagonists in cancer treatment. *N Engl J Med*. 2008 Mar

13;358(11):1160–74.

5. Cappuzzo F, Hirsch FR, Rossi E, Bartolini S, Ceresoli GL, Bemis L, et al. Epidermal Growth Factor Receptor Gene and Protein and Gefitinib Sensitivity in Non-Small-Cell Lung Cancer [Internet]. Vol. 97, JNCI: Journal of the National Cancer Institute. 2005. p. 643–55. Available from: <http://dx.doi.org/10.1093/jnci/dji112>
6. Jaisankar AI, Arivarasu L. Free Radical Scavenging and Anti-Inflammatory Activity of Chlorogenic Acid Mediated Silver Nanoparticle [Internet]. Journal of Pharmaceutical Research International. 2020. p. 106–12. Available from: <http://dx.doi.org/10.9734/jpri/2020/v32i1930715>
7. Karthik V, Arivarasu L, Rajeshkumar S. Hyaluronic Acid Mediated Zinc Nanoparticles against Oral Pathogens and Its Cytotoxic Potential [Internet]. Journal of Pharmaceutical Research International. 2020. p. 113–7. Available from: <http://dx.doi.org/10.9734/jpri/2020/v32i1930716>
8. Gohil KJ, Patel JA, Gajjar AK. Pharmacological Review on *Centella asiatica*: A Potential Herbal Cure-all. Indian J Pharm Sci. 2010 Sep;72(5):546–56.
9. Prakash V, Jaiswal N, Srivastava M. A REVIEW ON MEDICINAL PROPERTIES OF CENTELLA ASIATICA [Internet]. Vol. 10, Asian Journal of Pharmaceutical and Clinical Research. 2017. p. 69. Available from: <http://dx.doi.org/10.22159/ajpcr.2017.v10i10.20760>
10. Park BC, Bosire KO, Lee E-S, Lee YS, Kim J-A. Asiatic acid induces apoptosis in SK-MEL-2 human melanoma cells. Cancer Lett. 2005 Jan 31;218(1):81–90.
11. Thariny E, Arivarasu L, Rajeshkumar S. GREEN SYNTHESIS, ANTIOXIDANT AND ANTI-INFLAMMATORY ACTIVITY OF *Adathoda vasica* MEDIATED COPPER NANOPARTICLES. PLANT CELL BIOTECHNOLOGY AND MOLECULAR BIOLOGY. 2020 Nov 28;32–8.
12. Shree MK, Kavya Shree M, Arivarasu L, Rajeshkumar S. Cytotoxicity and Antimicrobial Activity of Chromium Picolinate Mediated Zinc Oxide Nanoparticle [Internet]. Journal of Pharmaceutical Research International. 2020. p. 28–32. Available from: <http://dx.doi.org/10.9734/jpri/2020/v32i2030726>
13. Shankar SB, Barani Shankar S, Arivarasu L, Rajeshkumar S. Biosynthesis of Hydroxy Citric Acid Mediated Zinc Nanoparticles and Its Antioxidant and Cytotoxic Activity [Internet]. Journal of Pharmaceutical Research International. 2020. p. 108–12. Available from: <http://dx.doi.org/10.9734/jpri/2020/v32i2630845>
14. Anand U, Jacobo-Herrera N, Altemimi A, Lakhssassi N. A Comprehensive Review on Medicinal Plants as Antimicrobial Therapeutics: Potential Avenues of Biocompatible Drug Discovery. Metabolites [Internet]. 2019 Nov 1;9(11). Available from: <http://dx.doi.org/10.3390/metabo9110258>
15. Herbal Sources Used by The Public Against Infections [Internet]. Vol. 12, International

Journal of Pharmaceutical Research. 2020. Available from:
<http://dx.doi.org/10.31838/ijpr/2020.sp1.015>

16. Nagoor Meeran MF, Goyal SN, Suchal K, Sharma C, Patil CR, Ojha SK. Pharmacological Properties, Molecular Mechanisms, and Pharmaceutical Development of Asiatic Acid: A Pentacyclic Triterpenoid of Therapeutic Promise. *Front Pharmacol.* 2018 Sep 4;9:892.
17. Rajeshkumar S, Kumar SV, Ramaiah A, Agarwal H, Lakshmi T, Roopan SM. Biosynthesis of zinc oxide nanoparticles using *Mangifera indica* leaves and evaluation of their antioxidant and cytotoxic properties in lung cancer (A549) cells. *Enzyme Microb Technol.* 2018 Oct;117:91–5.
18. Nandhini NT, Rajeshkumar S, Mythili S. The possible mechanism of eco-friendly synthesized nanoparticles on hazardous dyes degradation. *Biocatal Agric Biotechnol.* 2019 May 1;19:101138.
19. Vairavel M, Devaraj E, Shanmugam R. An eco-friendly synthesis of *Enterococcus* sp.–mediated gold nanoparticle induces cytotoxicity in human colorectal cancer cells. *Environ Sci Pollut Res.* 2020 Mar 1;27(8):8166–75.
20. Gomathi M, Prakasam A, Rajkumar PV, Rajeshkumar S, Chandrasekaran R, Anbarasan PM. Green synthesis of silver nanoparticles using *Gymnema sylvestre* leaf extract and evaluation of its antibacterial activity [Internet]. Vol. 32, *South African Journal of Chemical Engineering.* 2020. p. 1–4. Available from: <http://dx.doi.org/10.1016/j.sajce.2019.11.005>
21. Rajasekaran S, Damodharan D, Gopal K, Rajesh Kumar B, De Poures MV. Collective influence of 1-decanol addition, injection pressure and EGR on diesel engine characteristics fueled with diesel/LDPE oil blends. *Fuel.* 2020 Oct 1;277:118166.
22. Santhoshkumar J, Sowmya B, Venkat Kumar S, Rajeshkumar S. Toxicology evaluation and antidermatophytic activity of silver nanoparticles synthesized using leaf extract of *Passiflora caerulea*. *S Afr J Chem Eng.* 2019 Jul;29:17–23.
23. Raj R K, D E, S R. β -Sitosterol-assisted silver nanoparticles activates Nrf2 and triggers mitochondrial apoptosis via oxidative stress in human hepatocellular cancer cell line. *J Biomed Mater Res A.* 2020 Sep;108(9):1899–908.
24. Saravanan M, Arokiyaraj S, Lakshmi T, Pugazhendhi A. Synthesis of silver nanoparticles from *Phenerochaete chrysosporium* (MTCC-787) and their antibacterial activity against human pathogenic bacteria. *Microb Pathog.* 2018 Apr;117:68–72.
25. Gheena S, Ezhilarasan D. Syringic acid triggers reactive oxygen species–mediated cytotoxicity in HepG2 cells. *Hum Exp Toxicol.* 2019 Jun 1;38(6):694–702.
26. Ezhilarasan D, Sokal E, Najimi M. Hepatic fibrosis: It is time to go with hepatic stellate cell-specific therapeutic targets. *Hepatobiliary Pancreat Dis Int.* 2018 Jun;17(3):192–7.
27. Ezhilarasan D. Oxidative stress is bane in chronic liver diseases: Clinical and experimental

perspective. Arab J Gastroenterol. 2018 Jun;19(2):56–64.

28. Gomathi AC, Xavier Rajarathinam SR, Mohammed Sadiq A, Rajeshkumar S. Anticancer activity of silver nanoparticles synthesized using aqueous fruit shell extract of *Tamarindus indica* on MCF-7 human breast cancer cell line. J Drug Deliv Sci Technol. 2020 Feb 1;55:101376.
29. Dua K, Wadhwa R, Singhvi G, Rapalli V, Shukla SD, Shastri MD, et al. The potential of siRNA based drug delivery in respiratory disorders: Recent advances and progress. Drug Dev Res. 2019 Sep;80(6):714–30.
30. Ramesh A, Varghese S, Jayakumar ND, Malaiappan S. Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study. J Periodontol. 2018 Oct;89(10):1241–8.
31. Arumugam P, George R, Jayaseelan VP. Aberrations of m6A regulators are associated with tumorigenesis and metastasis in head and neck squamous cell carcinoma. Arch Oral Biol. 2021 Feb;122:105030.
32. Joseph B, Prasanth CS. Is photodynamic therapy a viable antiviral weapon against COVID-19 in dentistry? Oral Surg Oral Med Oral Pathol Oral Radiol. 2021 Jul;132(1):118–9.
33. Ezhilarasan D, Apoorva VS, Ashok VN. Syzygium cumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells. J Oral Pathol Med [Internet]. 2019 Feb [cited 2021 Sep 15];48(2). Available from: <https://pubmed.ncbi.nlm.nih.gov/30451321/>
34. Duraisamy R, Krishnan CS, Ramasubramanian H, Sampathkumar J, Mariappan S, Navarasampatti Sivaprakasam A. Compatibility of Nonoriginal Abutments With Implants: Evaluation of Microgap at the Implant-Abutment Interface, With Original and Nonoriginal Abutments. Implant Dent. 2019 Jun;28(3):289–95.
35. Gnanavel V, Roopan SM, Rajeshkumar S. Aquaculture: An overview of chemical ecology of seaweeds (food species) in natural products. Aquaculture. 2019 May 30;507:1–6.
36. Markov A, Thangavelu L, Aravindhan S, Zekiy AO, Jarahian M, Chartrand MS, et al. Mesenchymal stem/stromal cells as a valuable source for the treatment of immune-mediated disorders. Stem Cell Res Ther. 2021 Mar 18;12(1):192.
37. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. J Immunol Methods. 1983 Dec 16;65(1-2):55–63.
38. Du Q, Jerz G, Chen P, Winterhalter P. Preparation of Ursane Triterpenoids from *Centella asiatica* Using High Speed Countercurrent Chromatography with Step-Gradient Elution [Internet]. Vol. 27, Journal of Liquid Chromatography & Related Technologies. 2004. p. 2201–15. Available from: <http://dx.doi.org/10.1081/jlc-200025707>
39. Danda AK, Krishna TM, Narayanan V, Siddareddi A. Influence of primary and secondary

closure of surgical wound after impacted mandibular third molar removal on postoperative pain and swelling--a comparative and split mouth study. *J Oral Maxillofac Surg* [Internet]. 2010 Feb [cited 2021 Sep 15];68(2). Available from: <https://pubmed.ncbi.nlm.nih.gov/20116700/>

40. Ramadurai N, Gurunathan D, Samuel AV, Subramanian E, Rodrigues SJL. Effectiveness of 2% Articaine as an anesthetic agent in children: randomized controlled trial. *Clin Oral Investig* [Internet]. 2019 Sep [cited 2021 Sep 15];23(9). Available from: <https://pubmed.ncbi.nlm.nih.gov/30552590/>
41. Sathivel A, Raghavendran HR, Srinivasan P, Devaki T. Anti-peroxidative and anti-hyperlipidemic nature of *Ulva lactuca* crude polysaccharide on D-galactosamine induced hepatitis in rats. *Food Chem Toxicol* [Internet]. 2008 Oct [cited 2021 Sep 15];46(10). Available from: <https://pubmed.ncbi.nlm.nih.gov/18706469/>
42. Panda S, Doraiswamy J, Malaiappan S, Varghese SS, Del Fabbro M. Additive effect of autologous platelet concentrates in treatment of intrabony defects: a systematic review and meta-analysis. *J Investig Clin Dent* [Internet]. 2016 Feb [cited 2021 Sep 15];7(1). Available from: <https://pubmed.ncbi.nlm.nih.gov/25048153/>
43. Neelakantan P, Varughese AA, Sharma S, Subbarao CV, Zehnder M, De-Deus G. Continuous chelation irrigation improves the adhesion of epoxy resin-based root canal sealer to root dentine. *Int Endod J* [Internet]. 2012 Dec [cited 2021 Sep 15];45(12). Available from: <https://pubmed.ncbi.nlm.nih.gov/22612994/>
44. Govindaraju L, Neelakantan P, Gutmann JL. Effect of root canal irrigating solutions on the compressive strength of tricalcium silicate cements. *Clin Oral Investig* [Internet]. 2017 Mar [cited 2021 Sep 15];21(2). Available from: <https://pubmed.ncbi.nlm.nih.gov/27469101/>
45. Sekhar CH, Narayanan V, Baig MF. Role of antimicrobials in third molar surgery: prospective, double blind, randomized, placebo-controlled clinical study. *Br J Oral Maxillofac Surg* [Internet]. 2001 Apr [cited 2021 Sep 15];39(2). Available from: <https://pubmed.ncbi.nlm.nih.gov/11286448/>
46. DeSouza SI, Rashmi MR, Vasanthi AP, Joseph SM, Rodrigues R. Mobile phones: the next step towards healthcare delivery in rural India? *PLoS One* [Internet]. 2014 Aug 18 [cited 2021 Sep 15];9(8). Available from: <https://pubmed.ncbi.nlm.nih.gov/25133610/>
47. Nasim I, Neelakantan P, Sujeer R, Subbarao CV. Color stability of microfilled, microhybrid and nanocomposite resins--an in vitro study. *J Dent* [Internet]. 2010 [cited 2021 Sep 15];38 Suppl 2. Available from: <https://pubmed.ncbi.nlm.nih.gov/20553993/>
48. Danda AK, Muthusekhar MR, Narayanan V, Baig MF, Siddareddi A. Open versus closed treatment of unilateral subcondylar and condylar neck fractures: a prospective, randomized clinical study. *J Oral Maxillofac Surg* [Internet]. 2010 Jun [cited 2021 Sep 15];68(6). Available from: <https://pubmed.ncbi.nlm.nih.gov/20303209/>
49. Molecular structure and vibrational spectra of 2,6-bis(benzylidene)cyclohexanone: A

density functional theoretical study. *Spectrochim Acta A Mol Biomol Spectrosc.* 2011 Jan 1;78(1):113–21.

50. Putchala MC, Ramani P, Herald J. Sherlin, Premkumar P, Natesan A. Ascorbic acid and its pro-oxidant activity as a therapy for tumours of oral cavity – A systematic review [Internet]. Vol. 58, *Archives of Oral Biology.* 2013. p. 563–74. Available from: <http://dx.doi.org/10.1016/j.archoralbio.2013.01.016>
51. Neelakantan P, Grotra D, Sharma S. Retreatability of 2 mineral trioxide aggregate-based root canal sealers: a cone-beam computed tomography analysis. *J Endod.* 2013 Jul;39(7):893–6.
52. Suresh P, Marimuthu K, Ranganathan S, Rajmohan T. Optimization of machining parameters in turning of Al-SiC-Gr hybrid metal matrix composites using grey-fuzzy algorithm [Internet]. Vol. 24, *Transactions of Nonferrous Metals Society of China.* 2014. p. 2805–14. Available from: [http://dx.doi.org/10.1016/s1003-6326\(14\)63412-9](http://dx.doi.org/10.1016/s1003-6326(14)63412-9)

UNDER PEER REVIEW