

## Case study

### CASE REPORT ON CARCINOMA ON HARD PALATE

#### **Abstract:**

**Background:** The hard palate is a thin horizontal bony plate made up of two bones of the facial skeleton, located in the roof of the mouth. The bones are the palatine process of the maxilla and the horizontal plate of palatine bone. The hard palate spans the alveolar arch formed by the alveolar process that holds the upper teeth (when these are developed). If cancer is detected early, the overall 5-year survival rate for all persons is 85 percent. If the cancer has progressed to other tissues or organs, as well as regional lymph nodes, the overall 5-year survival rate is 67%. **Objective:** want to highlight the importance of primary prevention; education on risks of alcohol/tobacco use and oral hygiene. Secondary prevention; Early biopsy of any ulceration of the hard palate that does not regress with medical treatment. This would allow early diagnosis and minimal treatment with less morbidity and better survival chances. **Case presentation:** A male of 58 yrs., old came with Pain in moth, difficulty in breathing, difficulty in mastication, difficulty in deglutition, balm application, change in voice tooth exfoliation, lack of appetite. aggravates on mastication, difficulty in deglutition for 2 months, burning sensation on consumption of spicy food for 2 months change in voice, nasal discharge, loss of appetite, weight loss, tooth exfoliation in upper front region of jaw and balm application (2-3episodes, 15 days back). all necessary investigation carried out such as X- ray, MRI, CT scan, biopsy of tissue sample, and diagnosed Carcinoma on hard Palate. Treatment of this patient received antiemetic, analgesic, antibiotic, vitamin supplementary. **Prognosis:** After treatment, the patient shows great improvement. **Conclusion:** The therapeutic management for malignant tumors of the hard palate is essentially surgical, with or without postoperative radiotherapy, discussed on a case-by-case basis. Survival rate depends on several factors, including early diagnosis, histological characteristic and appropriate management. When ca of palate is detected primary, the management is very effective.

**Key words:** Carcinoma, bony plate, neoplasm staging, Alcohol drinking, smoking.

---

#### **INTRODUCTION**

Hard plate cancer (HPC) is an uncommon malignant tumour. Etiologic factors dominated by alcohol and tobacco consumption, are similar to those of other oral cavity cancers. The anatomical and histological constitution of the hard palate, with firm attachment of the mucosa to the underlying periosteum and the abundance of minor salivary glands, make the

hard palate a site of different histopathologic type of neoplasms. HPC represents approximately 1–3.5% of oral cavity cancers and is most often a squamous cell carcinoma.<sup>1</sup> A variety of treatments have been used to treat hard palate cancer, including surgery, radiotherapy, chemoradiation and combinations of these modalities.<sup>2</sup> Surgery is the treatment of choice.<sup>3</sup> It is usually performed via a transoral approach without the need to use facial incisions. The choice of the technique is based on location and size of the tumour in our work we present four cases of HPC treated by a transoral procedure.<sup>4</sup>

Hard palate cancer is a type of head and neck cancer that begins when cells that make up the bony part of the roof of the mouth grow out of control and form lesions or tumours. The hard palate creates a barrier between the mouth and the nasal cavity. Cancers that develop there tend to spread into the nasal cavity when they become more advanced. Hard palate squamous cell carcinoma develops from the thin, flat squamous cells lining the mucosal surface of the bone (periosteum) here. The exact cause of hard palate cancer is unknown. Tobacco use, alcohol consumption, precancerous lesions, and weakened immune system are the main risk factors for the formation of hard palate cancers.<sup>5</sup>

Using tobacco products and regularly drinking too much alcohol can increase your chances for developing hard palate cancer. Dentists are typically the first to notice the signs of hard palate cancer, often during a routine dental exam. The most common sign of hard palate cancer is an ulcer on the roof of the mouth. As the cancer grows, the ulcer may bleed. Other symptoms of hard palate cancer include the following: bad breath, loose teeth or pain around your teeth, dentures that no longer fit, changes in speech, difficulty swallowing, difficulty moving your jaw, a lump in the neck. Hard palate cancer is highly curable when diagnosed early. Treatment often involves surgery performed by a head and neck cancer surgeon.<sup>6</sup>

The goals of the treatment of hard palate cancer are to cure the cancer, preserve your appearance and the functions of your mouth, prevent the cancer from coming back. The extent and depth of the cancer guides your plan of care. An operation called a maxillectomy is often part of the treatment. If the cancer is more advanced, radiation, chemotherapy, or both may be used to shrink the tumour before or after surgery. This reduces the risk of the cancer coming back. For some people, radiation may be the only treatment needed.

The surgical treatment of malignant tumours of the hard palate must completely remove, with at least 1 cm margin, all the involved tissues with a resection of the mucosa and the underlying bone of the hard palate. In our study cervical lymph node metastases were found

in three patients, of whom one has Level I Lymph node metastasis with polymorphous low-grade adenocarcinoma and two had Level III lymph node metastasis with squamous cell carcinoma and mucosal melanoma as histological types. Surgical resection is performed trans-orally in almost all cases. The external approach is necessary for very advanced malignant neoplasms and when extending into the parapharyngeal space, pterygo-palatal fossa or masticatory space. Therapeutic neck dissection should be reserved for clinically and radiographically positive cervical lymph nodes. When unclear margins or perineural or perivascular invasion is present, with cervical lymph node involvement. Patients treated for hard palate cancer require a means of restoring the oral-nasal separation lost by removal of the palate and alveolar bone in order to improve patient's quality of life.<sup>7</sup>

Four major means of rehabilitation have been described in the literature. It includes: dental prosthetic when the cancer does not involve more than half the hard palate, mucoperiosteal local flap of palatal island flap, regional flaps by temporalis flap allows immediate reconstruction and can be used to correct unilateral and total defects. Otherwise, microvascular free tissue transfer is an option for larger defects.<sup>8</sup>

**PRESENTATION OF CASE:** A male of 58 yrs. old came with patient complaint of painful non healing ulcer in the upper front region of jaw. Patient was apparently alright months back when he experienced a painful non healing over upper front region of jaw over his left nostril area which was initially small in size and gradually increased to present size. Pain in mouth, difficulty in breathing, difficulty in mastication, difficulty in deglutition, balm application, change in voice tooth exfoliation, lack of appetite. aggravates on mastication, difficulty in deglutition for 2 months, burning sensation on consumption of spicy food for 2 months change in voice, nasal discharge, loss of appetite, weight loss, tooth exfoliation in upper front region of jaw and balm application (2-3episodes, 15 days back). History of asthma in the last 2 years, Patient is not on any medication for the same. No any history of hypertension, diabetes, jaundice, TB, or bleeding disorder. patient had past history of treatment in private hospital treatment. Patient first visited to a private hospital at Brahmapuri where Incisional Biopsy of lesion was done under LA on 5/10/21. and CT PNS was done. Patient was then referred to SPDC where slide and block review was done r/s/o "Moderately differentiated squamous cell carcinoma," All necessary investigation carried out such as X- ray, MRI, CT scan, biopsy of tissue sample, after testing Negative for COVID -19, patient got admitted to oral surgery ward for further management. Physical examination carried out There is not much abnormality found in head to foot examination, he is lean and thin and having dull look

and not active. He is weak and not cooperative. Hb%-9.9 gm%, RBC-3.63millions/cu.mm, WBC-8400/cu.mm, platelets-2.05 lacs/cu.mm, MCV-76.7n, MCH 27.7pico-gm, MCHC-35.6%, Monocytes-04%, Granulocytes-75%, Lymphocytes-20%, Eosinocytes-01%, Biasophiles-00%, urea 15mg%, creatine 0.6mgdl, sodium 133mEq/L, potassium 2.8mEq/L, albumin2.3gm/dl, total bilirubin 0.5mg/dl. and diagnosed Carcinoma on hard Palate. Treatment of this patient received antiemetic, analgesic, antibiotic, vitamin supplementary. Inj. Tramadol 50mg in 100 ml NS stat, Inj. dexamethasone 8mg IV, Inj. Emetrol 4mg BD, Inj. Neomol 1mg IV BD×3days, Inj. Perinorm 10mg IV BD×2 days, Inj. Ceftriaxone 1.5 gm IV BD×7 days, Inj. Metro 500 mg IV BD×5 days, Inj. Amikacin 500 mg IV BD×5 days, Inj. Pan 40 mg IV BD×7 days, Tab. Chymoral forte BD×7 days, Tab. Supradyn OD×7 days, Tab. Limcee 500 mg BD×15 days.

**PROGNOSIS:** After getting treatment, she shows great improvement and the treatment was still going on till my last date of care.

#### **DATA EXTRACTION**

Data Extracted from Handbook, PUB MED, Medline, And Cochrane Database Library.

#### **DISCUSSION:**

A male of 58 years from admitted to AVBRH with a rare case hard palate carcinoma. His weight is 45 kg and his height is 158 cm and BMI is 18.03. Admitted in hospital and all investigation and treatment were started. After getting treatment, she shows great improvement and the treatment was still going on. It is important to note that a patient could have one or more of these symptoms and not have hard palate cancer.<sup>9</sup> There are several non-cancerous causes of the same symptoms. That's why it's especially important to seek medical advice from a specialist. After diagnosing a patient with hard palate cancer (cancer involving the hard palate and the maxilla, or upper jaw), a doctor will need to determine what type of cancer it is, the grade of the tumour, and the stage of the cancer based on a biopsy or more often by the pathology after surgery.<sup>10</sup> It is important to note that oral cancers can sometimes be difficult to diagnose. If a doctor is having a hard time determining what type of cancer it is, he or she might ask for a second opinion and send some pieces of the tumour off to a specialist in head and neck pathology who deals more frequently with these types of tumors.<sup>11</sup> The most common type of hard palate cancer is squamous cell carcinoma. More than 90% of mouth cancers are squamous cell carcinoma. Squamous cell carcinoma is a cancer that starts from abnormal cells on the surface layer of the lips or the lining of the

mouth. If the cancer is discovered at an early stage, before invading past the deepest layer of the mouth lining, then it is called carcinoma in situ and has a good prognosis when removed. Another subtype is called verrucous carcinoma. This subtype usually has a slow growth pattern and is less likely to spread to lymph nodes in the neck or other parts of the body. For hard palate cancer, complete surgical removal of the tumour is almost always the first treatment, unless a doctor decides that it is not possible or safe to proceed with surgery. Hard palate cancers may be treated with a soft tissue resection, a maxillectomy, or a neck dissection.<sup>12</sup>

The surgery that a doctor recommends will depend on the location and extent of the cancer, as well as the stage. The management of the maxilla and hard palate bones, as well as the lymph nodes and other structures in the neck, are important factors to consider when planning a surgery for hard palate cancer.<sup>12</sup>

Restoration of function to this region is vitally important, since most cancer operations involving this region led to an opening in the palate and a loss of some of the upper teeth. Various forms of restoration of these types of defects have been employed and may include the use of a prosthesis, known as a palatal obturator, or any of a variety of reconstructive techniques. Patients and their care teams should discuss the types of surgeries that may be required for the treatment of their cancer.<sup>9</sup> The most common use of radiation for the treatment of hard palate cancer is called adjuvant radiation, which is radiation given after surgery in order to decrease the chances that the tumour will come back.

A doctor may recommend this type of post-surgical radiation for the following reasons, among others: If the tumour was not completely removed or if the surgical margins were positive for cancer. If the type of cancer was determined to be aggressive or of a high grade or T-stage. If the cancer had spread to lymph nodes or other structures, such as nerves or vessels. In some cases, complete surgical removal of a hard palate cancer may be impossible or unsafe, and a doctor may recommend radiation therapy as the primary treatment. In this type of treatment, an external beam of radiation is directed at the tumour in order to destroy the rapidly dividing cancer cells.

Chemotherapy is not commonly used to treat hard palate cancer. In some cases, it is used in combination with radiation as additional treatment following surgery. Chemotherapy is usually only added to adjuvant radiation therapy if there is extranodal extension from cancerous lymph nodes in the neck, or if there is cancer left behind during surgery.<sup>14</sup>

Patients should visit their head and neck specialist on a regular schedule (or earlier if they have any concerning symptoms). This allows doctors to monitor the patient for any sign that the cancer has returned. The best timeline for follow-up will be determined by the doctor.

#### Standard Follow-up Schedule

- For the first year, go every 1-3 months.
- For the second year, go every 2-6 months.
- For the third to fifth year, go every 4-8 months.
- After five years, start going once annually.

Doctors may select a scan to be performed in the first 6 months after treatment. The first scan serves as a “baseline” study for the purpose of comparing future studies. This will depend on the type and location of the cancer. Imaging could range from something as simple as a quick chest X-ray to more extensive tests such as a CT, MRI, or PET scan. If something suspicious comes up, a patient may need a biopsy.<sup>15</sup>

#### CONCLUSION

Squamous cell carcinoma is the most common malignant tumour of the hard palate. The clinical diagnosis is based on a search of risk factors, a complete ENT examination and endoscopic examination. CT scan allows a better analysis of bone invasion, while MRI retains its place in the evaluation of local and regional tumour extension. Surgical biopsy with immunohistochemical study confirms the diagnosis and exclude another diagnostic hypothesis.

Surgical excision is the treatment of choice. All our cases have been treated through an intraoral approach with elective neck dissection when pathological lymph nodes are found. Radiotherapy is considered in case of advanced stage disease, buccal extension, positive margins, perineural spread, or multiple lymph node metastases.

The reconstruction of defect after surgical resection is advisable. It restores almost immediately the phonetic and masticatory functions and it allows direct visualization of the primary site for recurrence detection as in case.<sup>16</sup>

#### CONSENT

As per international standard or university standard guideline patients consent and ethical approval has been collected and preserved by the authors.

## REFERANCE

1. Hammouda Y, Halily S, Oukessou Y, Rouadi S, Abada R, Roubal M, Mahtar M. Malignant tumors of the hard palate: Report of 4 cases and review of the literature. *International Journal of Surgery Case Reports*. 2020 Dec 16.
2. Pittman D. Recurrent Squamous Cell Carcinoma Hard Palate: Palatectomy and Reconstruction of Palate V/SNeoadjuvant Radiotherapy-A Case Study. *European Journal of Molecular & Clinical Medicine*. 2021 Mar 17;7(11):7179-83.
3. Pindborg JJ, Reichart PA, Smith CJ, Van der Waal I. *Histological Typing of Cancer and Precancer of the Oral Mucosa: In Collaboration with LH Sobin and Pathologists in 9 Countries*. Springer Science & Business Media; 2012 Dec 6.
4. Shah JP, Gil Z. Current concepts in management of oral cancer–surgery. *Oral oncology*. 2009 Apr 1;45(4-5):394-401.
5. Mahajan A, Kulkarni M, Parekh M, Khan M, Shah A, Gabhane M. Adenoid cystic carcinoma of hard palate: A case report. *Oral Maxillofac Pathol J*. 2011 Jan 1;2:127-31.
6. Kolokotronis A, Konstantinou N, Christakis I, Papadimitriou P, Matiakis A, Zaraboukas T, Antoniadis D. Localized B-cell non-Hodgkin's lymphoma of oral cavity and maxillofacial region: a clinical study. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology*. 2005 Mar 1;99(3):303-10.

7. Santos EP, Cavalcante DR, Melo AU, Pereira JC, Gomes MZ, Albuquerque RL. Plasmacytoid myoepithelioma of minor salivary glands: report of case with emphasis in the immunohistochemical findings. *Head & face medicine*. 2011 Dec;7(1):1-6.
8. Vranic S, Caughron SK, Djuricic S, Bilalovic N, Zaman S, Suljevic I, Lydiatt WM, Emanuel J, Gatalica Z. Hamartomas, teratomas and teratocarcinosarcomas of the head and neck: report of 3 new cases with clinico-pathologic correlation, cytogenetic analysis, and review of the literature. *BMC Ear, Nose and Throat Disorders*. 2008 Dec;8(1):1-0.
9. Napier SS, Speight PM. Natural history of potentially malignant oral lesions and conditions: an overview of the literature. *Journal of oral pathology & medicine*. 2008 Jan;37(1):1-0.
10. Giuliani M, Troiano G, Cordaro M, Corsalini M, Gioco G, Lo Muzio L, Pignatelli P, Lajolo C. Rate of malignant transformation of oral lichen planus: A systematic review. *Oral diseases*. 2019 Apr;25(3):693-709.
11. Miller AB, Hoogstraten BF, Staquet MF, Winkler A. Reporting results of cancer treatment. *cancer*. 1981 Jan 1;47(1):207-14.
12. Morita T, Tsunoda J, Inoue S, Chihara S. The Palliative Prognostic Index: a scoring system for survival prediction of terminally ill cancer patients. *Supportive care in cancer*. 1999 Apr;7(3):128-33.
13. Murphy BA, Gilbert J. Dysphagia in head and neck cancer patients treated with radiation: assessment, sequelae, and rehabilitation. In *Seminars in radiation oncology* 2009 Jan 1 (Vol. 19, No. 1, pp. 35-42). WB Saunders.
14. Freifeld AG, Bow EJ, Sepkowitz KA, Boeckh MJ, Ito JI, Mullen CA, Raad II, Rolston KV, Young JA, Wingard JR. Clinical practice guideline for the use of antimicrobial agents in neutropenic patients with cancer: 2010 update by the Infectious Diseases Society of America. *Clinical infectious diseases*. 2011 Feb 15;52(4):e56-93.
15. Scott SE, McGurk M, Grunfeld EA. The process of symptom appraisal: cognitive and emotional responses to detecting potentially malignant oral symptoms. *Journal of psychosomatic research*. 2007 Jun 1;62(6):621-30.