

# EVALUATION OF CHANGES IN HAEMATOLOGICAL PARAMETERS OF OBESE INDIVIDUAL AT OMISANJANA AREA OF ADO EKITI, EKITI STATE, NIGERIA

## ABSTRACT

**Background:** Obesity is a recurring illness which has reached all over the world and endangers public global health.

**Aim:** The study was done to evaluate the haematological parameters of obese individual at Omisanjana area of Ado Ekiti, Ekiti State. Nigeria.

**Materials and methods:** The study is a hospital based cross-sectional study among obese individual and non-obese individuals. This study was carried out at Omisanjana area of Ado Ekiti, Ekiti State. Fifty (50) obese individuals and fifty (50) apparently non-obese individuals were recruited as controls and enrolled in this study.

**Results:** The results showed significant difference of MXD ( $p=0.017$ ), haemoglobin ( $p=0.000$ ), MCV ( $p=0.006$ ), MCH ( $p=0.006$ ), MCHC ( $p=0.008$ ) and no significant difference in PCV ( $p=0.064$ ), WBC ( $p=0.896$ ), LYM ( $p=0.069$ ), GRAN ( $p=0.488$ ), RBC ( $p=0.820$ ), HCT ( $p=0.061$ ), PLT ( $p=0.819$ ), when compared between obese individuals and non-obese individuals respectively.

**Conclusion:** Subjects with obesity had lower mean haematological parameters such as haemoglobin mean cell volume, mean cell haemoglobin, mean cell haemoglobin concentration, neutrophils values compared to non-obese subjects. While a great majority of the parameters were of normal range for the subjects some were decreased e.g. total WBC count, Platelet and PCV.

**Keywords:** changes, BMI, haematological parameters, obese individual, Omisanjana Area

## INTRODUCTION

Obesity is a recurring illness which has reached all over the world and endangers public global health. Obesity is defined as body mass index (BMI) of  $30\text{kg/m}^2$ , while a value greater than  $40\text{kg/m}^2$  is considered extreme or morbid obesity [1]. Body Mass Index (BMI) is referred as the trendiest of many anthropometric indices. Indeed, it is accredited as an internationally accepted index for evaluating obesity [2] and is a degree of weight adjusted for height, calculated as weight in kilograms divided by the square of height in meters ( $\text{kg/m}^2$ ). Overweight and obesity are defined as abnormal or excessive fat accumulation in the body that may impair health, so Body Mass Index (BMI) of  $>25\text{ kg/m}^2$  and  $\geq 30\text{ kg/m}^2$  are considered to be overweight and obese respectively in adults irrespective of gender and age [2]. The phenomenon of obesity has drawn the attention of the scientific community, organizations and governments worldwide because it affects people's lives negatively and imposes excessive financial implications in every health system [3]. It is known that the obesity distribution is very huge worldwide. According to the World Health Organization's report in 2005, there were about 400 million obese adults worldwide and 1.6 billion overweight [4]. More than 1.1 billion adults are overweight, of which 312 million are obese. According to estimates of the International Obesity Task Force, 1.7 billion people are exposed to health risks related to body weight, while the increase in Body Mass Index (BMI) is responsible for more than 2.5 million deaths annually, which is expected to double by 2030 [5]. However, Dietary patterns such as a high consumption of nutrient-dense foods such as cereals, fruits, vegetables and low-fat meat and dairy products have been related to a number of favorable health outcomes in adults including a decreased prevalence of obesity [6].

Obesity is a chronic medical condition. It can lead to several untoward health effects, involving different organ systems [7]. Obesity is an important risk factor for hypertension, coronary heart disease and cerebrovascular diseases. The symptoms of obesity may resemble other medical problems or conditions. Psychological disturbances are also very common as well as stress, social pressure and doing developmental chores. Obesity is considered to be the root cause of many lifestyle diseases [8].

Obesity has been described as a state of low-grade inflammation [1], with leukocytes playing an important role in this state. Salma *et al.* [1] showed that a high WBC count in obese patients is associated with insulin resistance. Platelets, red blood cells (RBCs) and haemoglobin are associated with cardio respiratory conditions [9] in obese individuals. However, limited studies

have been conducted to describe any possible associations between body fat and haematological variables and so this topic is aimed at filling such gap by evaluating the haematological parameters of obese individual at Omisanjana area of Ado Ekiti, Ekiti State.

The study was done to evaluate the haematological parameters of obese individual at Omisanjana area of Ado Ekiti, Ekiti State. Nigeria.

## **MATERIALS AND METHOD**

### **Research design**

The study is a hospital based cross-sectional study among obese individual and non-obese individuals. The subjects were selected using a well-structured questionnaire who were age and sex matched.

### **Study area**

This study was carried out at Omisanjana area of Ado Ekiti, Ekiti State.

### **Target population**

This study was conducted at Omisanjana area of Ado Ekiti, Ekiti State. Fifty (50) obese individuals and fifty (50) apparently non-obese individuals were recruited as controls and enrolled in this study.

### **Blood collection**

Five (5ml) of venous blood was collected from each participant into an Ethylenediaminetetraacetic acid (EDTA) bottle which was then used for the determination of FBC.

**Body weight:** Body weight was measured while the subject minimally clothed and without shoes, standing steady on a weighing scale and it was recorded to the nearest 0.1kg.

**Height:** Height was measured to the nearest 0.1 cm while the subject was standing barefoot in erect position with a wall-mounted stadiometer.

**Body mass index:** BMI was measured by weight in kilograms divided by square of height in meters (kg/m<sup>2</sup>). (BMI in the range of 18.50 to 24.99 kg/m<sup>2</sup> is considered to be normal)

### **Method of test**

This was carried out using an automated analyser; KX-2IN (Sysmex Corporation, Kobe, Japan) Haematology analyser.

### **Method of data analysis**

The data were presented in tables and were presented as mean  $\pm$  standard deviation and analyzed using statistical packages for social sciences (SPSS, Version 20.0) and level of significance set at as  $p \leq 0.05$ .

### **Informed consent**

Informed consent was obtained from the subjects who participated in the study, the purpose of the study was explained to all participants. Participation in the study was entirely voluntary. Anonymity and confidentiality was ensured and maintained.

## **RESULTS**

**Table 1: Background Characteristics of obese individuals**

Demographic profile	Frequency (Percentages)
Gender	
Male	50(50%)
Female	50(50%)
Age in Years	
18-30	50(50%)

31-65	50(50%)
Education Qualification	
Primary	4 (8%)
Secondary	21(42%)
Tertiary	25(50%)

**Table 2: Mean values of haematological parameters of obese and non-obese subjects**

Parameter	Obese	Non-obese	t-value	P -value
PCV (%)	36.72±5.02	38.42±3.99	1.872	0.064
WBC( $10^9$ /L)	6.14±2.45	6.08±2.42	-0.131	0.896
LYM#( $10^9$ /L)	2.03±0.83	2.33±0.83	1.837	0.069
GRAN#( $10^9$ /L)	3.94±3.85	3.52±1.87	-0.695	0.488
MXD#( $10^9$ /L)	0.83±1.20	0.41±0.02	-2.431	0.017*
RBC( $10^9$ /L)	4.52±0.83	4.55±0.58	0.228	0.820

HGB(g/dl)	11.15±3.19	12.99±1.55	3.670	0.000*
HCT(%)	36.66±5.06	38.40±4.02	1.899	0.061
MCV(fl)	79.13±10.37	84.82±7.02	3.210	0.002*
MCH(Pg)	26.65±3.87	28.60±2.97	2.827	0.006*
MCHC(g/dl)	32.20±2.00	33.39±2.33	2.727	0.008*
PLT(10 <sup>9</sup> /L)	173.53±71.46	176.74±67.46	0.230	0.819

The results showed significant difference of MXD ( $0.83 \pm 1.20 \times 10^9/L$ ,  $0.41 \pm 0.02 \times 10^9/L$ ,  $p=0.017$ ), HGB ( $11.15 \pm 3.19g/dl$ ,  $12.99 \pm 1.55g/dl$ ,  $p=0.000$ ), MCV ( $79.13 \pm 10.37fL$ ,  $84.82 \pm 7.02fL$ ,  $p=0.006$ ), MCH ( $26.65 \pm 3.87Pg$ ,  $28.60 \pm 2.97Pg$ ,  $p=0.006$ ), MCHC ( $32.20 \pm 2.00g/dl$ ,  $33.39 \pm 2.33g/dl$ ,  $p=0.008$ ) and no significant difference in PCV ( $36.72 \pm 5.02\%$ ,  $38.42 \pm 3.99\%$ ,  $p=0.064$ ), WBC ( $6.14 \pm 2.45 \times 10^9/L$ ,  $6.08 \pm 2.42 \times 10^9/L$ ,  $p=0.896$ ), LYM ( $2.03 \pm 0.83 \times 10^9/L$ ,  $2.33 \pm 0.83 \times 10^9/L$ ,  $p=0.069$ ), GRAN ( $3.94 \pm 3.85 \times 10^9/L$ ,  $3.52 \pm 1.87 \times 10^9/L$ ,  $p=0.488$ ), RBC ( $4.52 \pm 0.83 \times 10^9/L$ ,  $4.55 \pm 0.58 \times 10^9/L$ ,  $p=0.820$ ), HCT ( $36.66 \pm 5.06\%$ ,  $38.40 \pm 4.02\%$ ,  $p=0.061$ ), PLT ( $173.53 \pm 71.46 \times 10^9/L$ ,  $176.74 \pm 67.46 \times 10^9/L$ ,  $p=0.819$ ), when compared between obese individuals and non-obese individuals respectively.

## DISCUSSION

Results obtained show reduction in haemoglobin, Mean Cell volume (MCV), Mean Cell haemoglobin (MCH) and Mean Cell haemoglobin concentration (MCHC). These were in contrast to Mei-Chu *et al.* [10] that suggested that red blood cell (RBC) count and haematocrit were positively associated with obesity. This shows anaemia in obese individuals and may lead to microcytic hypochromic anaemia. This situation could lead to cardiovascular disease. Also, previous literature has traced increased packed cell volume (PCV) in obesity to be one of the risk factors for cardiovascular and other diseases. Furthermore, PCV is the most important indicator to determine viscosity of the blood although; there was no change in PCV of the subjects in this study. Viscosity of the blood is an indicator of vascular risks, and increased body mass index (BMI) is known to increase viscosity of the blood [11] and the lower value may indicate that the subjects of the study are not at risk of cardiovascular disease.

Leukocytosis has often been linked with atherosclerotic disease and has also been accepted as a risk factor for cardiovascular disease (CVD) [12]. Haematological parameters have established as indicators for health and disease [13, 14]. The association between leukocyte count and risk of atherosclerotic disease is plausible because leukocytes present a major contribution to the rheologic properties of blood. This is achieved by altering the adhesive properties under stress and participating also in the case of endothelial injury. However, there is no significant different of WBC of obese individual with non- obese individual while other parameters are significantly differently different. Obesity per se is not independently associated with altered RBC, Hb and Ht,

and the association between BMI and hematological parameters is mediated by their associations with abdominal fat and insulin resistance markers [15].

## CONCLUSION

Subjects with obesity had lower mean haematological parameters such as haemoglobin mean cell volume, mean cell haemoglobin, mean cell haemoglobin concentration, neutrophils values compared to non-obese subjects. While a great majority of the parameters were of normal range for the subjects some were decreased e.g. total WBC count, Platelet and PCV.

Knowledge of the relationship between BMI and hematological indices of apparently healthy individuals within any population is therefore crucial for healthcare planning, as a justification for appropriate early prognosis and important counseling policy strategically important for reducing the incidence of obesity in the country. Obese individual should be aware of the risk of them developing cardiovascular disease due to the viscosity of their blood and as such haematological parameters such as full blood count would be a great diagnostic pointer to such disease condition. Thus, it becomes pertinent for the obese individual to assess their haematological parameters at nearby laboratory on monthly bases as a routine checkup to curb the risk of developing arthelerosclerosis which may aggravate as a result of obesity.

## REFERENCES

1. Salma S, Alqassas M, Othman S, Shihon SB, Alqalayta L, Alqusair S, Mohamad Q. . Obesity correlates with neutrophilia. *Hematology Transfusion International Journal*.2016; **3**(2):159–162.
2. Olutayo IA, David BA, Adaobi MO. Correlation between Body Mass Index and Hematological Indices in Young Adult Nigerians with Different Hemoglobin Genotypes.*American Journal of Biomedical Sciences*. 2017; **9**(1), 38-46.
3. Polikandrioti M, Stefanou E..Obesity Disease.*Health Science Journal*. 2009; **3**(3)132-136.
4. Low S, Chin MC, Ma S, Heng D, Deurenberg-Yap M..Rationale for redefining obesity in Asians.*AnnalsAcademy of Medicine Singapore*. 2009; **38**(1): 66-69.
5. Berghöfer A, Pischon T, Reinhold T, Apovian CM, Sharma AM, Willich SN. Obesity prevalence from a European perspective: a systematic review. *Bio Medical CentralPublic Health*. 2008; **5**(8):200.
6. Yanoff LB, Menzie CM, Denkinger BI..Inflammation and iron deficiency in the hypoferremia of obesity.*International Journal of Obesity*.2007; **31**(9):1412–1419.

7. Ranjani H, Mehreen TS, Pradeepa R, Anjana RM, Garg R, Anand K, Mohan V. Epidemiology of childhood overweight and obesity in India: a systematic review. *Indian Journal of Medical Research*. 2016; **143**(2): 160-174.
8. Chambers JC, Elliott PZ, Delilah ZW. Common genetic variation near MC4R is associated with waist circumference and insulin resistance. *Nature Genetics*. 2017; **71**(7):850-857.
9. Zamai CA, Rodrigues AA, Filocomo M, Braga LE, de Oliveira JF, Zago C. Avaliação da aptidão cardio-respiratória através do teste de caminhada em esforço controlado (TCEC). *Conexões: Rev Fac Educ Fis*. 2010;8(2):146-58.
10. Mei-Chu YJ, Chia-Chang H, Wei-Chin H, Yung-Chuan L, Chao-Ping W, Ting T, Cheng L, Yi-Hsin H, Fu-Mei C, Yau-Jiunn L, Teng-Hung Y. Association between lifestyle and hematological parameters: A study of Chinese male steelworkers. *Journal of Clinical Laboratory Analysis*. 2019; 33(e22946)1-9.
11. Salih K, Jouda J, El-Haboby B. Obesity average and its relation with packed cell volume (PCV) value in a section of students and employees of Al-Mustansiriyah University in Iraq. *Journal of Biology, Agriculture and Healthcare*. 2016; **6**(6): 24-29.
12. Madjid M, Awan I, Willerson JT, Casscells SW. Leukocyte Count and Coronary Heart Disease; Implications for Risk Assessment, *JACC*. 2004; **44**(10): 1945–1956.
13. Obeagu EF, Obeagu GU, Chijioke UO, Ofor IB (2017) Analysis of Alterations in Selected Haematological Parameters of Ascariasis Patients in Umudike, Abia State, Nigeria. *Ann Clin Lab Res*. 2017; **5**(3):193. DOI: 10.21767/2386-5180.1000193.
14. Obeagu EI, Azuonwu O, Didia BC and Obeagu GU. Determination of Haematological Changes Associated with Syphilis in Subjects in Umudike, Abia State, Nigeria. *Cohesive Journal of Microbiology and Infectious Disease*. 2018; **1**(1): 505.
15. Barazzoni R, Gortan Cappellari G, Semolic A, Chendi E, Ius M, Situlin R, et al. The Association between Hematological Parameters and Insulin Resistance Is Modified by Body Mass Index – Results from the North-East Italy MoMa Population Study. *PLoS ONE*. 2014; **9**(7): e101590. <https://doi.org/10.1371/journal.pone.0101590>