

### CASE REPORT ON HODGKIN'S LYMPHOMA

#### Abstract

**Introduction:** Hodgkin lymphoma, which also accounts up 11% of all lymphomas, is named Hodgkin's disease. This malignant disease is known as the Reed-Sternberg-cell, which is situated in lymph-nodes and is characterized by the proliferation of irregular gait. The disease occurs more often in those between 15-30 years and over 55 and is double prevalent in women, in men and women. The sickness is bimodal age specific. The frequency in India is around 1 in 25,000 inhabitants. However, for all stages, long-term survival is around 80%. **Clinical findings:** The onset of symptoms is usually insidious but usually used for frequent diagnosis are: Extension of lymph nodes in area of cervical, axillary or inguinal, extraordinary weight loss, Fatigue, Weakness, fever, chills, tachycardia, sweating or nocturnal sweat. **Diagnostic evaluation:** Peripheral analysis of the blood, Biopsy of Excision of lymph node, Examination of bone marrow, Radiological study **Therapeutic investigation:** Chemotherapy and radiation therapy or some may need immunotherapy or stem cell transplant. **Outcome:** After treatment, the child showed great improvement, her low grade fever and night sweats. **Conclusion:** A male child 11 year was admitted to pediatric ward no.22 in AVBRH on 25/05/2021 with a unique case of Hodgkin's lymphoma. She was admitted in hospital and all investigation and treatment were started. After getting treatment, she shows great improvement and the treatment was still going on till my last date of care.

**Key words:** Hodgkin's lymphoma, Child (female)

#### Introduction

Hodgkin lymphoma, which also accounts up 11% of all lymphomas, is named Hodgkin's disease. (1)This malignant disease is known as the Reed-Sternberg-cell, which is situated in lymph-nodes and is characterized by the proliferation of irregular gait. (2)The disease occurs more often in those between 15-30 years and over 55 and is double prevalent in women, in men and women.

The sickness is bio-modal age specific.(3) The frequency in India is around 1 in 25,000 inhabitants. However, for all stages, long-term survival is around 80%.(4)

**Patients Identification:** A male child 11 year was admitted to pediatric ward no.22 in AVBRH on 25/05/2021 with a complaint of Left hip discomfort for 2 months. In the last 3 months, he has recorded heavy night sweats, mild temperature, pruritus rash and a roughly 15% weight loss. In the last month, also, he noticed a growing lump in his left hip.

**Present medical history:** A male child 11 year was admitted to pediatric ward no.22 in AVBRH on 25/05/2021 with a complaint of Left hip discomfort for 2 months. In the last 3 months, he has recorded heavy night sweats, mild temperature, pruritus rash and a roughly 15% weight loss. In the last month, also, he noticed a growing lump in her left hip. Also, he found an enlarging mass in his left hip during the latest month at presently he is well and is under observation with continuous medication.

**Past medical history:** The past medical history of my patient she have swelling on right side of neck, light pain. His ongoing general treatment in private hospital. He was admitted to hospital due to increase swelling. Till then, she was undergone on treatment in AVBRH Sawangi (M) Wardha.

**Family history:** There are 4 members in his family. All other members of the family were not having any complaints in their health. In his family no other member is having any problem of hypertension, diabetes or any other disease condition.

**Past investigation and Outcome:** The past medical history of my patient he have swelling on right side of neck, light pain... His ongoing general treatment in private hospital. He was admitted to hospital due to increase swelling and fever. Till then, he was undergone on treatment in AVBRH Sawangi (M) Wardha.

**Clinical findings:** Swelling in right side of neck and Left hip discomfort for 2 months. In the last 3 months, she has recorded heavy night sweats, mild temperature, pruritus rash and a roughly 15% weight loss. In the last month. In the last month, also, he noticed a growing lump in his left hip.

**Etiology:** Unknown is the exact etiologic agent of Hodgkin lymphoma. There is however an elevated risk of “Hodgkin's lymphoma” for the illness, autoimmune disorders and immunosuppression of Epstein-Barr (EBV). Family susceptibility to Hodgkin's lymphoma has also demonstrated. In the mixed cell and lymphocyte deficient subtypes of Hodgkin's lymphoma EBV has been reported to be more prevalent. The potential cause of EBV illness loss of the immune surveillance was postulated. No other virus played a significant role in pathogenesis of the illness. .The increased risk of developing Hodgkin lymphoma includes secondary immune suppression following strong organ or hematopoietic cell transplants, therapy with immunosuppressive medications, and human immunodeficiency (HIV). HIV patients usually have a high stage, atypical lymph nodes and bad prognostics. Studies have revealed that HL growth in homosexual siblings in patients with Hodgkin lymphoma has been ten-fold, indicating a gene-environmental involvement in the susceptibility of Hodgkin's lymphoma.

**Physical examination:** There is not much abnormality found in head to toe examination, the child is lean and thin and having dull look. He is weak and not so cooperative. Though it is found that the child is having swelling on right-side of neck.

**Diagnostic assessment:** Blood test: Hb – 10.6 gm.%, Total RBC count – 3.87millions/cumm, RDW – 15.3%, HCT – 33.5%, Total WBC count – 6600/cumm, Monocytes – 08%, Granulocytes – 75%, Lymphocytes – 15%, AST(SGOT) – 46U/L. Peripheral Smear: RBCs –Normocytic, normonucliotic. Platelets – adequate on smear. No Haemoparasite seen.

**Ultrasonography:** No obvious retroperitoneal lymphadenopathy seen.

**Therapeutic Intervention:**Blood transfusion, Inj. Ceftriaxone 500mg IV x BD, Tab. Paracetamol 500mg x OD Inj. Neomol 100mg x BD, Inj. Pantop 40mg x OD. IVF - NS, RL.

**Discussion:**A male child admitted in AVBRH on 25/05/2021 in pediatric ward no 15 with chief complaint of Swelling in right side of neck and Left hip discomfort for 2 months, history of night sweating, low grade fever, pruritus rash across the body, and weight loss of about 15% during the last three months. In the last month, he also detected a growing lump in her left hip. As soon as he was admitted to hospital investigations were done and appropriate treatment were started. After getting treatment, he shows great improvement and the treatment was still going on till my last date of care.

In a study published "On a malignant appearance of the absorbing glands and spleen" Thomas Hodgkin gave the earliest description of "Hodgkin's lymphoma" in 1832. The name "Hodgkin's disease" was created by Samuel Wilk in 1856. (5) 6. Epstein Barr Classical Virus Association In developmental children with a frequency of 91 to 98 per cent, Hodgkin's lymphoma is prevalent. (6) About 4.5% of patients are susceptible to hereditary disease. Of course, in humans infected by immunodeficiency viruses, there is a 10-fold increase in the incidence of classical lymphoma of Hodgkin. It may be linked to autoimmune disorders and can develop lymphoma of Non-Hodgkin. (7)

Neoplastic cells make up less than 1% in the lymph node of Hodgkin's lymphoma. While cells of similar phenotypes but of various morphologies are widespread, classic reed-sternberg cells are rather unusual, as are Hodgkin/Reed-Sternberg cells. (8) Immunophenotypic investigations reveal that B cells in the germ center of the lymph node are generated from (CD15); positive (CD30) cells. The etiology of Hodgkin lymphoma is significant for interaction between Hodgkin/Reed-Sternberg cells and the surrounding inflammatory cells. (9)

Pediatric Occurrence of Hodgkin's cervical lymphatic lymphoma is usually unilateral, painless. The main site is subdiaphragmatic in 3 percent of the cases. Those that are mediastinal with chronic toxicity (10). Rarely, an axillary or inguinal node may occur with splenomegaly; 4. 'B' symptoms include a fever above 38°C, nocturnal sweat drenching and unclear loss of body weight more than 10 percent in the last six months in roughly 50 percent of developing country youngsters. (11)

**Conclusion:** : The prognosis and increasing prevalence in developing countries of juvenile Hodgkin lymphoma appear to be worse and general professional should be conscious of early clinical events. In order to identify Hodgkin's lymphoma early, there is a high suspicion index, as many youngsters are in severe stages. The extent of illness is defined by Radiological Studies. In the early evaluation of a kid with "Hodgkin's lymphoma" with his to-pathological exam, fine needle-aspiration plays the crucial function of providing a definite immune-supplemented diagnosis. The scale of the condition helps stratify risks and determine drug treatment options. My patient show great improvement after getting the treatment and the treatment was still going on till my last date of care.

**Ethical approval:** This information was obtained from the institution's ethics committee.

**Reference:**

1. What Is Hodgkin Lymphoma? | Define Hodgkin Lymphoma [Internet]. [cited 2021 Nov 24]. Available from: <https://www.cancer.org/cancer/hodgkin-lymphoma/about/what-is-hodgkin-disease.html>
2. Hodgkin's Lymphoma - Physiopedia [Internet]. [cited 2021 Nov 24]. Available from: [https://www.physio-pedia.com/Hodgkin%27s\\_Lymphoma](https://www.physio-pedia.com/Hodgkin%27s_Lymphoma)
3. McDonald CM. CLINICAL APPROACH TO THE DIAGNOSTIC EVALUATION OF HEREDITARY AND ACQUIRED NEUROMUSCULAR DISEASES. *Phys Med Rehabil Clin N Am*. 2012 Aug;23(3):495–563.
4. Understanding Statistics Used to Guide Prognosis and Evaluate Treatment | Cancer.Net [Internet]. [cited 2021 Nov 24]. Available from: <https://www.cancer.net/navigating-cancer-care/cancer-basics/understanding-statistics-used-guide-prognosis-and-evaluate-treatment>
5. Pediatric Lymphomas | SpringerLink [Internet]. [cited 2021 Nov 24]. Available from: <https://link.springer.com/book/10.1007%2F978-3-540-68753-5>
6. Frequency of EBV associated classical Hodgkin lymphoma decreases over a 54-year period in a Brazilian population [Internet]. [cited 2021 Nov 24]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5789833/>
7. HIV infection and lymphoma [Internet]. [cited 2021 Nov 24]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2095580/>
8. Hodgkin's lymphoma: the pathologist's viewpoint [Internet]. [cited 2021 Nov 24]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1769601/>
9. Classical Hodgkin Lymphoma - an overview | ScienceDirect Topics [Internet]. [cited 2021 Nov 24]. Available from: <https://www.sciencedirect.com/topics/medicine-and-dentistry/classical-hodgkin-lymphoma>
10. Malignant pulmonary and mediastinal tumors in children: differential diagnoses [Internet]. [cited 2021 Nov 24]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2967153/>
11. Lymphoma - Hodgkin: Symptoms and Signs | Cancer.Net [Internet]. [cited 2021 Nov 24]. Available from: <https://www.cancer.net/cancer-types/lymphoma-hodgkin/symptoms-and-signs>

UNDER PEER REVIEW