

Original Research Article

Gender Difference in Blood Pressure Control and Cardiovascular Risk Factors With Diagnosed Hypertension

Abstract:

Objective: Our study was designed to compare the gender difference in Blood Pressure Control and Cardiovascular Risk Factors Mohammad Bin Khalifa bin Salman Al Khalifa Specialist Cardiac Centre Bahrain population With Diagnosed Hypertension.

Methodology: This cross-sectional study was conducted in Mohammad Bin Khalifa bin Salman Al Khalifa Specialist Cardiac Centre Bahrain from December 2019 to December 2020. Blood pressure was measured twice by trained physicians using aneroid sphygmomanometers after a standardized protocol. Patients were asked to sit with both feet on the floor for ≥ 5 minutes before the first BP measurement. Both the two BP measurements were taken 60 seconds apart. For this research we defined hypertension as systolic BP ≥ 140 mm Hg, diastolic BP ≥ 90 mm Hg. All the participants were divided into two groups normotensive participants, having no medical history of hypertension, and hypertensive participants, those who reported hypertension with antihypertensive medications use. Weight status was another variable for analyzing risk factors. We defined weight factor by body mass index as defined by WHO experts for overweight (BMI ≥ 24 kg/m²) and obesity (BMI ≥ 28 kg/m²).

Results: We observed that the male population reported a high prevalence of cardiovascular risk factors as compared to men in terms of obesity, total cholesterol level and also reported low HDL levels both in hypertensive and normotensive groups. However, in females, the HDL-C remained similar in both hypertensive and normotensive groups. The prevalence of hypertension increased in the men population with increasing BMI due to unhygienic food and poor diet plans.

Conclusion: Hypertension is one of the leading causes of cardiovascular disorders in many regions of the world. Our results indicated significant sex-related homogeneity in terms of hypertension.

Keywords:

Gender prevalence, Cardiovascular risk Factors, Hypertension

Introduction:

Rapid urbanization, aging, poor lifestyle, unhealthy diet patterns, and globalization are one of the most challenging disorders in developing countries which lead to hypertension disorder¹. Hypertension become a world emergency especially in developing countries that need proper treatment to reduce the risk of cardiovascular events and occurrence². In the modern world, hypertension is one of the major challenging disorders leading to heart attack, stroke, and vascular complications. The high mortality ratio due to cardiovascular disease usually happened due to elevated blood pressure levels³. In 2008, approximately 1 billion deaths were reported in the world adult population due to hypertension⁴. Medical experts claims that in 2025, hypertension become an alarming situation that may cause 1.7 billion adult death. In low-income countries, annually 6 million deaths are reported due to hypertension⁵. Hypertension is now identified as a major contributor to disease burden in many parts of the world. Nearly 2/3rd hypertensive patients live in underdeveloped countries. One of the studies stated that 3.4% male adult population and 6.8% female population are suffering from hypertension worldwide enhances the risk of cardiovascular disorders⁶. In 2000, 2-4% urban population of low-income Asian countries were suffering from hypertension in which their blood levels ranges $\geq 160/\geq 95$ mmHg⁷. The global burden of disease claimed that the cardiovascular mortality ratio would increase in 2023 due to hypertension. Benjamin's study reported a low prevalence of hypertension in women younger than 65 years of age group when compared to men⁸. But when women reached above 65 years of age majority of them face elevated blood pressure levels in their elderly life period⁸. Our study was designed to compare the gender difference in Blood Pressure Control and Cardiovascular Risk Factors in population With Diagnosed Hypertension.

Methodology:

This cross-sectional study was conducted in Mohammad Bin Khalifa bin Salman Al Khalifa Specialist Cardiac Centre Bahrain from December 2019 to December 2020. In this study, BP screening was done among the adult population aged 18 years or over. Before initiating the research, ethical approval was obtained from the hospital research ethics committee and research was conducted by following Helsinki principles. All the participants were well-known about the objectives and nature of the research. Written and verbal consent were obtained from every participant. The female proportion of participants aged <40 years was 10.8% and male participants were 4.7%. At the initial survey majority of the young participant's withdrawal from the study so for the final analysis participants ≥ 40 years were included. Survey analysis

was used for gathering data related to Cardiovascular Disease Risk Factors. The questionnaire was obtained from the WHO MONICA and used for further analysis^{9,10}. Blood pressure was measured twice by trained physicians using aneroid sphygmomanometers after a standardized protocol. Patients were asked to sit with both feet on the floor for ≥ 5 minutes before the first BP measurement. Both the two BP measurements were taken 60 seconds apart. For this research we defined hypertension as systolic BP ≥ 140 mm Hg, diastolic BP ≥ 90 mm Hg. We also used antihypertensive medication under the definition of hypertension. On the other hand, we defined BP control as diastolic BP < 90 mm Hg and systolic BP ≥ 140 mm Hg. All the participants were divided into two groups normotensive participants, having no medical history of hypertension, and hypertensive participants, those who reported hypertension with antihypertensive medications use. Weight status was another variable for analyzing risk factors. We defined weight factor by body mass index as defined by WHO experts for overweight (BMI ≥ 24 kg/m²) and obesity (BMI ≥ 28 kg/m²)¹¹. Participants who currently smoking more than one cigarette and hookah were categorized as current daily smokers. At the time of the survey, blood samples were taken from each participant for analyzing levels of serum total cholesterol (TC), triglyceride (TG). Furthermore, laboratory analyses of low-density lipoprotein cholesterol (LDL-C), high-density lipoprotein cholesterol (HDL-C), urea, creatinine, and uric acid (UA) were also performed.

Statistical analysis was performed by using SPSS 23.0. Continuous variables were measured in mean and standard deviations whereas a two-tailed student t-test was used to compare the differences between the two groups. Categorical variables were measured by using the Chi-square test whereas column proportion was measured by using the z test. To determine independent risk factors of hypertension we used multivariate binary logistic regression with a 95% confidence interval for adjusted odds ratios.

Results:

In this study total of 2652 patients were recruited of which 72% were female and 28% were male. We observed that females aged 50 or above had a high frequency of hypertension as compared to men. However, overall the weighted prevalence indicated high hypertension in the male population (61.2% in men vs. 48.8% in the women population). The prevalence of hypertension increased with age in both men and women. But the hypertension prevalence of males was lower than female participants at 60-69 and 70-79. We observed that hypertension prevalence remained unchanged in the male population but continuously shifted in female participants. total cholesterol level and table 2 represents the similar systolic and diastolic blood pressure among male and female populations. In the table 3, We observed that the male population reported a high prevalence of cardiovascular risk factors as compared to men in terms of obesity, total cholesterol level and also reported low HDL levels both in hypertensive

and normotensive groups. However, in females, the HDL-C remained similar in both hypertensive and normotensive groups. The prevalence of hypertension increased in the men population with increasing BMI due to unhygienic food and poor diet plans. In our multivariate analysis, we observed older age, higher BMI, higher LDL-C levels as independent risk factors of hypertension in the female population (Table 4).

Table 1: Demographic characteristics of recruited patients

Age (years)	Female N= 1936	Male N= 716	Total N= 2652	Weighted prevalance in female population (95% C.I)	Weighted prevalance in male population (95% C.I)
≥80	44 (2.3%)	46 (6.4%)	90 (3.4)	72.7% (57.2-85.0)	78.3% (63.6-89.1)
70-79	199 (10.3%)	166 (23.2%)	365 (13.8)	65.8 % (58.8-72.4)	60.8% (53.0-68.3)
60-69	521 (26.9%)	304 (42.5%)	825 (31.1)	62.8 % (58.5-66.9)	61.2% (55.5-66.7)
50-59	659 (34%)	128 (17.9%)	787 (29.7)	53.6% (49.7-57.4)	61.7% (52.7-70.2)
40-49	513 (26.5%)	72 (10.1%)	585 (22.1)	34.1% (30.0-38.4)	59.7% (47.5-71.7)
Smoking	100 (3.77%)	358 (13.49%)	458 (17.2%)	-	-

Table 2: Systolic and diastolic blood pressure in male and female population

	Male		Female	
	Hypertensive	Normotensive	Normotensive	Hypertensive

	group	group	group	group
Diastolic blood pressure	92 ± 9	75 ± 7	92 ± 9	76 ± 6
Systolic blood pressure	114 ± 18	116 ± 10	114 ± 18	116 ± 11

Table 3: Blood parameters of male and female population associated with risk of cardiovascular disorders.

Blood parameters	Males		Females	
	Hypertensive	Normotensive	Normotensive	Hypertensive
UA (µmol/L)	351 ± 92	317 ± 90	291 ± 70	303 ± 75
TC (mmol/L)	5.45 ± 1.01	5.25 ± 1.01	5.04 ± 0.89	5.15 ± 0.91
CREA (µmol/L)	72.2 ± 52.9	66.6 ± 16.1	64.8 ± 12.7	65.9 ± 16.4
TG (mmol/L)	1.42 ± 0.85	1.24 ± 0.71	1.15 ± 0.72	1.29 ± 0.87
Urea (mmol/L)	6.57 ± 2.13	6.26 ± 1.53	6.04 ± 1.60	6.25 ± 1.78
LDL-C (mmol/L)	2.56 ± 0.70	2.40 ± 0.61	2.23 ± 0.57	2.30 ± 0.63
HDL-C (mmol/L)	1.33 ± 0.33	1.36 ± 0.32	1.38 ± 0.43	1.37 ± 0.38

Table 4: Multivariate Binary Logistic regression of risk factors contributing in hypertension

	Odd ratio	95% CI for OR		Wald	p-value
		Lower	Upper		
Age					
≥80	3.302	2.008	5.429	22.161	<0.001
70-79	1.687	1.255	2.268	12.024	0.001

60-69	1.609	1.269	2.040	15.434	<0.001
50-59	1.333	1.090	1.630	7.850	0.005
40-49	-	-	-	37.480	<0.001
Male Vs Female	1.353	1.111	1.647	9.077	0.003
BMI					
Obesity	3.299	2.609	4.172	99.388	<0.001
Overweight	1.517	1.263	1.821	19.956	<0.001
Underweight/ normal	-	-		99.762	<0.001
Constant	0.052	-	--	103.128	0.001
UA (umol/L)	1.003	1.002	1.004	25.202	0.001
Urea (mmol/L)	1.015	0.969	1.064	0.406	0.524
LDL-C (mmol/L)	1.295	1.142	1.469	16.206	<0.001
TG (mmol/L)	1.104	0.987	1.234	2.993	0.084
Smoking	1.235	0.984	1.551	3.302	0.069

Discussion:

Our study reported recent trends in blood pressure control and highlights the cardiovascular risk factors associated with hypertension. Our study found a slightly high prevalence of hypertension among males but excluded large gender differences. We observed that the male population reported a high prevalence of cardiovascular risk factors as compared to men in terms of obesity, total cholesterol level and also reported low HDL levels. Elevated blood pressure level is one of the major causes of mortality in many regions. A study conducted in 2015 reports 4.5 million deaths in men and 4.0 million deaths in women due to high systolic

blood pressure in low-income regions¹³. A study of Reckelhoff¹⁴ observed high blood pressure among the male population as compared to females but this relationship could be varied by age. A study by Whelton¹⁵ observed higher systolic blood pressure in women when compared with men. Furthermore, they observed women had lower diastolic blood pressure than men. In our study, we did not find any significant gender differences in terms of systolic and diastolic blood pressure. In both genders, we observed similar systolic blood pressure (144 ± 18 vs. 144 ± 18 mmHg) and diastolic blood pressure (92 ± 9 vs. 92 ± 9 mmHg). This similarity occurred due to the sample selection. We only include patients 40 years or above age group for analysis. Our results are similar to the worldwide survey of Zhou et al¹³, in which he reported similar SBP and DBP among the male and female populations aged ≥ 50 years.

Multivariate analysis of Hyman et al¹⁶, reported uncontrolled blood pressure among the male population due to lack of awareness. In our multivariate analysis, we observed older age, higher BMI, higher LDL-C levels as independent risk factors of hypertension in the female population. Sex hormones are the major contributors to blood pressure. Androgens levels could increase the blood pressure level in the male population¹⁴. Our study did not observe the blood pressure control rate in premenopausal and postmenopausal women because 80% of women diagnosed with hypertension were postmenopausal.

Aging is another risk factor for hypertension. Increasing age is highly associated with uncontrolled blood pressure levels¹⁷. After menopause women usually observed elevated systolic blood pressure which is thought to be secondary to the withdrawal of vasodilator effects of endogenous estrogen. This effect also increased arterial stiffness, reduces the production of endothelial nitric oxide, and causes salt sensitivity. Researchers observed increased levels of systolic blood pressure and pulse pressure in postmenopausal women as compared to the elder age male population^{18,19}. But our results are in contradiction of these results and we did not observe any significant difference in systolic blood pressure of both genders. However, the DBP level remains the same in both genders¹⁸. After menopause large population of women are suffering from obesity which is another contributory factor of hypertension-related cardiovascular diseases. The previous study of Framingham²⁰ reported a high prevalence of cardiovascular diseases in the female population after menopause. Menopause increases waist circumference and fat mass. The changes in body composition play a vital role in uncontrolled blood pressure²⁰. Our study reported high obesity levels in the male population due to poor and unhygienic diet plans which may threaten them with cardiovascular diseases in the future.

Conclusion:

Hypertension is one of the leading causes of cardiovascular disorders in many regions of the world. Our results indicated significant sex-related homogeneity in terms of hypertension. The

weighted prevalence of our results shows a high male prevalence of hypertension due to increased levels of androgens levels of men. Interestingly, 80% of our female population was postmenopausal but they had less elevated blood pressure levels due to early diagnosis, and awareness. We recommend that further research would be needed to understand sex-specific outcomes associated with hypertension.

References:

1. Suchard MA Schuemie MJ Krumholz HM et al. Comprehensive comparative effectiveness and safety of first-line antihypertensive drug classes: a systematic, multinational, large-scale analysis. *Lancet*. 2019; 394: 1816-1826
2. Ettehad D Emdin CA Kiran A et al. Blood pressure lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis. *Lancet*. 2016; 387: 957-967
3. Olsen MH Angell SY Asma S et al. A call to action and a lifecourse strategy to address the global burden of raised blood pressure on current and future generations: the Lancet Commission on hypertension. *Lancet*. 2016; 388: 2665-2712
4. Zhang Y Moran AE. Trends in the prevalence, awareness, treatment, and control of hypertension among young adults in the United States, 1999 to 2014. *Hypertension*. 2017; 70: 736-742.
5. NCD Risk Factor Collaboration (NCD-RisC). Long-term and recent trends in hypertension awareness, treatment, and control in 12 high-income countries: an analysis of 123 nationally representative surveys. *Lancet*. 2019; 394: 639-651
6. UN Department of Economic and Social Affairs Population Division World population prospects 2019, online edition. Rev 1. <https://population.un.org/wpp/Download/Standard/CSV> Date: 2019. Date accessed: November 23, 2020

7. Mills KT Bundy JD Kelly TN et al. Global disparities of hypertension prevalence and control: a systematic analysis of population-based studies from 90 countries. *Circulation*. 2016; 134: 441-450
8. Benjamin EJ, Blaha MJ, Chiuve SE, et al.; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart Disease and Stroke Statistics-2017 Update: A Report From the American Heart Association. *Circulation*. 2017; 135:e146–e603.
9. D. A. Calhoun, J. N. Booth, S. Oparil et al. Determination of prevalence, risk factors, and comorbidities in a large, population-based Hypertension. 2014; vol. 63 (3):451–458.
10. C. Chen and F. C. Lu. The guidelines for prevention and control of overweight and obesity in Chinese adults. *Biomedical and Environmental Sciences*. 2004;17: 1–36.
11. W. He, Q. Li, M. Yang et al. Lower BMI cutoffs to define overweight and obesity in China. *Obesity* 2015; 23 (3):684–691.
12. Baiting Liu, Hainiang Liu, Rongmei Na, Xiaofei Li, Qianxiao Li, Libo Chen, Wencheng Tu, Jiahui Hu, Dong Cheng, Yalan Cao, Zhu Li, Weiyi Fang, Ning Zhu, Qin Yu, "A Comparison on Prevalence of Hypertension and Related Risk Factors between Island and Rural Residents of Dalian City, China. *International Journal of Hypertension*. 2019; 19: 8.
13. Zhou B Perel P Mensah GA Ezzati M. Global epidemiology, health burden and effective interventions for elevated blood pressure and hypertension. *Nat Rev Cardiol*. 2021; (published online May 28.)
14. Reckelhoff, J. F. Gender differences in the regulation of blood pressure. *Hypertension*. 2001; 37:1199–1208.
15. He J, Whelton PK. Elevated systolic blood pressure and risk of cardiovascular and renal disease: overview of evidence from observational epidemiologic studies and randomized controlled trials. *Am Heart J*. 1999; 138: 211–219.
16. Hyman DJ, Pavlik VN. Characteristics of patients with uncontrolled hypertension in the United States. *N Engl J Med*. 2001; 345: 479–486.
17. Izumi Y, Matsumoto K, Ozawa Y, Kasamaki Y, Shinndo A, Ohta M, Jumabay M, Nakayama T, Yokoyama E, Shimabukuro H, Kawamura H, Cheng Z, Ma Y, Mahmut M. Effect of age at menopause on blood pressure in postmenopausal women. *Am J Hypertens*. 2007; 20:1045–1050.
18. Lobo RA. Metabolic syndrome after menopause and the role of hormones. *Maturitas*. 2008; 60:10–18.
19. Steiner M, Dunn E, Born L. Hormones and mood: from menarche to menopause and beyond. *J Affect Disord*. 2003; 74:67–83.
20. Kannel WB. The Framingham Study: historical insight on the impact of cardiovascular risk factors in men versus women. *J Gend Specif Med*. 2002; 5: 27–37.

UNDER PEER REVIEW