

Overview on Neurological manifestations of thyroid disease

Abstract:

Hypothyroidism and hyperthyroidism are both prevalent medical conditions that are frequently accompanied with neurologic and neuromuscular dysfunction. Triiodothyronine (T3) and thyroxine (T4) hormones are produced by the thyroid gland and play a crucial role in tissue formation and metabolism. Both of these hormones have a variety of impacts on the brain and neuromuscular system. hypothyroidism can have serious consequences for neuropsychiatric function. The pathophysiological processes underlying the neurological symptoms of hypothyroidism and hyperthyroidism are likely to be multifactorial, in this review we will be looking at multiple neurological as well as psychiatric manifestations related to thyroidal hormones disorders.

Introduction:

Hypothyroidism and hyperthyroidism are both prevalent medical conditions that are frequently accompanied with neurologic and neuromuscular dysfunction. Some of these related conditions are well-known to doctors, but others are not. All of these neurologic symptoms are significant since they might serve as vital indicators to a thyroid disorder's diagnosis. They are also frequently linked to the patients' presenting concerns. Furthermore, they are frequently reversible with the restoration of the patient to a euthyroid state, much like other indications of thyroid disease. [1]

Triiodothyronine (T3) and thyroxine (T4) hormones are produced by the thyroid gland and play a crucial role in tissue formation and metabolism. Both of these hormones have a variety of impacts on the brain and neuromuscular system. Hypothyroidism can generate a variety of neurological indications and symptoms as a result. Muscle weakness, weariness, physical and mental sluggishness, weight gain, constipation, and cold sensitivity are all common symptoms of hypothyroidism. Neuromuscular dysfunctions were shown to be common in thyroid diseases, ranging from 20% to 80%. [2,3,4]

Thyroid hormones have a primary target organ in the brain, and adult-onset hypothyroidism can have serious consequences for neuropsychiatric function. Patients with moderate hypothyroidism, on the other hand, may link unrelated

symptoms to their thyroid status. This can lead to overtreatment or the use of nonstandard thyroid hormone formulations, both of which have hazards associated with them. In this context, according to a recent research from the United Kingdom, L-T4 is being recommended more frequently and for milder forms of hypothyroidism. This indicates that L-T4 is frequently recommended for minor causes, most of which are neuropsychiatric in origin. [5]

Hypothyroidism usually affects both the central and peripheral nervous systems. Patients have reflex loss, proximal muscular weakness, numbness, paresthesia, reduced sensations, and slower muscle contraction and relaxation as a result of their peripheral neuropathy. [2] Thyroid diseases affect the general public as well as hospitalised patients. Thyroid illness can manifest with neurological consequences initially, or it can develop alongside other neurological diseases, especially those with an autoimmune origin. As a result, neurologists frequently meet thyroid illness patients. [6] Polyneuropathy has been linked to hypothyroidism in numerous previous studies. Peripheral polyneuropathy is caused by a deficiency in the nerve cell body, axons, or myelin sheath, and it causes reduced nerve conduction velocities and amplitudes. The sural and median nerves are the most usually impacted nerves, as the distal and sensory nerves are affected first. Carpal tunnel syndrome, caused by entrapment of the median nerve, is the most common cause of peripheral nerve injury in hypothyroidism. [2]

Hypothyroidism is often classified into two types. Overt hypothyroidism is characterised as a high serum TSH level combined with a low free T4 (fT4) level, whereas moderate or "subclinical" hypothyroidism is described as a high TSH along with a normal fT4. Although these symptoms are all part of a larger picture of hypothyroidism, they will be explored separately. Subclinical hypothyroidism is especially important since it is frequent, especially among the elderly with cognitive problems. [5]

Sensory neural hearing loss and ophthalmopathy are two symptoms of cranial nerve involvement in thyroid disease. Patients with hypothyroidism were found to have hearing loss in about 37% of cases. The brainstem auditory evoked potential (BAEP) is a technique for determining the functional integrity of the thalamocortical projections to the primary and association cortex. Even with

asymptomatic hypothyroidism, several investigations have found that central and peripheral conduction latencies are prolonged. [2]

The pathophysiological processes underlying the neurological symptoms of hypothyroidism and hyperthyroidism are likely to be multifactorial, (1) autoimmune mechanisms such as Hashimoto's encephalopathy (thyroid peroxidase, thyroglobulin or thyroid microsomes). Thyroid autoantibodies, associations with autoimmune disorders (thyrotropin receptor antibodies) and thyroid dysfunction associated with myasthenia gravis; (2) "channelopathy" seen in periodic paralysis of thyroid toxicity. (3) Adrenergic hypersensitivity associated with hyperthyroidism; (4) Ischemia in vascular attacks in patients with hyperthyroidism and atrial fibrillation. [7]

Longer delay and lower amplitude of visual evoked potentials have been linked to hypothyroidism. VEPs are potential changes recorded from the scalp by electrodes in response to visual stimuli. It's a noninvasive, straightforward electrophysiological test for determining the impact of hypothyroidism on the central nervous system (CNS). If electrodiagnostic techniques are to be employed in the early diagnosis of nerve involvement in hypothyroidism, it is necessary to understand the anomalies in VEP. [2]

thyroid function has become one of the "regular bloods" in neurology practise. Other symptoms of potential significance in terms of thyroid disease may be overlooked in the busy general neurology clinic, or abnormal thyroid tests may be assumed to be incidental, even though conditions such as carpal tunnel syndrome prompt thyroid testing despite no clear evidence for this approach. [8]

Genetic Disorders of thyroidal metabolism

Allan-Herndon-Dudley Syndrome: The X-linked disorder Allan-Herndon-Dudley syndrome (AHDS) is characterised in males by neurologic observations (hypotonia and feeding difficulties in infancy, developmental delay / intellectual disability varies from minor to profound) and later-onset pyramidal signs, extrapyramidal findings (dystonia, choreoathetosis, paroxysmal movement disorder, hypokinesia, masked facies), and seizures, which are often drug resistant. Dysthyroidism (which manifests as poor weight growth, decreased muscle mass, and fluctuating cold sensitivity, sweating, high heart rate, and irritability) and pathognomonic thyroid test results are other possible findings. [9]

In 1944, William Allan, Nash Herndon, and Florence Dudley published a six-generation North American pedigree with 24 afflicted boys who had a specific mix of dysmorphic traits, intellectual incapacity, and neurological abnormalities. After the discovery of several family male cases with identical clinical characteristics and additional thyroid hormone abnormalities, Allan-Herndon-Dudley syndrome was recognised as a separate X-linked intellectual impairment disease. [7]

Diagnosis/testing: AHDS is diagnosed in a male proband with suggestive signs and a hemizygous SLC16A2 pathogenic variation revealed by molecular genetic testing, and in a female proband with a heterozygous SLC16A2 pathogenic variant detected by molecular genetic testing. [9] Increased blood lactate concentrations, perhaps representing a hyperthyroid metabolic myopathy, and increased serum sex hormone-binding globulin concentrations, likely reflecting a thyroid hormone impact on the liver, are two less specific test abnormalities. In certain people, higher ferritin levels and lower cholesterol levels have been found. [7]

Treatment: Treatment with antiepileptic medications by a neurologist with experience. Thyroid hormone replacement treatment in children offers little benefit and may even be harmful by exacerbating dysthyroidism. Assess developmental progress and educational needs in children every six months until they are four years old, then once a year: neurologic examination for new manifestations (e.g., seizures, changes in tone, movement disorders); spine for scoliosis and hips for dislocation; mobility and self-help skills. L-T4 or L-T3 administration alone can aggravate elevated serum T3 levels and the consequent hypermetabolism. [9]

Benign Hereditary Chorea (Brain-Lung-Thyroid Disease): Benign hereditary chorea (BHC) is an autosomal dominant condition characterised by nonprogressive chorea that develops early in life. BHC is distinguished from Huntington disease by its early start and benign history. [10] NKX2-1-related illnesses include choreoathetosis, congenital hypothyroidism, and infant respiratory distress, among others. The hallmark of NKX2-1-related illnesses, childhood chorea, may or may not be linked to respiratory distress syndrome or congenital hypothyroidism. Chorea usually begins in early infancy or around the age of one year (in most cases), or later in childhood or adolescence, and persists until the second decade, after which it remains static or remits. The second most

prevalent symptom is pulmonary illness, which includes respiratory distress syndrome in newborns, interstitial lung disease in young children, and pulmonary fibrosis in the elderly. generally speaking, BHC individuals have normal IQ, and dementia is not present, as it is in HD patients. Early start of symptoms, usually before the age of one year, indicates a developmental problem. Choreic motions seem to reduce in certain BHC families as they become older. Harper observed an almost full penetrance in men and a 0.75 penetrance in females (1978). The condition is less common than HD. [10]

Laboratory tests may reveal thyroid function problems in BHC patients, such as low thyroxine and excessive thyroid-stimulating hormone. Due to the ease with which hypothyroidism may be treated with l-thyroxine, individuals with early onset chorea, as well as other movement disorders and neurodevelopmental delays, should have thyroid function tests performed as part of their initial diagnostic work-up. Patients with mutation-positive BHC should also have their thyroid function tested on a regular basis. [7]

Treatment: Tetrabenazine and levodopa have been found to be the most effective therapy in a few individual patients with BHC. Some individuals were given these drugs, but only experienced adverse effects and saw no or little improvement in their hyperkinetic movements. As a result, no results or recommendations can be derived from this or past studies. Physical exercise, physiotherapy, interdisciplinary facilitation at school and at work, and yearly lung and thyroid checks and medication, if needed, are probably still the best treatment choices for most people with BHC. [11]

Psychiatric and cognitive effects of hypothyroidism:

Many clinic patients complain of low quality of life and weariness, which are frequently linked to mental co-morbidity and urge a TSH check. If the TSH level is modestly high, the patient and physician presume the symptoms are due to mild hypothyroidism. However, these symptoms frequently do not improve with L-T4 therapy, raising the question of whether they were initially caused by the thyroid abnormalities. Subclinical hypothyroid individuals have been found to have higher rates of depression and anxiety symptoms than the general population. [5,12-16] This is not, however, a general finding. In fact, no changes in sadness or anxiety

were detected between euthyroid and subclinical hypothyroid patients in the biggest investigations. [5,17,18]

Overt hypothyroidism can have an impact on a variety of cognitive functions. General intellect, attention/concentration, memory, perceptual function, language, psychomotor function, and executive function have all been found to be impaired in studies. Memory is the most consistently impacted domain, with verbal memory losses being the most prominent. Although there may not be total reversal, L-T4 therapy is typically beneficial in correcting these decrements. [5]

Neurologic complications of thyroid dysfunction:

In England and Wales, the outpatient "incidence" for thyrotoxicosis was 1.1 per 1,000 in 1955 and 1.7 per 1,000 in 1956. Inpatient "incidence" in the United States in 1971 was 0.16 per 1,000 for thyrotoxicosis and 0.13 per 1,000 for myxedema. In Olmsted County, Minnesota, the average yearly incidence of Graves' illness for females was 30.5 per 100,000 from 1935 to 1967. Clinical evidence of mild or severe muscular weakness can be found in more than half of hyperthyroid individuals. The weakness is usually proximal, and electromyography and muscle biopsy both show the presence of a myopathic process. Severe muscle weakness with an immediate start is rather uncommon, occurring in only around 1% of hyperthyroid individuals. Patients with ophthalmoplegia and psychosis account for 4% and 2% of all cases, respectively. Despite its widespread awareness, myasthenia gravis is thought to affect less than 1% of people. TPP is almost non-existent in the West; nevertheless, it is seen in 2 to 8% of hyperthyroid patients in the Orient, and is 20 to 60 times more common in hyperthyroid men than in hyperthyroid women. Myxedema's neurologic symptomatology is more comprehensive, and there is little agreement across the numerous series. [19]

Neurological manifestations of hyperthyroidism:

Clinical signs of thyrotoxicosis in children and adolescents include tremor, which is commonly noticed in the hands, face, and legs. This result can be noticed in the majority of thyrotoxicosis individuals of any age. Because beta blockers generally give considerable alleviation, tremor is assumed to be due to beta-adrenergic system excitation. It may be viewed at rest as well as with intent. In one research evaluating the prevalence of thyrotoxicosis in children, 58.2 percent of individuals

with the disease developed tremor. Tremor is more prevalent in pubertal/postpubertal children than in preschoolers. Choriform movements associated with thyrotoxicosis have been recorded in children and adolescents, however they are rare. Involuntary but seemingly well-coordinated spastic movements like flexing and extending fingers, raising and lowering shoulders, or grimacing may improve with beta blockade, and in youth totally resolve once thyroid function is normalised with appropriate antithyroid medication or definitive treatment like radioiodine ablation or surgical thyroidectomy. [20-25]

Hashimoto Encephalopathy:

Shaw , who gathered five cases with comparable symptoms such as seizures, disorientation, recurrent episodes of alternating hemiparesis, elevated protein levels in the cerebral fluid, and electrocardiographic abnormalities, coined the name "Hashimoto encephalopathy" (HE) in 1991. These individuals, however, showed hypothyroidism as well as positive thyroid antibodies. A 63-year-old man with seizures, disorientation, recurrent bouts of alternating hemiparesis, elevated protein levels in the cerebral fluid, electrocardiographic abnormalities, and biopsy-confirmed Hashimoto thyroiditis was documented in 1966. [26]

The pathogenesis of Hashimoto encephalopathy, on the other hand, is poorly known, and there is presently no consensus on diagnostic criteria among neurologists and endocrinologists. Although documented cases of the illness improve with thyroid hormone replacement, autopsy investigations in adults reveal lymphocytic infiltration of the central nervous system, results that are comparable to the well-described pathological alterations of thyroiditis found in Hashimoto disease. Furthermore, based on these results, glucocorticoids in anti-inflammatory dosages appear to relieve encephalopathy. [20]

Conclusion:

Thyroid is critical organ due to role of its thyroidal hormones, both hypothyroidism and hyperthyroidism induce some serious neurological and phycological disorders, some of which is genetic, hypothyroidism can impact the development of child and thus regular thyroidal hormones testing is recommended in children who demonstrates any signs of neurological psychiatric or cognitive disease.

References:

1. Swanson JW, Kelly JJ Jr, McConahey WM. Neurologic aspects of thyroid dysfunction. *Mayo Clin Proc.* 1981 Aug;56(8):504-12. PMID: 7266060.
2. Gupta N, Arora M, Sharma R, Arora KS. Peripheral and Central Nervous System Involvement in Recently Diagnosed Cases of Hypothyroidism: An Electrophysiological Study. *Ann Med Health Sci Res.* 2016 Sep-Oct;6(5):261-266. doi: 10.4103/amhsr.amhsr_39_16. PMID: 28503341; PMCID: PMC5414436.
3. Hall JE, Guyton AC. *Textbook of Medical Physiology.* 12th ed. India Elsevier Publications; 2013. pp. 565–76.
4. Parkhad SB, Palve SB, Chandrashekar MC, Shankari P. Assessment and comparison of electrophysiological findings in thyroid dysfunction patients of Kanchipuram District. *Int J Biol Med Res.* 2015;6:5068–71.
5. Samuels MH. Psychiatric and cognitive manifestations of hypothyroidism. *Curr Opin Endocrinol Diabetes Obes.* 2014 Oct;21(5):377-83. doi: 10.1097/MED.000000000000089. PMID: 25122491; PMCID: PMC4264616.
6. Wood-Allum CA, Shaw PJ. Thyroid disease and the nervous system. *Handb Clin Neurol.* 2014;120:703-35. doi: 10.1016/B978-0-7020-4087-0.00048-6. PMID: 24365348.
7. Kurian MA, Jungbluth H. Genetic disorders of thyroid metabolism and brain development. *Dev Med Child Neurol.* 2014 Jul;56(7):627-34. doi: 10.1111/dmcn.12445. Epub 2014 Mar 26. PMID: 24665922; PMCID: PMC4231219.
8. When to consider thyroid dysfunction in the neurology clinic. Mistry N, Wass J, Turner MR *Pract Neurol.* 2009 Jun; 9(3):145-56.
9. Sarret C, Oliver Petit I, Tonduti D. Allan-Herndon-Dudley Syndrome. 2010 Mar 9 [Updated 2020 Jan 16]. In: Adam MP, Ardinger HH, Pagon RA, et al., editors. *GeneReviews*[®] [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2021. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK26373/>

10. de Vries BB, Arts WF, Breedveld GJ, Hoogeboom JJ, Niermeijer MF, Heutink P. Benign hereditary chorea of early onset maps to chromosome 14q. *Am J Hum Genet.* 2000 Jan;66(1):136-42. doi: 10.1086/302725. PMID: 10631144; PMCID: PMC1288318.
11. Koht J, Løstegaard SO, Wedding I, Vidailhet M, Louha M, Tallaksen CM. Benign hereditary chorea, not only chorea: a family case presentation. *Cerebellum Ataxias.* 2016 Feb 2;3:3. doi: 10.1186/s40673-016-0041-7. PMID: 26839702; PMCID: PMC4736661.
12. Roberts LM, Pattison H, Roalfe A, et al. Is subclinical thyroid dysfunction in the elderly associated with depression or cognitive dysfunction? *Ann Intern Med.* 2006;145:573–581.
13. Gussekloo J, van Exel E, de Craen AJ, et al. Thyroid status, disability and cognitive function, and survival in old age. *JAMA.* 2004;292:2591–2599.
14. Bell RJ, Rivera-Woll L, Davison SL, et al. Well-being, health-related quality of life and cardiovascular disease risk profile in women with subclinical thyroid disease - a community-based study. *Clin Endocrinol (Oxf)* 2007;66:548–556.
15. Baldini IM, Vita A, Mauri MC, et al. Psychopathological and cognitive features in subclinical hypothyroidism. *Prog Neuropsychopharmacol Biol Psychiatry.* 1997;21:925–935.
16. Park YJ, Lee EJ, Lee YJ, et al. Subclinical hypothyroidism (SCH) is not associated with metabolic derangement, cognitive impairment, depression or poor quality of life (QoL) in elderly subjects. *Arch Gerontol Geriatr.* 2010;50:68–73.
17. de Jongh RT, Lips P, van Schoor NM, et al. Endogenous subclinical thyroid disorders, physical and cognitive function, depression, and mortality in older individuals. *Eur J Endocrinol.* 2011;165:545–554.
18. Panicker V, Evans J, Bjoro T, et al. A paradoxical difference in relationship between anxiety, depression and thyroid function in subjects on and not on T4: findings from the HUNT study. *Clin Endocrinol.* 2009;71:574–580.
19. Kudrjavcev T. Neurologic complications of thyroid dysfunction. *Adv Neurol.* 1978;19:619-36. PMID: 742545.
20. Nandi-Munshi D, Taplin CE. Thyroid-related neurological disorders and complications in children. *Pediatr Neurol.* 2015 Apr;52(4):373-82. doi: 10.1016/j.pediatrneurol.2014.12.005. Epub 2014 Dec 31. PMID: 25661286.

21. Kung A.W. Neuromuscular complications of thyrotoxicosis. *Clin Endocrinol (Oxf)*. 2007; 67: 645-650.
22. Williamson S. Greene S.A. Incidence of thyrotoxicosis in childhood: A national population based study in the UK and Ireland. *Clin Endocrinol (Oxf)*. 2010; 72: 358-363.
23. Lazar L. Kalter-Leibovici O. Pertzalan A. Weintrob N. Josefsberg Z. Phillip M. Thyrotoxicosis in prepubertal children compared with pubertal and postpubertal patients. *J Clin Endocrinol Metab*. 2000; 85: 3678-3682.
24. Seeherunvong T. Diamantopoulos S. Berkovitz G.D. A nine year old girl with thyrotoxicosis, ataxia, and chorea. *Brain Dev*. 2007; 29: 660-
25. Pozzan G.B. Battistella P.A. Rigon F. et al. Hyperthyroid-induced chorea in an adolescent girl. *Brain Dev*. 1992; 14: 126-127.
26. Payer J, Petrovic T, Lisy L, Langer P. Hashimoto encephalopathy: a rare intricate syndrome. *Int J Endocrinol Metab*. 2012 Spring;10(2):506-14. doi: 10.5812/ijem.4174. Epub 2012 Apr 20. PMID: 23843812; PMCID: PMC3693614.