

## **Atherogenic Plasma Index Levels of Colon Cancer Patients After Chemotherapy**

### **Abstract**

Atherogenic Plasma Index values increase with increasing coronary artery disease risk. There are studies showing that it may increase the risk of cardiovascular disease in cancer, it is thought that cancer and lipoprotein metabolism are closely related. In accordance with the literature, we thought that atherosclerosis may increase with chemotherapy, in which the cardiovascular risk increases in colon cancer patients.

**Keywords:** Atherogenic plasma index, lipoprotein, cancer, colon

### **Introduction**

Despite the newly developed treatment modalities, colorectal cancer is still the 3rd in cancer deaths (1,2). Studies have been carried out in recent years that lipid metabolism may play a role in cancer development and metastasis. A relationship has been found between lipids and disease formation in cancers such as breast prostate lung (3-7). The molecular mechanisms of drug resistance include alteration of the drug-specific binding site, decreased drug permeability that can enzymatically deactivate the drug, and/or gene mutations that can increase the pumping out of drugs across the plasma membrane. Many of these processes are associated with altered lipid metabolism. Lipid metabolism has been found to play a central role in both conventional chemotherapy and resistance to targeted therapies. Atherosclerosis and cancer have been associated with chronic inflammation. Uncontrolled cell proliferation and oxidative stress are common factors in the formation of both diseases. Inflammation plays a central role in the formation of atherosclerosis. Changes in blood lipid values after chemotherapy cause negative effects on chronic inflammation and may contribute to recurrence and metastasis in later periods. Monitoring lipid changes after treatment may contribute to disease control. The atherogenic plasma index (AIP) is an index that consists of triglycerides and high-density lipoprotein cholesterol. It has been used to measure blood lipid levels and is widely used as an optimal indicator of cardiovascular diseases associated with dyslipidemia (8). In our study, we planned to compare the atherogenic plasma index levels of

patients diagnosed with colon cancer before chemotherapy and at the 6th month of chemotherapy. The effect of colon cancer chemotherapy treatment on lipids is unclear. We evaluated the effect on the atherogenic plasma index after chemotherapy and thought that lipid-regulating treatments would contribute to both cancer development and disease control.

## Material Method

The blood lipid profile and hemogram parameters, which were routinely checked in cases over the age of 18 with a histopathological diagnosis of colon cancer, were scanned and recorded until 01.06.2017 -30.06.2021. The lipid parameters, which are routinely checked in the patients who received chemotherapy for the treatment of colon cancer, were recorded before the treatment and at the 6th month of the treatment. The atherogenic plasma index (log (triglyceride/HDL-C)) was calculated before and after the chemotherapy and compared. The study was approved by Afyon University of Health Sciences clinical research ethics committee with the number 2011-KAEK-2.

## Analyzes and Result

**Table 1.** Descriptive statistics for participants

Variable	Groups	F	%
Gender	Male	25	67,6
	Female	12	32,4
	Total	37	100,0

As seen in Table 1, 67.6% of the 37 individuals participating in the research were men and 32.4% were women.

**Table 2.** Descriptive statistics for the participants

individuals	N	M <sub>age</sub>	sd
Weight	37	69,59	11,47
Age	37	61,40	11,16

As seen in Table 2, the average weight of the individuals participating in the research was 69.59±11.47, and the average age of the individuals was 61.40±11.16.

**Table 3. Analysis of the patients' pre/post treatment values**

Measurement	Variables	N	$\bar{x}$	Std. Deviation	Std. Error	t	df	p
Pair 1	B.T. monocyte	37	,5711	,25136	,04132	,914	36	,367

	A.T. monocyte	$\frac{3}{7}$	,5378	,26250	,04315		
Pair 2	B.T. HDL-C	$\frac{3}{7}$	46,4595	14,69050	2,41510	1,778	,84
	A.T. HDL-C	$\frac{3}{7}$	41,1351	12,43463	2,04424		
Pair 3	B.T. LDL-C	$\frac{3}{7}$	109,486 5	36,74735	6,04123	-,660	,51 3
	A.T. LDL-C	$\frac{3}{7}$	113,378 4	31,73620	5,21740		
Pair 4	B.T.Total Cholesterol	$\frac{3}{7}$	186,054 1	39,45106	6,48571	-	,16 5
	A.T. Total Cholesterol	$\frac{3}{7}$	200,351 4	71,67256	11,7829 0	1,416	
Pair 5	B.T. PAİ	$\frac{3}{7}$	,5543	,28587	,04700	-	,17 9
	A.T. PAİ	$\frac{3}{7}$	,6288	,29495	,04849	1,372	
Pair 6	B.T. TG	$\frac{3}{7}$	181,702 7	122,20631	20,0905 9	-,484	,63 1
	A.T. TG	$\frac{3}{7}$	191,648 6	98,99248	16,2742 6		
Pair 7	B.T. Glucose	$\frac{3}{7}$	122,459 5	49,76589	8,18146	-,062	,95 1
	A.T. Glucose	$\frac{3}{7}$	122,945 9	31,16796	5,12398		

BT: before treatment, AT; After treatment, PAİ; Plasma atherogenic index.

As can be seen in Table 3, according to the results of the paired sample t-test analysis performed on the values, between B.T. Monocyte and A.T. Monocyte, between B.T. HDL-C and A.T. HDL-C, between B.T.. LDL-C and A.T. LDL-C, between B.T.Total Cholesterol and A.T.

## Discussion

There was no effect on the atherogenic plasma index measured at the 6th month of our patients who received FOLFOX (5 fluorouracil, folinic acid, oxaliplatin) chemotherapy for the treatment of colon cancer. No significant changes were detected in lipid parameters after chemotherapy. It has been determined in many studies that chemotherapy treatment causes changes in lipid metabolism.

In a study conducted in 18 lymphomas, 18 breast cancers, 14 small cell lung cancers, and 7 urethelial carcinomas, total cholesterol and low-density lipoprotein cholesterol (LDL-C) were increased after chemotherapy in patients other than breast cancer patients. The tendency to increase in serum triglyceride levels was found to be statistically significant in breast cancer patients (9). In a study, a decrease in LDL-C and an increase in high-density lipoprotein cholesterol (HDL-C) were found in patients with less metastatic colon cancer who were given combined chemotherapy with antiangiogenic therapy (10). An increase in triglycerides and a

decrease in HDL-C were detected in bexarotene chemotherapy used for the treatment of cutaneous T-cell lymphoma (11).

It is thought that the change in lipids in colorectal cancer affects the prognosis. In a study, an increase in cholesterol levels and a decrease in triglyceride and HDL-C levels were found in patients with colorectal cancer after adjuvant chemotherapy. In addition, HDL-C elevation was found to be prognostic (12). In our study, no change was detected in the serum lipid profile. Radiotherapy and chemotherapy treatment are side effects of heart failure, hypertension, thromboembolism and atherosclerosis (13). Both radiotherapy and chemotherapy treatment cause inflammation and endothelial cell activation, which leads to the onset of atherosclerosis. It also causes inhibition of thrombolysis. It causes instability in previously formed plaques. Altered expression of thrombolysis-related proteases is involved in atherosclerotic plaque progression and in the process of cancer invasion and metastasis (14).

There are a limited number of studies in the literature examining lipid changes after colorectal cancer chemotherapy. Lipid change and its mechanism after chemotherapy treatment remain unclear. The number of patients in our study is limited. We think that there is a need for more extensive research by increasing the number of patients.

## References

- 1.Siegel RL, Miller KD, Fedewa SA, Ahnen DJ, Meester RGS, Barzi A, et al. Colorectal cancer statistics, 2017. *CA Cancer J Clin.* 2017;67:177–93.
- 2.Thélin C, Sikka S. Epidemiology of colorectal Cancer — incidence, lifetime risk factors statistics and temporal trends. In: Ettarh R, editor. *Screening for colorectal Cancer with colonoscopy.* London: IntechOpen Limited; 2015. p 61-77
- 3.Yamashita Y, Nishiumi S, Kono S, Takao S, Azuma T, Yoshida M. Differences in elongation of very long chain fatty acids and fatty acid metabolism between triple-negative and hormone receptor-positive breast cancer. *BMC Cancer.* 2017;17:589.
- 4.Yang B, Ren X-L, Fu Y-Q, Gao J-L, Li D. Ratio of n-3/n-6 PUFAs and risk of breast cancer: a meta-analysis of 274135 adult females from 11 independent prospective studies. *BMC Cancer.*
- 5.Llorente A, Skotland T, Sylvänne T, Kauhanen D, Róg T, Orłowski A, et al. Molecular lipidomics of exosomes released by PC-3 prostate cancer cells. *Biochim Biophys Acta - Mol Cell Biol Lipids.* 2013;1831:1302–9.
- 6.Skotland T, Ekroos K, Kauhanen D, Simolin H, Seierstad T, Berge V, et al. Molecular lipid species in urinary exosomes as potential prostate cancer biomarkers. *Eur J Cancer.* 2017;70:122–32.
- 7.Xia S-H, Wang J, Kang JX. Decreased n -6/ n -3 fatty acid ratio reduces the invasive potential of human lung cancer cells by downregulation of cell adhesion/invasion-related genes. *Carcinogenesis.* 2005;26:779–84

8. Cai G, Shi G, Xue S, Lu W. The atherogenic index of plasma is a strong and independent predictor for coronary artery disease in the Chinese Han population. *Medicine (Baltimore)*. 2017;96:e8058..
9. Alexopoulos CG, Pournaras S, Vaslamatzis M, Avgerinos A and Raptis S. Changes in serum lipids and lipoproteins in cancer patients during chemotherapy. *Cancer Chemother Pharmacol*. 1992; 30:412-416.
10. Melichar B, Kalabova H, Krcmova L, Kasparova M, Malirova E, Melicharova K, Pecka M, Hyspler R and Solichova D. Serum homocysteine, cholesterol, retinol, alpha-tocopherol, glycosylated hemoglobin and inflammatory response during therapy with bevacizumab, oxaliplatin, 5-fluorouracil and leucovorin. *Anticancer Res*. 2009; 29:4813-4820
11. Basani S and Garg A. Marked lowering of high-density lipoprotein cholesterol levels due to high dose bexarotene therapy. *J Clin Lipidol*. 2015; 9:832-836. *Oncotarget*, Vol. 7, No. 35
12. Y. Wang, Z. Wang, F. Wang, X. Lei, S. Yan, D. Wang<sup>1</sup>, F. Zhang, R. Xu, L. Wang, Y. Li. *Oncotarget* 2016 Aug 30; 7(35): 57290-57300 .Predictive value of chemotherapy-related high-density lipoprotein cholesterol (HDL) elevation in patients with colorectal cancer receiving adjuvant chemotherapy: an exploratory analysis of 851 cases
13. Moslehi JJ. Cardiovascular toxic effects of targeted cancer therapies. *N Engl J Med* 2016; 375:1457–1467. Review of cardiovascular toxicity of chemotherapeutic agents.
14. G. Stoll, M. Bendszus. Inflammation and atherosclerosis: novel insights into plaque formation and destabilization. *Stroke*, 37 (2006), pp. 1923-1932