

# **Effects of Kinesio-Taping in Lateral Epicondylitis: A Narrative Review**

---

## **ABSTRACT**

Lateral epicondylitis is a familiar musculoskeletal disorder that primarily affects the extensors of the wrist. Kinesio-taping (KT) is a newer generation taping technique that aid's the body's healing process whilst supporting and giving stability to the muscles and joints without hindering the range of motion of the body. The objective of this narrative review is to assess the effects of Kinesio-Taping in lateral epicondylitis in pain reduction and enhancing functional ability of the wrist. Three databases were utilised for this narrative review. All studies that encompassed Kinesio-Taping in the treatment of lateral epicondylitis only were included. 6 articles were utilized for this review in which it showed a reduction in pain and, thereby, improving pain pressure threshold. The current review recognises the need for Kinesio-Taping in individuals with lateral epicondylitis. In spite of the encouraging results, more vigorous randomized control trials are to be conducted to pinpoint the mechanism behind Kinesio-Taping.

*Keywords: Lateral epicondylitis; pain reduction; Kinesio-Taping; proprioceptive stimulation; hand grip strength.*

## **1. INTRODUCTION**

Lateral Epicondylitis commonly known as Tennis Elbow or Lateral Elbow Tendinopathy is an agonising elbow condition that is caused by excessive use of wrist extensors. [1] This can be a work-related repetitive strain/ sports specific injury. [2, 3] With an incidence rate of 15.1 per 10000 patients (in the United States of America alone), lateral epicondylitis affects men and women equitably anywhere between 30 to 60 years of age, peaking in the early 5<sup>th</sup> decade. [4, 5]

A renowned chiropractor named Dr. Kenzo Kase, invented Kinesio-taping (KT) in 1976 that mimics the elastic properties of the human skin. The KT mechanism sprung from the speculation that an external unit can assist in the functions of muscles and other tissues. By design, the tape is elastic and thin, stretching up to 40% - 60% of the actual length, making it extremely pliable compared to its ancestors. The tension created in the skin improves the transmission with mechanoreceptors when stretched and increases the motor unit recruitment amid a muscle contraction. Muscle function is improved by this process and facilitates the contraction of inactive muscles. The employment of KT over the wrist extensors could well and truly be another curative approach. [6]

### **1.1 PATHOGENESIS**

An established pathology has not yet been identified but is repeatedly linked with any work-related activities involving repeated movements/ straining of the wrist. Movements such as excessive gripping, wrist extension, and forearm supination lead to microtrauma of extensor tendons (particularly the Extensor carpi radialis brevis). Over time it leads to hyperplasia of wrist extensor tendons, ultimately leading to tendon degeneration and resulting in Lateral Epicondylitis. [5, 7, 8, 9]

### **1.2 CLINICAL PRESENTATION**

The patient will usually present with pain and tenderness at the lateral epicondyle, decreased grip strength, and inability to extend the wrist. It can aggravate along the upper forearm and cascading through the outer part of the forearm. The pain can also be aggravated with resisted dorsiflexion and radial deviation of the wrist and also wrist extension and fingers extension with elbow extension. [10, 11]

### **1.3 DIAGNOSIS**

During physical examination, special tests like Cozen's (sensitivity = 91%, specificity = 80%) and Mill's (sensitivity = 76%, specificity = 85%) tests are used to determine if the patient has lateral epicondylitis. [12, 13, 14, 15] Other imaging diagnosis like X-Ray shows calcification around the lateral epicondyle with MRI and Ultrasound presenting with fluid (hypoechoic) in the extensor carpi radialis brevis tendon origin plus degenerative changes. [16, 17]

### **1.4. KINESIO-TAPING**

KT is generally placed over and around the region to block off further contraction of the muscles. It is hypothesised to decrease pain and inflammation through the improvement of lymphatic and hematological systems without the restriction of the affected part's range of motion. This method lessens pressure and pain that is caused by irritation of the neurosensory receptors by lifting the skin minutely. [18]

The function is refined by allowing muscle support without hindering any movement. This is attained by stimulating a sudden rise in muscle strength through a concentric pull on the

fascia which may then trigger a muscle contraction or improves the muscle alignment that is instrumental in muscle strength increment. The pain is reduced within the first 24 hours and hopefully for up to a week. [18]

This endorses the context that kinesio-taping can be as a therapeutic approach for patients having lateral epicondylitis. The review is aimed at supporting the need for KT in lateral epicondylitis.

## **2. SEARCH METHODOLOGY**

To examine the effects of Kinesio-Taping in Lateral epicondylitis, PubMed, Google Scholar, and ScienceDirect were used as databases from the year 2016 – 2019 using the following keywords: "Lateral epicondylitis", "Kinesiotaping", "Kinesiotaping in lateral epicondylitis" and "effects of kinesio taping in lateral epicondylitis". 6 articles were later finalised based on the selection criteria.

### **2.1 Inclusion criteria:**

- Studies conducted on human population.
- Studies that included Kinesio-Taping as the main intervention.
- Studies that included participants with lateral epicondylitis only.

### **2.2 Exclusion criteria:**

- Non-English articles in full-text format.

## **3. DISCUSSION**

### **3.1 Kinesio-Taping in pain reduction:**

KT exhibited a notable recovery in pain whilst wrist extension with resistance and also improved pain free grip strength which was conducted by Cho YT et al., in individuals with chronic lateral epicondylitis. The results obtained endorsed it for a short-term pain management strategy with its mechanism still unclear. The first hypotheses points to the gate control theory as the non-neuronal cells that act as the signaling pathway is blocked via the stretch of the tape through keratinocytes by stimulating an inhibitory pain mechanism. [19]

The second hypotheses is aimed at reduction of pain through improved blood circulation that was analyzed by Giray et al., where they noted that this theory was important in choosing a placebo taping and also found that KT plus exercises were superior in reducing pain and disability. They also proposed that eccentric training increased tendon strength by stimulation of mechanoreceptors to produce collagen. [13]

### **3.2 Kinesio-Taping in enhancing grip strength:**

Gracias A et al., performed a study to differentiate the effects of pain pressure threshold and grip strength where KT was effectual in increasing grip strength and decreasing pain in people with lateral epicondylitis. This lead to another hypotheses the muscle technique assisted in reducing muscle tension bettering the grip strength and diminishing pain only after subsequent application of KT. [14]

This is also backed by Shaheen H et al., study that compared the effects between ultrasound and KT in which KT was superiorly better in improving the hand grip strength only for a short term effect. [20]

These two studies exemplified the usage of KT for a shorter duration.

### 3.3 Kinesio-Taping in improving the functionality/ disability of the wrist:

Eraslan L et al., conducted a study to prove if KT improves functionality and pain in lateral epicondylitis patients where KT was utilized from insertion to origin of the extensor carpi radialis brevis and the base of fascia correction was put at the point of pain. This provided sensorimotor and proprioceptive feedback mechanisms which are presumed to boost the lymphatic drainage. The rise in number of muscle fibers were seen to improve the functionality of the wrist through proprioceptive stimulus formulated another theory for KT. [21]

Bhambani S et al., study upheld this theory of improved functionality of the wrist as here KT supported the muscles around the elbow during movement that decreased tension around the area, pain and gave a proprioceptive feedback to the patient. [22]

The selected articles were then subjected to the PEDro (Physiotherapy Evidence Database) scale for quality assertion of articles with total scores ranging from 0 – 11 [23]. A mean score of 7 for the 6 articles utilized further indicate that KT can be implemented for patients with lateral epicondylitis [Table 1.].

**Table 1. Summary of utilised articles**

Study	Subjects	Groups	Results	PEDro Scale
Esra Giray et al. (2019) [13]	31 patients were selected and randomized into 3 groups	Group 1 – KT plus exercises, Group 2 – sham taping plus exercises, Group 3- only exercises	KT showed better pain reduction and functionality than sham taping.	10 ~ Excellent
Alicia Gracias and S Shobhalakshmi (2019) [14]	30 individuals were selected and randomized into 2 groups	Control group and experimental group	Improved grip strength and pain reduction.	7~ Good
Sonam Bhambhani et al. (2019) [22]	24 participants were randomly assigned to 2 groups	Group A – KT with conventional physiotherapy, Group B – conventional physiotherapy only	Group A yielded better results in terms of improved functionality and pain reduction.	6 ~ Good

Hamza Shaheen et al. (2019) [20]	20 patients aged between 20 – 50 years were randomly selected and divided into 2 groups	Group A – ultrasound and exercises, Group B – KT and exercises for 12 sessions over a 4 week period; 3 sessions/ week.	Group B showed significant improvement in hand grip strength and is better than ultrasound.	7 ~ Good
Yeng-Tin Cho et al. (2018) [19]	15 participants	Control group and Sham group	Reduction in pain and improved pain-free grip strength.	7~ Good
Leyla Eraslan et al. (2017) [21]	45 patients randomized into 3 groups	Group 1 – icing, TENS & home exercise program, Group 2 – KT with physiotherapy 3 – Extracorporeal shockwave therapy with physiotherapy	Improved pain-free grip strength and wrist functionality.	7~ Good

#### 4. CONCLUSION

Based on this narrative review and mostly with a PEDro score of 7, KT has been proved to reduce the pain intensity by blocking the afferent pathway of the pain gate neurons, improve the pain pressure threshold, hand-grip strength by increasing the tone of the tendons through peripheral feedback regulation and improve the overall functionality of the wrist extensors by acting at the mechanoreceptors and improving the proprioception of the joint. It is hypothesized that the tape when applied from muscle insertion to origin causing pain inhibition through feeble muscle contractions. All these are hypotheses and more in-depth research should be done to find out the exact physiological mechanism. KT could be implemented in the effective management of Lateral Epicondylitis as a short-term intervention.

#### CONSENT

NIL

#### ETHICAL APPROVAL

NIL

#### REFERENCES

1. Hubbard MJ, Hildebrand BA, Battafaran MM, Battafarano DF. Common soft tissue musculoskeletal pain disorders. Prim Care. 2018 Jun; 45(2): 289–303.

2. Van Rijn RM, Huisstede BM, Koes BW, Burdorf A. Associations between work-related factors and specific disorders at the elbow: a systematic literature review. *Rheumatology (Oxford)*. 2009 May; 48(5):528-36.
3. Kafsas Ş, Kafsas ME, Çınarlı FS. The protective role of Kinesio Taping on Lateral epicondyle pain and handgrip strength performance during the tennis tournament of college players. *Journal of Athletic Performance and Nutrition*. 2018 May; 5(1): 32-43.
4. Degen RM, Conti MS, Camp CL, Altchek DW, Dines JS, Werner BC. Epidemiology and Disease Burden of Lateral Epicondylitis in the USA: Analysis of 85, 318 Patients. *HSS Journal*. 2017 Jun; 14(1):9-14.
5. Canale ST, Beaty JH, Azar FM. *Campbell's Operative Orthopaedics*. 13th ed. Philadelphia: Elsevier; 2017. p. 2330.
6. Lemos TV, Pereira KC, Protássio CC, Lucas LB, Matheus JPC. The effect of kinesio taping on handgrip strength. *J Phys Ther Sci*. 2015 Mar; 27(3): 567-570.
7. Dimitrios S. Lateral elbow tendinopathy: evidence of physiotherapy management. *World J Orthop*. 2016 Aug; 7(8): 463-466.
8. Olaussen M, Holmedal Ø, Mdala I, Brage S, Lindbæk M. Corticosteroid or placebo injection combined with deep transverse friction massage, mills manipulation, stretching and eccentric exercise for acute lateral epicondylitis: a randomised, controlled trial. *BMC Musculoskelet Disord*. 2015 May; 16: 122.
9. Keijsers R, De Vos RJ, Kuijjer PPFM, Van Der Bekerom MPJ, Van Der Woude HJ, Eygendaal D. Tennis Elbow. *Sage Journals*. 2018 Sep; 11(5): 384-92.
10. Geoffroy P, Yaffe MF, Rohan I. Diagnosing and treating lateral epicondylitis. *Can Fam Physician*. 1994 Jan; 40: 73-78.
11. Frontera WR, Delisa JA, Gans BM, Walsh NE, Robinson LR. *Delisa's Physical Medicine & Rehabilitation: Principles and Practice*. 5th ed. Vol 1. Philadelphia: Lippincott Williams and Wilkins; 2010. p. 916-919.
12. Giray E, Karali-Bingul D, Akyuz G. The effectiveness of Kinesiotaping, sham taping or exercises only in Lateral Epicondylitis treatment: A Randomized Controlled Study. *American Academy Of Physical Medicine and Rehabilitation*. 2019 Nov; 11: 681-693.
13. Gracias A, Shobhalakshmi S. Immediate effect of Kinesio Taping on pain and grip strength in individuals with Lateral Epicondylitis. *Indian Journal of Physical Therapy and Research*. 2019 Apr; 1(1): 24-28.
14. Dones VC, Grimmer KA, Milanese S, Kumar S. The Sensitivity of the Provocation Tests in replicating pain on the lateral elbow area of participants with lateral epicondylalgia. *Journal of Case Reports and Clinical Research Studies*. 2014; 1(1): 1-15.
15. Karanasios S, Korakakis V, Moutzouri M, Drakonaki E, Koci K, Pantazopolou V, Tsepis E, Gioftsos G. Diagnostic accuracy of examination tests for lateral elbow Tendinopathy (LET) – A systematic review. *J Hand Ther*. 2021 Apr; 4(3): 1-10.
16. Ahmad Z, Siddiqui N, Malik SS, Abdus-Samee M, Tytherleigh-Strong G, Rushton N. Lateral epicondylitis: A Review of pathology and management. *Bone Joint J*. 2013; 95-B: 1158-1164.
17. Faro F, Wolf JM, Lateral Epicondylitis: Review and Current Concepts. *J Hand Surg*. 2007 Oct; 32(8): 1271-1279.

18. Mostafavifar M, Wertz J, Borchers J. A systematic review of the effectiveness of Kinesio Taping for musculoskeletal injury. *Phys Sportsmed* 2012 Nov; 40(4): 33-40.
19. Cho YT, Hsu WY, Lin LF, Lin YN. Kinesio Taping reduces elbow pain during resisted wrist extension in patients with Chronic Lateral Epicondylitis: A Randomized, Double-Blinded, Cross-Over Study. *BMC Musculoskelet Disord*. 2018; 19: 193.
20. Shaheen H, Alarab A, Ahmad MS. Effectiveness of therapeutic ultrasound and kinesio tape in treatment of tennis elbow. *Journal of Novel Physiotherapy and Rehabilitation*. 2019 Apr; 3: 25-33.
21. Eraslan L, Yuce D, Erbilici A, Baltaci G. Does Kinesiotaping improve pain and functionality in patients with newly diagnosed Lateral Epicondylitis?. *European Society Of Sports Traumatology, Knee Surgery And Arthroscopy*. 2017 Aug; 26(3): 938-945.
22. Bhambhani S, Mitra M, Kaur A. Effectiveness of Kinesiotaping along with conventional physiotherapy for patients with tennis elbow. *Indian Journal of Physiotherapy and Occupational Therapy*. 2016 Sep; 10(3): 18-22.
23. Maher CG, Sherrington C, Herbert RD, Moseley AM, Elkins M. Reliability of the PEDro scale for rating quality of randomised controlled trials. *Phys Ther*. 2003; 83(8):713-21.