

Original Research Article

ESTIMATION OF BLOOD PRESSURE ON POSTPRANDIAL BREAKFAST, MEAL AND DINNER AMONG VEGETARIANS

Running title: Measuring BP among vegetarians

ABSTRACT:

BACKGROUND: The blood pressure is the pressure of the blood within the arteries. Based on the circadian rhythm the BP can change. DBP can change according to age. Processed foods, canned soups, snack foods, and condiments may increase BP. The main aim of this study is to estimate the difference in postprandial BP responses during breakfast, lunch and dinner.

MATERIALS AND METHODS: A total of 26 healthy volunteers were randomly selected, men and women aged 17-20 for the study. Systolic blood pressure, Diastolic blood pressure, Heart rate were measured with ambulatory Blood Pressure devices every 10 min before until 90 min after each meal. The measurements were taken to identify the difference in postprandial Blood pressure responses between breakfast, lunch and dinner. Before starting the study, informed consent was taken. Data were collected and analysed using SPSS software, independent sample t test.

RESULTS: In the present study, there was a mild increase observed before and after breakfast, lunch and dinner. There is a difference in BP before and postprandial during breakfast, lunch and dinner. In the present study, there was a mild increase observed before and after breakfast, lunch and dinner. In the morning the body releases hormones such as adrenaline and noradrenaline. These hormones boost energy which raises the BP. So there is decrease in BP before and postprandial during breakfast

CONCLUSION: From this study only a significant difference was observed in SBP, DBP, HR of dinner. There was no significant change seen in breakfast.

KEY WORDS: Blood pressure, Dietary patterns, Healthy diet, Hypertension, DASH diet, Innovative technology

INTRODUCTION:

Blood Pressure is the pressure of blood against blood vessels. Two blood pressure: Systolic blood pressure (SBP) and Diastolic blood pressure (DBP). Optimal systolic blood pressure is 120 mmHg and optimal diastolic blood pressure is 80 mmHg. SBP measures the

pressure that blood exerts on vessels while the heart is beating. DBP measures the pressure in the vessels between heartbeats. Normal blood pressure level is less than 120/80 mmHg.(1). Normally the blood pressure is lower at night while sleeping. But, a few hours before I wake up, my blood pressure starts to rise. Usually, BP will be peaking in the middle of the afternoon and in the evening BP drops again (2).

People with blood pressure will not experience any symptoms and hypertension also known as silent killer. When blood pressure reaches nearly 180/120 mmHg it is a hypertensive crisis. Getting less oxygen will damage blood walls and make it harder to regulate the blood pressure(3,4). Decreased secretion of thyroid hormones can slow the heart rate (5). High blood pressure is associated with high cardiovascular risk. In elder people, the most powerful predictor of risk is due to the increase in pulse pressure which causes decreased diastolic and increased systolic pressure. An increase in BP is an inevitable consequence of ageing(6). Temporary decrease in blood pressure can compensate by constricting the blood vessels present in the digestive tract (outside) which causes the heart to beat faster (7,8).

Water drinking increases systolic blood pressure by 100 mmHg in some patients. In such patients water should be avoided for 1.5 hours after retrieving. Citrus fruits and grapefruit may have powerful effects of lowering the blood pressure (9). When a person has their food, their body directs blood to the digestive tract to help digestion. If someone has high blood pressure, a DASH diet or Mediterranean diet it can lower it. Blood pressure can be raised by salt and anger.

Diet is an important factor, which can prevent hypertension. Vegetarian diet can lower the blood pressure. Vegetarian dietary patterns significantly reduce systolic and diastolic blood pressure. Vegetarians consume foods like vegetables, grains, legumes, fruits and dairy products. Some people rarely include eggs. Vegetarian people consume less protein foods. As vegetarians can consume protein rich vegetarian foods. As a vegetarian you cannot have meat, fish which are rich in protein content compared to vegetables and grains (10).

Our team has extensive knowledge and research experience that has translate into high quality publications (11–13)(4,14–18),(19)(20),(21)(22),(23)(24)(8,25–28) The main aim of the study is to estimate the difference in postprandial BP responses during breakfast, lunch and dinner.

MATERIALS AND METHODS:

A total of 26 healthy volunteers were randomly selected, men and women aged 17-20 for the study. Systolic blood pressure, Diastolic blood pressure, Heart rate were measured with ambulatory Blood Pressure devices every 10 min before until 90 min after each meal. The measurements were taken to identify the difference in postprandial Blood pressure responses between breakfast, lunch and dinner. All gave their written informed consent before beginning the trial. The study was approved by the institutional ethics committee and was conducted in accordance with the guidelines. Before starting the study, informed consent was taken. Data were collected and analysed using an independent sample t test.

RESULTS:

In the present study, there was a mild increase observed before and after breakfast, lunch and dinner (Table 1). There is a difference in BP before and postprandial during breakfast, lunch and dinner. There was a statistically significant difference found before and after dinner of all parameters. There was a statistically significant difference found before and after meals of DBP and HR. There was a statistically insignificant difference found before and after breakfast (Table 2). In the present study, there was a mild increase observed before and after breakfast, lunch and dinner (Figure 1, 2, 3). In the morning the body releases hormones such as adrenaline and noradrenaline. These hormones boost energy which raises the BP. So there is decrease in BP before and postprandial during breakfast




Figure 1: The bar graph represents the breakfast before and after. Blue represents the before and red represents after breakfast. There was no significant difference found in SBP, DBP, HR after breakfast.




Figure 2: The bar graph represents the lunch before and after. Blue represents the before meal

and red represents after meal.* represent statistically significant (p<0.05). There was a significant difference found in DBP, HR after meals.




Figure 3:The graph represents the before and after dinner. Blue colour represents the before dinner and red colour represents after dinner. * represent statistically significant (p<0.05). There was a significant difference found in SBP, DBP, HR after dinner.

Table 1: Mean and SD of before and after breakfast, lunch and dinner

	Breakfast		Lunch		Dinner	
	BEFORE	AFTER	BEFORE	AFTER	BEFORE	AFTER
SBP	100.23 ± 11.98	106.84 ± 12.08	107.21 ± 12.78	114.73 ± 12.53	98.52 ± 6.80	113.05 ± 15.11
DBP	70.25 ± 6.51	73.15 ± 7.01	65.36 ± 9.77	73.15 ± 9.61	66.21 ± 7.80	76.20 ± 8.97
HR	69.25 ± 8.54	73.26 ± 9.64	79.89 ± 13.59	93.47 ± 13.77	79.42 ± 13.99	94.42 ± 15.61

Table 2: Significance level of independent sample t test

Parameters	Breakfast	Lunch	Dinner
SBP	0.09	0.07	0.00
DBP	0.392	0.008	0.00
HR	0.183	0.004	0.00

DISCUSSION:

The main finding of this study is the relevant variation of before and postprandial BP response during the day among young adolescents. There is no significance in SBP, DBP, HR in breakfast and DBP, HR is significant in lunch, SBP, DBP, HR is significant in dinner.

Previous studies conducted among vegetarian meals and people consumed a comparable meal at both breakfast and lunch, dinner and had a standardized period of fasting before both the lunchtime and dinner time tests (29). There are some differences in before and postprandial BP in breakfast, lunch and dinner (30). But there is no significant difference found in SBP, DBP, HR of the breakfast and no significant difference in SBP of lunch and there is significant difference in DBP, HR of the lunch and also there is significant difference in SBP, DBP, HR of the dinner (31,32). However, results suggest that central hemodynamic measures are more sensitive to the feeding state than are standard BP measurements.

Many studies are done in postprandial time and at different meal times. This is the first study put forward to study the SBP, DBP, HR of breakfast, lunch and dinner.(33,34). Clinical BP measurements for the diagnosis and monitoring of hypertension should be performed in a standardized manner with respect to meals (35). This Random BP measurement may be appropriate for primary care, but the operator should be aware of the fasting and inform the patient the meaning of the result recorded. There is a controversial study to the present study that BP has increased from breakfast to lunch.

The limitations of this study was done on a limited population in a shorter duration. The study cannot compare vegetarian and non vegetarian individuals. It also cannot check the carbohydrate, fats and protein intake.

CONCLUSION:

From this study, only a significant difference was observed in SBP, DBP, HR of dinner. There was no significant change seen in breakfast. Most vegetarians have lower blood pressure than non-vegetarians.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of

knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES:

1. Dempsey E, Seri I. Definition of Normal Blood Pressure Range [Internet]. *Hemodynamics and Cardiology*. 2019. p. 47–64. Available from: <http://dx.doi.org/10.1016/b978-0-323-53366-9.00003-x>
2. Sheps SG. *Mayo Clinic on High Blood Pressure*. Orient Paperbacks; 2005. 192 p.
3. Egbuna C, Mishra AP, Goyal MR. *Preparation of Phytopharmaceuticals for the Management of Disorders: The Development of Nutraceuticals and Traditional Medicine*. Academic Press; 2020. 570 p.
4. Wadhwa R, Paudel KR, Chin LH, Hon CM, Madheswaran T, Gupta G, et al. Anti-inflammatory and anticancer activities of Naringenin-loaded liquid crystalline nanoparticles in vitro. *J Food Biochem*. 2021 Jan;45(1):e13572.
5. Oparil S, Acelajado MC, Bakris GL, Berlowitz DR, Cífková R, Dominiczak AF, et al. Hypertension. *Nat Rev Dis Primers*. 2018 Mar 22;4:18014.
6. Master AM. THE NORMAL BLOOD PRESSURE RANGE AND ITS CLINICAL IMPLICATIONS [Internet]. Vol. 143, *Journal of the American Medical Association*. 1950. p. 1464. Available from: <http://dx.doi.org/10.1001/jama.1950.02910520006004>
7. Acelajado MC, Oparil S. Hypertension and cardiovascular risk [Internet]. *Hypertension*. 2013. p. 182–94. Available from: <http://dx.doi.org/10.2217/ebo.12.367>
8. Wahab PUA, Madhulaxmi M, Senthilnathan P, Muthusekhar MR, Vohra Y, Abhinav RP. Scalpel Versus Diathermy in Wound Healing After Mucosal Incisions: A Split-Mouth Study. *J Oral Maxillofac Surg*. 2018 Jun;76(6):1160–4.
9. Blood Pressure Diary [Internet]. Vol. 10, *Blood Pressure*. 2001. p. 55–55. Available from: <http://dx.doi.org/10.1080/080370501750183417>
10. Yokoyama Y, Tsubota K, Watanabe M. Effects of vegetarian diets on blood pressure [Internet]. *Nutrition and Dietary Supplements*. 2016. p. 57. Available from: <http://dx.doi.org/10.2147/nds.s74715>
11. Saraswathi I, Saikarthik J, Senthil Kumar K, Madhan Srinivasan K, Ardhanaari M, Gunapriya R. Impact of COVID-19 outbreak on the mental health status of undergraduate medical students in a COVID-19 treating medical college: a prospective longitudinal study. *PeerJ*. 2020 Oct 16;8:e10164.
12. Santhakumar P, Roy A, Mohanraj KG, Jayaraman S, Durairaj R. Ethanolic Extract of *Capparis decidua* Fruit Ameliorates Methotrexate-Induced Hepatotoxicity by Activating

- Nrf2/HO-1 and PPAR γ Mediated Pathways. *Ind J Pharm Educ*. 2021 Mar 19;55(1s):s265–74.
13. Nambi G, Kamal W, Es S, Joshi S, Trivedi P. Spinal manipulation plus laser therapy versus laser therapy alone in the treatment of chronic non-specific low back pain: a randomized controlled study. *Eur J Phys Rehabil Med*. 2018 Dec;54(6):880–9.
 14. Rajakumari R, Volova T, Oluwafemi OS, Rajesh Kumar S, Thomas S, Kalarikkal N. Grape seed extract-soluplus dispersion and its antioxidant activity. *Drug Dev Ind Pharm*. 2020 Aug;46(8):1219–29.
 15. Clarizia G, Bernardo P. Diverse Applications of Organic-Inorganic Nanocomposites: Emerging Research and Opportunities: Emerging Research and Opportunities. IGI Global; 2019. 237 p.
 16. Solai Prakash AK, Devaraj E. Cytotoxic potentials of *S. cumini* methanolic seed kernel extract in human hepatoma HepG2 cells. *Environ Toxicol*. 2019 Dec;34(12):1313–9.
 17. Tahmasebi S, Qasim MT, Krivenkova MV, Zekiy AO, Thangavelu L, Aravindhan S, et al. The effects of oxygen-ozone therapy on regulatory T-cell responses in multiple sclerosis patients. *Cell Biol Int*. 2021 Jul;45(7):1498–509.
 18. Vivekanandhan K, Shanmugam P, Barabadi H, Arumugam V, Daniel Raj Daniel Paul Raj D, Sivasubramanian M, et al. Emerging Therapeutic Approaches to Combat COVID-19: Present Status and Future Perspectives. *Front Mol Biosci*. 2021 Mar 8;8:604447.
 19. Ezhilarasan D. Critical role of estrogen in the progression of chronic liver diseases. *Hepatobiliary Pancreat Dis Int*. 2020 Oct;19(5):429–34.
 20. Egbuna C, Mishra AP, Goyal MR. Preparation of Phytopharmaceuticals for the Management of Disorders: The Development of Nutraceuticals and Traditional Medicine. Academic Press; 2020. 574 p.
 21. Kamath SM, Manjunath Kamath S, Jaison D, Rao SK, Sridhar K, Kasthuri N, et al. In vitro augmentation of chondrogenesis by Epigallocatechin gallate in primary Human chondrocytes - Sustained release model for cartilage regeneration [Internet]. Vol. 60, *Journal of Drug Delivery Science and Technology*. 2020. p. 101992. Available from: <http://dx.doi.org/10.1016/j.jddst.2020.101992>
 22. Barabadi H, Mojab F, Vahidi H, Marashi B, Talank N, Hosseini O, et al. Green synthesis, characterization, antibacterial and biofilm inhibitory activity of silver nanoparticles compared to commercial silver nanoparticles [Internet]. Vol. 129, *Inorganic Chemistry Communications*. 2021. p. 108647. Available from: <http://dx.doi.org/10.1016/j.inoche.2021.108647>
 23. Bharath B, Perinbam K, Devanesan S, AlSalhi MS, Saravanan M. Evaluation of the anticancer potential of Hexadecanoic acid from brown algae *Turbinaria ornata* on HT–29 colon cancer cells [Internet]. Vol. 1235, *Journal of Molecular Structure*. 2021. p. 130229.

Available from: <http://dx.doi.org/10.1016/j.molstruc.2021.130229>

24. Gowhari Shabgah A, Ezzatifar F, Aravindhan S, Olegovna Zekiy A, Ahmadi M, Gheibihayat SM, et al. Shedding more light on the role of Midkine in hepatocellular carcinoma: New perspectives on diagnosis and therapy. *IUBMB Life*. 2021 Apr;73(4):659–69.
25. Sridharan G, Ramani P, Patankar S, Vijayaraghavan R. Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma. *J Oral Pathol Med*. 2019 Apr;48(4):299–306.
26. R H, Hannah R, Ramani P, Ramanathan A, Jancy MR, Gheena S, et al. CYP2 C9 polymorphism among patients with oral squamous cell carcinoma and its role in altering the metabolism of benzo[a]pyrene [Internet]. Vol. 130, *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology*. 2020. p. 306–12. Available from: <http://dx.doi.org/10.1016/j.oooo.2020.06.021>
27. J PC, Marimuthu T, C K, Devadoss P, Kumar SM. Prevalence and measurement of anterior loop of the mandibular canal using CBCT: A cross sectional study. *Clin Implant Dent Relat Res*. 2018 Aug;20(4):531–4.
28. Mudigonda SK, Murugan S, Velavan K, Thulasiraman S, Krishna Kumar Raja VB. Non-suturing microvascular anastomosis in maxillofacial reconstruction- a comparative study. *J Craniomaxillofac Surg*. 2020 Jun;48(6):599–606.
29. Nambi G, Kamal W, Es S, Joshi S, Trivedi P. Spinal manipulation plus laser therapy versus laser therapy alone in the treatment of chronic non-specific low back pain: a randomized controlled study. *Eur J Phys Rehabil Med*. 2018 Dec;54(6):880–9.
30. National Academies of Sciences, Engineering, and Medicine, Health and Medicine Division, Board on Population Health and Public Health Practice, Committee on Community-Based Solutions to Promote Health Equity in the United States. *Communities in Action: Pathways to Health Equity*. National Academies Press; 2017. 582 p.
31. Vloet LCM, Mehagnoul-Schipper DJ, Hoefnagels WHL, R W M. The Influence of Low-, Normal-, and High-Carbohydrate Meals on Blood Pressure in Elderly Patients With Postprandial Hypotension [Internet]. Vol. 56, *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2001. p. M744–8. Available from: <http://dx.doi.org/10.1093/gerona/56.12.m744>
32. Santhakumar P, Roy A, Mohanraj KG, Jayaraman S, Durairaj R. Ethanolic Extract of *Capparis decidua* Fruit Ameliorates Methotrexate-Induced Hepatotoxicity by Activating Nrf2/HO-1 and PPAR γ Mediated Pathways [Internet]. Vol. 55, *Indian Journal of Pharmaceutical Education and Research*. 2021. p. s265–74. Available from: <http://dx.doi.org/10.5530/ijper.55.1s.59>
33. Vloet LCM, Smits R, Jansen RWMM. The effect of meals at different mealtimes on blood pressure and symptoms in geriatric patients with postprandial hypotension. *J Gerontol A*

Biol Sci Med Sci. 2003 Nov;58(11):1031–5.

34. Fisher AA, Davis MW, Le Couteur DG. THE EFFECT OF MEALS AT DIFFERENT MEALTIMES ON BLOOD PRESSURE AND SYMPTOMS IN GERIATRIC PATIENTS WITH POSTPRANDIAL HYPOTENSION [Internet]. Vol. 60, The Journals of Gerontology Series A: Biological Sciences and Medical Sciences. 2005. p. 184–5. Available from: <http://dx.doi.org/10.1093/gerona/60.2.184>
35. Sridharan G, Ramani P, Patankar S, Vijayaraghavan R. Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma [Internet]. Vol. 48, Journal of Oral Pathology & Medicine. 2019. p. 299–306. Available from: <http://dx.doi.org/10.1111/jop.12835>

UNDER PEER REVIEW