

Antioxidant studies of thiazole ring bearing chalcone derivatives

Abstract

Oxidative stress is one of the common problems seen in a variety of diseases. Chalcones and in particular heteroaryl chalcones had reported with promising antioxidant activities. Hence, in the present work we reported the antioxidant activity of twenty thiazole ring bearing chalcone derivatives (**1-20**). Among the tested compounds, three compounds including 17, 19 and 20 containing 2-pyridinyl, 3-pyridinyl and 2-thiazolyl scaffolds showed superior antioxidant activity than the standard with IC₅₀ values 4±1µg/mL, 3±1 µg/mL and 5±1µg/mL respectively. The compound 19 is an interesting lead for the development of newer antioxidant agents.

Key Words: Oxidative stress; Chalcones; Heteroaryl chalcones; Antioxidant activity.

1. Introduction


Chalcones are a type of natural products containing two aryl rings connected through a three-carbon linkage i.e., ketovinyl group. Chalcones were found to possess broad spectrum of pharmacological activities and many authors have reviewed the same [1-4]. The activity of chalcones is attributed mainly due to the ketovinyl chain and the level or the intensity of the activity is due to the nature of aryl rings [5]. Medicinal chemists had manipulated the basic chalcone core by replacing the phenyl ring with a variety of substituted aryl or heteroaryl rings. It was observed that the presence of heteroaryl ring had improved the intensity of bioactivity over the trivial phenyl or the substituted phenyl rings [6-7]. This improved activity may be due to the excellent biological properties of the heteroaryl scaffolds including good pharmacokinetics, metabolic stability and effective binding with the target receptors [8]. Due to these properties scientists had prepared and reported a large number of heteroaryl chalcones with potential anticancer, antimicrobial, antitubercular activities [9-16]. Among the other bioactivities, the antioxidant activity of small molecules had been well studied as oxidative stress play a vital role in the infectious diseases, cardiovascular (CVS) and neurological disorders, infectious diseases and different forms of cancers. Chalcones were reported with significant antioxidant activity [17-18] and some of their biological profiles are due to their antioxidant effect[19]. Many reports had clearly revealed that the heteroaryl

chalcones comprise excellent antioxidant potential and these compounds are good starting points for the successful development of new drug candidates [20-25]. Chalcones display antioxidant action either by scavenging the highly reactive free radicals or the reactive oxygen species (ROS) or by antagonizing the enzyme aldose reductase [26]. By considering the aforementioned facts we here report the antioxidant activity of 20 previously synthesized chalcones bearing thiazole scaffold [27].

2. Materials and methods

2.1. Chemistry

Method for the synthesis of target thiazole ring bearing chalcone derivatives (1-20): The preparation and characterization of the target compounds was reported [27] wherein 1 mmol of the heteroaryl aldehyde i.e., 2,4-dichloro-5-carboxaldehyde was dissolved in glacial acetic acid and hydrochloric acid and to this mixture aryl/heteroaryl ketone (1 mmol) previously dissolved in ethanol was added and refluxed for a period of 4-6 h. At the end of the reaction, the precipitate that is formed was filtered and washed with 100 mL of cold water and dried. Further, the dry precipitate was purified by column chromatography to isolate the pure compounds (Scheme 1).



R

1. 2-chlorophenyl; 2. 3-chlorophenyl; 3. 4-chlorophenyl; 4. 2,3-dichlorophenyl; 5. 2,6-dichlorophenyl;
6. 2,5-dichlorophenyl; 7. 2,4-dichlorophenyl; 8. 3,4-dichlorophenyl; 9. 2-fluorophenyl; 10. 3-fluorophenyl;
11. 4-fluorophenyl; 12. 2,4-difluorophenyl; 13. 2,5-difluorophenyl; 14. 2,6-difluorophenyl; 15. 3,4-difluorophenyl;
16. 3,5-difluorophenyl; 17. 2-pyridinyl; 18. 3-pyridinyl; 19. 4-pyridinyl; 20. 2-thiazolyl

Scheme 1: Synthesis of target thiazole bearing chalcone derivatives (**1-20**)

2.2. *In vitro* Antioxidant activity

The DPPH assay is a rapid and simple spectrophotometric method for assessing antioxidants, and it may be used to analyse a large number of samples at once. The goal of this study was to use the DPPH free radical assay to assess the antioxidant activity (AA percent) of the target chalcones (**1-20**). The activity of DPPH radical scavenging was measured using the Brand-Williams et al. procedure [28]. In ethanol, the samples were treated with the stable

DPPH radical. A 0.1 mM solution of DPPH was prepared by dissolving it in methanol. The samples were treated with the stable DPPH radical in an ethanol solution. Methanol was used to create varied concentrations of test samples (5–100 g/mL) and standard (1.0, 2.5, and 5.0 g/mL). One millilitre of 0.1 mM DPPH solution was added separately to 3 mL of each concentration of test samples and standard. The absorbance of these combinations was measured at 517 nm after being kept in the dark for about 30 minutes [29]. The following formula was used to calculate the ability to scavenge the DPPH radical:

$$AA\% = 100 - \left[\frac{(\text{Absorbance of the sample} - \text{Absorbance of Control})}{\text{Absorbance of Control}} \right] \times 100$$

After 100 minutes of DPPH reaction, a colour shift from deep violet to light yellow was measured using a UV-VIS spectrophotometer at 517 nm after reducing DPPH with antioxidant molecules.

3. Results and discussion

3.1. Chemistry


Thiazole bearing chalcone derivatives (1-20) were obtained in 75-91% yields and are characterized by spectral methods (Reference). In their FT-IR spectrum the compounds displayed absorption bands for C=O and HC=CH around the wave numbers 1651-1698 cm⁻¹ and 1506-1520 cm⁻¹ respectively whereas the ¹H NMR revealed two diagnostic doublet peaks corresponding to the vinyl protons at chemical shift values 7.27-7.89 ppm and 7.66-8.16 ppm. For these doublets, the coupling constant value J was between 15 and 17 Hz indicating the trans geometry of the chalcone linkage. The synthesis of chalcones was further verified by the molecular ion peak in the mass spectrum. Furthermore, all of the compounds had an isotopic M+2 peak that was one-third intense as the molecular ion peak.

2.2. *In vitro* Antioxidant activity

The antioxidant activity of the target analogues was performed by DPPH assay using gallic acid as a standard and the results are summarized in Table 1. The studied compounds had shown an interesting antioxidant activity with IC₅₀ values ranging between 3-61 µg/mL. These values indicate that there is a difference in the level of the antioxidant activity exhibited by the compounds. For instance, it was observed that compounds **17** and

19 containing heteroaryl 2-pyridinyl and 4-pyridinyl rings were more active than the standard gallic acid ($IC_{50} = 5 \mu\text{g/mL}$) with IC_{50} values $4 \mu\text{g/mL}$ and $3 \mu\text{g/mL}$ respectively whereas compound **20** containing 2-thiazolyl motif was equipotent with gallic acid. The compound **18** bearing 3-pyridinyl ring was close in its activity to the standard with an IC_{50} value of $8 \mu\text{g/mL}$. The other compounds containing halogenated phenyl ring shown optimum antioxidant activity but is not equal or more than gallic acid. From the above results the structure activity relationships suggests that thiazole containing chalcone is optimal for activity and the nature of aryl or heteroaryl ring connected to the chalcone bridge is crucial in determining the antioxidant potency. The results indicate that the thiazole bearing chalcones containing heteroaromatic rings like pyridinyl or thiazolyl scaffolds are much useful for the antioxidant activity over the phenyl substituted rings.

Table 1. Antioxidant activities of thiazole ring bearing chalcone derivatives (1-20). ($IC_{50} \pm SD, \mu\text{g/mL}$)^a



Entry	R	Antioxidant activity IC_{50} ($\mu\text{g/mL}$)
1	2-chlorophenyl	32 ± 1
2	3-chlorophenyl	46 ± 1
3	4-chlorophenyl	23 ± 2
4	2,3-dichlorophenyl	39 ± 1
5	2,6-dichlorophenyl	28 ± 2
6	2,5-dichlorophenyl	33 ± 2
7	2,4-dichlorophenyl	18 ± 1
8	3,4-dichlorophenyl	30 ± 2
9	2-fluorophenyl	36 ± 1
10	3-fluorophenyl	61 ± 2
11	4-fluorophenyl	12 ± 1
12	2,4-difluorophenyl	14 ± 1
13	2,5-difluorophenyl	31 ± 2
14	2,6-difluorophenyl	35 ± 1
15	3,4-difluorophenyl	28 ± 1
16	3,5-difluorophenyl	49 ± 2
17	2-pyridinyl	4 ± 1
18	3-pyridinyl	8 ± 1
19	4-pyridinyl	3 ± 1
20	2-thiazolyl	5 ± 1
Gallic acid	-	5 ± 1

^a IC_{50} are the mean values of three independent experiments

Conclusion

In the present study we have assessed the antioxidant potential of 20 of the previously synthesized thiazole bearing chalcone derivatives containing different kinds of substituted aryl and heteroaryl rings. The compounds 17, 19 and 20 bearing pyridinyl and thiazolyl scaffolds elicited promising antioxidant activity that is more than the standard gallic acid. The most active compounds that have been identified by this study will be useful for the development of prospective antioxidant agents that will be of greater use in the treatment of different disorders.

References

1. Salehi B, Quispe C, Chamkhi I, El Omari N, Balahbib A, Sharifi-Rad J, Bouyahya A, Akram M, Iqbal M, Docea AO, Caruntu C. Pharmacological Properties of Chalcones: A Review of Preclinical Including Molecular Mechanisms and Clinical Evidence. *Frontiers in Pharmacology*. 2020;11.
2. Yazdan SK, Sagar GV, Shaik AB. Biological and synthetic potentiality of Chalcones. *Journal of Chemical and Pharmaceutical Research*. 2015;7(11):829-42.
3. Gaonkar SL, Vignesh UN. Synthesis and pharmacological properties of chalcones: a review. *Research on chemical intermediates*. 2017 Nov;43(11):6043-77.
4. Matos MJ, Vazquez-Rodriguez S, Uriarte E, Santana L. Potential pharmacological uses of chalcones: a patent review (from June 2011–2014). *Expert opinion on therapeutic patents*. 2015 Mar 4;25(3):351-66.
5. Ammaji S, Masthanamma S, Richie RB, Sivakumar A, Afzal BS. Antitubercular and antioxidant activities of hydroxy and chloro substituted chalcone analogues: Synthesis, biological and computational studies. *Arabian Journal of Chemistry*. 2022;15(2):103581.
6. Jaiswal P, Pathak DP, Bansal H, Agarwal U. Chalcone and their heterocyclic analogue: A review article. *J. Chem. Pharm. Res*. 2018;10:160-73.
7. Shaik A, Bhandare RR, Palleapati K, Nissankararao S, Kancharlapalli V, Shaik S. Antimicrobial, antioxidant, and anticancer activities of some novel isoxazole ring containing chalcone and dihydropyrazole derivatives. *Molecules*. 2020 Jan;25(5):1047.
8. Gomtsyan A. Heterocycles in drugs and drug discovery. *Chemistry of heterocyclic compounds*. 2012 Apr 1;48(1):7-10.

9. Farghaly TA, Masaret GS, Muhammad ZA, Harras MF. Discovery of thiazole-based-chalcones and 4-hetarylthiazoles as potent anticancer agents: Synthesis, docking study and anticancer activity. *Bioorganic chemistry*. 2020 May 1;98:103761.
10. Thiriveedhi A, Venkata Nadh R, Srinivasu N, Kaushal K. Novel hybrid molecules of isoxazole chalcone derivatives: Synthesis and study of in vitro cytotoxic activities. *Letters in Drug Design & Discovery*. 2018 Jun 1;15(6):576-82.
11. Konidala SK, Kotra V, Danduga RC, Kola PK, Bhandare RR, Shaik AB. Design, multistep synthesis and in-vitro antimicrobial and antioxidant screening of coumarin clubbed chalcone hybrids through molecular hybridization approach. *Arabian Journal of Chemistry*. 2021 Jun 1;14(6):103154.
12. Mazimba O. Antimicrobial activities of heterocycles derived from thienylchalcones. *Journal of King Saud University-Science*. 2015 Jan 1;27(1):42-8.
13. Salotra R, Utreja D. A Comprehensive Appraisal of Chalcones and Their Heterocyclic Analogs as Antimicrobial Agents. *Current Organic Chemistry*. 2020 Dec 1;24(23):2755-81.
14. Hamada NM, Sharshira EM. Synthesis and antimicrobial evaluation of some heterocyclic chalcone derivatives. *Molecules*. 2011 Mar;16(3):2304-12.
15. Desai V, Desai S, Gaonkar SN, Palyekar U, Joshi SD, Dixit SK. Novel quinoxaliny chalcone hybrid scaffolds as enoyl ACP reductase inhibitors: Synthesis, molecular docking and biological evaluation. *Bioorganic & medicinal chemistry letters*. 2017 May 15;27(10):2174-80.
16. Ramesh D, Joji A, Vijayakumar BG, Sethumadhavan A, Mani M, Kannan T. Indole chalcones: Design, synthesis, in vitro and in silico evaluation against Mycobacterium tuberculosis. *European journal of medicinal chemistry*. 2020 Jul 15;198:112358.
17. Vásquez-Martínez YA, Osorio ME, San Martín DA, Carvajal MA, Vergara AP, Sanchez E, Raimondi M, Zacchino SA, Mascayano C, Torrent C, Cabezas F. Antimicrobial, anti-inflammatory and antioxidant activities of polyoxygenated chalcones. *Journal of the Brazilian Chemical Society*. 2019;30:286-304.
18. Sökmen M, Khan MA. The antioxidant activity of some curcuminoids and chalcones. *Inflammopharmacology*. 2016 Jun;24(2):81-6.
19. Machala M, Kubínová R, Hořavová P, Suchý V. Chemoprotective potentials of homoisoflavonoids and chalcones of *Dracaena cinnabari*: modulations of drug- metabolizing enzymes and antioxidant activity. *Phytotherapy Research*. 2001 Mar;15(2):114-8.

20. Kumar CS, Loh WS, Ooi CW, Quah CK, Fun HK. Structural correlation of some heterocyclic chalcone analogues and evaluation of their antioxidant potential. *Molecules*. 2013 Oct;18(10):11996-2011.
21. Aktar BS, Sıcak Y, Tok TT, Oruç-Emre EE, Yağlıoğlu AŞ, İyidoğan AK, Öztürk M, Demirtaş I. Designing heterocyclic chalcones, benzoyl/sulfonyl hydrazones: An insight into their biological activities and molecular docking study. *Journal of Molecular Structure*. 2020 Jul 5;1211:128059.
22. Stepanić V, Matijašić M, Horvat T, Verbanac D, Kučerová-Chlupáčová M, Saso L, Žarković N. Antioxidant activities of alkyl substituted pyrazine derivatives of chalcones—in vitro and in silico study. *Antioxidants*. 2019 Apr;8(4):90.
23. Santosh R, Selvam MK, Kanekar SU, Nagaraja GK. Synthesis, Characterization, Antibacterial and Antioxidant Studies of Some Heterocyclic Compounds from Triazole- Linked Chalcone Derivatives. *ChemistrySelect*. 2018 Jun 22;3(23):6338-43.
24. Ardiansah B. Chalcones bearing N, O, and S-heterocycles: Recent notes on their biological significances. *J. Appl. Pharm. Sci*. 2019 Aug 1;9:117-29.
25. Kumari S, Paliwal SK, Chauhan R. An improved protocol for the synthesis of chalcones containing pyrazole with potential antimicrobial and antioxidant activity. *Current Bioactive Compounds*. 2018 Mar 1;14(1):39-47.
26. Kucerova-Chlupacova M, Dosedel M, Kunes J, Soltesova-Prnova M, Majekova M, Stefek M. Chalcones and their pyrazine analogs: synthesis, inhibition of aldose reductase, antioxidant activity, and molecular docking study. *Monatshefte für Chemie-Chemical Monthly*. 2018 May;149(5):921-9.
27. Kasetti AB, Singhvi I, Nagasuri R, Bhandare RR, Shaik AB. Thiazole–Chalcone Hybrids as Prospective Antitubercular and Antiproliferative Agents: Design, Synthesis, Biological, Molecular Docking Studies and In Silico ADME Evaluation. *Molecules*. 2021 Jan;26(10):2847.
28. Brand-Williams W, Cuvelier ME, Berset CL. Use of a free radical method to evaluate antioxidant activity. *LWT-Food science and Technology*. 1995 Jan 1;28(1):25-30.
29. Garcia EJ, Oldoni TL, Alencar SM, Reis A, Loguercio AD, Grande RH. Antioxidant activity by DPPH assay of potential solutions to be applied on bleached teeth. *Brazilian dental journal*. 2012;23:22-7.