

Comparative Anti inflammatory activity of hexane and aqueous extract of *Mucuna pruriens*

RUNNING TITLE: Anti Inflammatory activity of *Mucuna pruriens*

ABSTRACT:

INTRODUCTION: *Mucuna pruriens* is a tropical legume native to Africa and tropical Asia. It is used as a herbalism. The plant is an annual climbing shrub with long vines that can reach a height of 15 meters in length. *Mucuna pruriens* belong to the family legume. It is a herbal drug that is used as a medicine for the treatment of male infertility, nervous disorder, and also as an aphrodisiac. It has been shown that it potentially has more medical importance. It causes orange hair that causes a severe itch when in contact with the skin. It releases a protein called mucanain.

AIM: To determine the anti-inflammatory activity of aqueous and hexane extract of *Mucuna pruriens*.

MATERIALS AND METHODOLOGY: *Mucuna pruriens* dried seeds in powder form were collected. Using a weighing machine, 5 gms of *Mucuna pruriens* were weighed and taken in a conical flask [A]. Another 5gms of *Mucuna pruriens* were weighed and poured into the separate conical flask [B]. In [A] conical flask they add 50ml of distilled water and in another [B] conical flask they add 50ml of hexane solution and cover the mouth of both conical flasks with foil paper. Keep the conical flask inside the shaker for 2 days. After 2 days observations were noted and transferred to a separate test tube and added 10µl, 20µl, 30µl, 40µl, 50µl of the solution in the separate test tube for both hexane and aqueous solution. An anti-inflammatory agent is added to each of the test tubes. Keep all the test tubes in a beaker with a small amount of water. Heat it for 15 min at 50 degrees Celsius and allow it cool. Using a spectrophotometer to observe the reading of both aqueous and hexane solutions. Correlation and analysis were done in SPSS.

RESULT: The hexane solution and aqueous solution of *Mucuna pruriens* have a high percentage of inhibition at the 30µl, 40µl and 50 µl concentration when compared with a standard solution. And hexane has a high percentage of inhibition at 50µl when compared with a standard solution. The obtained data were analysed using Spearman correlation analysis and the non-parametric correlation was statistically significant at $p < 0.05$.

CONCLUSION: The aqueous and hexane extract of *Mucuna pruriens* show good anti-inflammatory activity. In that hexane extract has good anti-inflammatory activity when compared with aqueous extract. And it is used for disease. And further research should be done with the action and mechanism of *Mucuna pruriens*.

KEYWORDS: *Mucuna pruriens*, anti-inflammation, medicinal treatment, Parkinson's disease, a nerve disorder, innovative technology.

HIGHLIGHTS:

- *Mucuna pruriens* belong to the legume family.
- It is used for anti-inflammatory activity and diabetic disorders.
- It is an ornamental plant and used for many medicinal values.
- In that hexane extract has good anti-inflammatory activity when compared with aqueous extract.

- This plant is pharmacologically studied for varied activities like medication, aphrodisiac, antineoplastic, anticonvulsant, antimicrobial activities, etc

INTRODUCTION

Mucuna pruriens may be a common Indian medicative plant that has been employed in ancient Ayurvedic Indian medication for diseases like brain disorder (1). *Mucuna pruriens* belong to the legume family and it is a flavourer drug used for the treatment of male sterility, nervous disorders, and as an aphrodisiac. Its seed area unit is probably of considerable medicative importance. The plant *M. pruriens* named as “velvet bean,” could also be a lively annual climb legume originally from Southern China, Japan, and Asian countries, wherever it's only one occasion widely cultivated as a vegetable crop. The plant has long, slender branches, alternate, lance-like leaves, and white flowers with a bluish-purple, butterfly-shaped whorl. The pods or legumes are bushy, thick, and leatherlike, average it's four inches long, and has formed like string sound holes, and contains seeds. They are brown with thick coats and stiff hairs (2) Emulsifiers are proteins found in living things that are capable of interacting with carbohydrates(3,4)

This plant is pharmacologically studied for varied activities like medication, aphrodisiac, antineoplastic, anticonvulsant, antimicrobial activities, etc (5). Its medicinal drug agents, the nice effort has been spent on the event of medicine for the treatment of inflammation and has many medicinal values. (6)clinical effects and levodopa pharmacokinetics following two different doses of *Mucuna* preparations and comparing them with standard L dopa /carbidopa. It has onset action and has a long time effect without a concomitant increase in dyskinesias.(7)*M.pruriens* have an intrinsic DOPA -decarboxylase inhibitor or act by the mechanism which is independent of levodopa(8). Velvet beans are used like coffee in America. It is a crop and provides fodder and green manure and is also used as an ornamental plant. And treatment reduced the anti-nutritional value of the seed by autoclaving. (9,10)

Inflammation can be defined as the response of the body against various nanoparticles like exogenous and endogenous inciting agents like pathogens, damaged cells, or toxic compounds (11). Inflammation is related to conditions like trauma, heat, chemical, and microorganism, arthritis, gout, and cancer. Inflammation signs are heat, redness, swelling, pain, and loss of

performance. Medicine employed in the inflammation area unit is effective and has facet effects. (12) Medicine like NSAIDs, corticosteroids area units want to manage the pain and inflammation Anti-inflammatory activity was tested in carrageenan evoked rat paw edema model and cotton pellet technique in rat. (13,14). Anti-inflammatory assays were analyzed by a hot plate, formalin, and used probably as a centrally mediated opioid antagonist against analgesia (15). The study involves evaluating the impact of the methanolic extract of *M.pruriens* seed in mice with relevancy delayed hypersensitivity wherever in vitro inflammatory leukocytes mobilize. (16).

This study is needed to reduce the inflammation by the plant extract because plant extract has no toxicity and has fewer side effects which are good for health. This will fulfill various health issue problems and treatment of various nerve disorders. It can be used as an alternative drug in the future. Our team has extensive knowledge and research experience that has translated into high quality publications (17–36). The study aims to determine the anti-inflammatory activity of aqueous extract and hexane extract of *Mucuna pruriens*.

MATERIALS AND METHODOLOGY:

PREPARATION OF THE EXTRACT:

Mucuna pruriens dried seeds in powder form were collected in readymade form. Using a weighing machine, 5 gms of *mucuna pruriens* were weighed and taken in a conical flask [A]. And another 5gms of *Mucuna pruriens* were weighed and poured into the separate conical flask[B]. In [A] conical flask they add 50ml of distilled water and in another [B] conical flask they add 50ml of hexane solution. And cover the mouth of both conical flasks with foil paper. And kept the conical flask inside the shaker for 2 days. After 2 days observations were noted and transferred to a separate test tube and added 10 μ l, 20 μ l, 30 μ l, 40 μ l, 50 μ l of the solution in the separate test tube for both hexane and aqueous solution. Then add an anti-inflammatory agent to each of the test tubes. And keep all the test tubes in a beaker with a small amount of water. And heat the beaker for 15 min at 50 degrees. And cool the beaker. And using a spectrophotometer observe the reading of both aqueous and hexane solutions. The error should be avoided while the

collection of samples. It belongs to a randomized control study. This study is only on anti-inflammatory effects. Correlation and analysis were done in spss and Non parametric correlation was significant at p value less than 0.05. Validations were done by nano experts.

Reagent and Chemicals :

Anti inflammatory :

0.45 mL of Bovine serum albumin (1% aqueous solution)

0.05 mL of *Pterocarpus santalinus* of various fixation (10 μ L,20 μ L,30 μ L,40 μ L,50 μ L)

1N hydrochloric acid

Positive control : Diclofenac Sodium

Negative control : DMSO : Dimethyl sulfoxide.




Figure:1 Sample A is an aqueous extract of *Mucuna pruriens*, where Sample B is a hexane extract of *Mucuna pruriens*. Sample A shows a pale yellow turbid appearance of aqueous extract, where sample B shows a pale yellow translucent appearance of hexane extract.




Figure:2 Anti-inflammatory activity assessed using albumin denaturation assay . Sample A test tube shows an anti-inflammatory activity of hexane and sample B test tube shows the anti-inflammatory activity of aqueous extract.

RESULTS:

ANTI- INFLAMMATORY ACTIVITY:

The present study revealed that at the concentration of 10 μ l, hexane extract has 58% of inhibition and aqueous extract has 57% of inhibition which is more than the standard solution 46%. At a concentration of 20 μ l, hexane extract has 65% of inhibition and aqueous has 61% of inhibition when compared to this standard solution has less zone of inhibition of 60%. At the concentration of 30 μ l, hexane extract has more zone of inhibition 75% when compared with a standard solution, 71%, when compared with this aqueous, has less percentage of inhibition of 69%. At 40 μ l standard solution has more zone of inhibition 78% when compared with the percentage of

inhibition with hexane extract of 76% and an aqueous solution of 70%. At 50 μ l of concentration, the standard solution has more inhibition of 82 % when compared with the %inhibition with hexane extract 78% and an aqueous solution of 71%.




Figure-3: The spectrophotometric absorbance value. When the concentration increases the absorbance level also increases which shows that it has good anti-inflammatory activity in the hexane extract. Non parametric correlation was significant at p value less than 0.05. X axis shows the concentration of the extract in microlitre and the Y axis shows the percentage of inhibition ($r = +1$).




Figure 4: The spectrophotometric value of the percentage of inhibition shows when the concentration increases the percentage of inhibition decreases which shows that it has a good inflammatory effect. Non parametric correlation was significant at p value less than 0.05. X axis shows the concentration of the extract in microlitre and the Y axis shows the percentage of inhibition ($r = -1$)




Figure-5 The Line graph depicts the percentage of inhibition and absorbance level at different concentrations of the extract.




Figure -6 The above bar diagram is on the level of absorbance. When the concentration increases the absorbance level also increases which shows that it has good anti-inflammatory activity in the aqueous extract. Non parametric correlation was significant at p value less than 0.05. X axis shows the concentration of the extract in microlitre and the Y axis shows the percentage of inhibition ($r=+1$).




Figure 7 The above bar graph is on the percentage of inhibition. Concentration increases the percentage of inhibition decreases which shows that it has a good inflammatory effect among the bovine serum albumin. Non parametric correlation was significant at p value less than 0.05. X axis shows the concentration of the extract in microlitre and the Y axis shows the percentage of inhibition ($r = -1$).




Figure 8: the line graph shows the percentage of inhibition and absorbance levels at different concentrations of the extract. In this when the concentration increases the absorbance value also increases and the percentage of the inhibition decreases and it shows the good anti-inflammatory activity

Table: 1

The below table shows the anti -inflammatory activity of mucuna pruriens in the hexane extract along with % of inhibition.

S.No	Concentration (µl)	Wavelength (nm)	Absorbents (wt%)	% of inhibition
1.	10	660	0.112	75.7
2.	20	660	0.11	73.2
3.	30	660	0.111	71.4
4.	40	660	0.112	57.5
5.	50	660	0.11	48.1

Table :2

The below table shows the anti -inflammatory activity of mucuna pruriens in aqueous extract along with % of inhibition

S.No	Concentration (µl)	Wavelength (nm)	Absorbents (wt%)	% of inhibition
1.	10	660	0.161	69
2.	20	660	0.18	66.1

3.	30	660	0.212	61.4
4.	40	660	0.276	52.9
5.	50	660	0.326	47.2

DISCUSSION:

The present study revealed that at the concentration of 10 μ l, hexane extract has 58% of inhibition and aqueous extract has 57% of inhibition which is more than the standard solution 46%. At a concentration of 20 μ l, hexane extract has 65% of inhibition and aqueous has 61% of inhibition when compared to this standard solution has less zone of inhibition of 60%. At the concentration of 30 μ l, hexane extract has more zone of inhibition 75% when compared with standard solution 71%, when compared with this aqueous has less percentage of inhibition of 69%. At 40 μ l standard solution has more zone of inhibition 78% when compared with the percentage of inhibition with hexane extract of 76% and an aqueous solution of 70%. At 50 μ l of concentration, the standard solution has more inhibition of 82 % when compared with the %inhibition with hexane extract 78% and an aqueous solution of 71%. So, Hexane extract shows in (figure 3)more percentage of inhibition when compared to the aqueous solution. In the hexane extract The spectrophotometric absorbance value. When the concentration increases the absorbance level also increases which shows that it has good anti-inflammatory activity in the hexane extract. In (Figure 4) The spectrophotometric value of the percentage of inhibition Shows when the concentration increases the percentage of inhibition decreases which shows that it has a good inflammatory effect. In (Figure 5) the spectrophotometric value shows the percentage of inhibition and absorbance level at different concentrations of the extract. In this when the concentration increases the absorbance value also increases and the percentage of the inhibition decreases and it shows good anti-inflammatory activity. In (Figure 6)the spectrophotometric shows the level of absorbance. When the concentration increases the absorbance level also increases which shows that it has good anti-inflammatory activity in the aqueous extract. In (Figure 7)the spectrophotometric value shows the percentage of inhibition. In this when the

concentration increases the percentage of inhibition decreases which shows that it has a good inflammatory effect among the bovine serum albumin. And in (Figure 8) the spectrophotometric value shows the percentage of inhibition and absorbance level at different concentrations of the extract. In this when the concentration increases the absorbance value also increases and the percentage of the inhibition decreases and it shows good anti-inflammatory activity.

In the previous article, the author concluded that coriander oleoresin with selenium nanoparticles has good anti-inflammatory activity. And it is easy to find and it is eco-friendly. (37). The zinc oxide nanoparticles using the grape seed extract acts as an anti-inflammatory agent with the least side effects. It is used as an anti-inflammatory agent in labs(38). increasing concentrations of has increased anti-inflammatory activity. Chitosan nanoparticles have better anti-inflammatory activity. It can be used as an effective anti-inflammatory agent. Further formulations in various forms such as gel and microspheres can be potentially used for controlled drug delivery in the treatment of periodontitis. (39) The previous author concluded that the effect of aqueous extract of *Mucuna pruriens* leaves on liver enzymes studies. *Mucuna pruriens* extract being used traditionally to improve liver disease conditions and improve antioxidant activity. Aqueous extracts of *M. pruriens* leaves have demonstrated antimicrobial and antioxidant activities with bioactive compounds such as phenols, polyphenols, and tannins. (40) in vitro test was done for anti-inflammatory property. 1% of BSA was taken and 2ml of it was mixed with the plant extract. and it is incubated and kept in the water bath for a few minutes after that absorbance value was estimated in that they concluded that zinc oxide nanoparticle has anti-inflammatory properties(41). Antioxidant properties were done with *M. pruriens*. In vitro study, it has high phenolic content and it has high antioxidant activity. It has the benefit of levodopa where it is used for the treatment of snake bites, Parkinson's disease. And in this, it fails to describe each test, and the hydrophilic compound is used as an identification. DPPH and ABTS assay harm human health with the high phenolic activity of antioxidants. (42) *M. presents* leaves as used for hallucinogenic tryptamines antinutritional activity with phenol tannin. In the invitro scavenge DPPH has radical reactivity with oxygen species (43).

No adverse effect was found in animal blood samples. The seed comet assay has negative results which cause damage to DNA. Alteration plant Wistar albino mice weigh 20.25gm used as a

standard propylene cage and controlled temp and fed and its standard lab diet and water and libitum (44) It poses significant analgesic activity and increases the dose of seed powder. It doesn't observe effects including mutagenicity seed. It has future scope such as it can isolate pure drugs which may be more potent(45). Parkinson's disease (PD) is the most common neurodegenerative and movement disorder, a black tea polyphenol exhibited neuroprotective effects on rotenone-induced apoptosis in SH-SY5Y cells. Advanced studies can be done with this plant extract with a different activity. (46). The limitations of the present study consider a small number of concentration and in vitro study. In the future, this study can be conducted in increased concentration, and clinical studies can be done and animal studies can also be done. It can be used as an alternative drug in the future.

CONCLUSION: The aqueous and hexane extract of *Mucuna pruriens* show good anti-inflammatory activity. In that hexane extract has good anti-inflammatory activity when compared with aqueous extract. It is used for the treatment of various diseases and further research should be done with the action and mechanism of *Mucuna pruriens*.

REFERENCE:

1. Kavitha C, Thangamani C. Amazing bean *Mucuna pruriens*: A comprehensive review [Internet]. Vol. 8, Journal of Medicinal Plants Research. 2014. p. 138–43. Available from: <http://dx.doi.org/10.5897/jmpr2013.5036>
2. Lampariello LR, Cortelazzo A, Guerranti R, Sticozzi C, Valacchi G. The Magic Velvet Bean of *Mucuna pruriens*. Afr J Tradit Complement Altern Med. 2012 Oct;2(4):331–9.
3. Lawal OS, Adebawale KO. Effect of acetylation and succinylation on solubility profile, water absorption capacity, oil absorption capacity and emulsifying properties of mucuna bean (*Mucuna pruriens*) protein concentrate [Internet]. Vol. 48, Nahrung/Food. 2004. p. 129–36. Available from: <http://dx.doi.org/10.1002/food.200300384>

4. Harsha L, Brundha MP. Prevalence of Dental Developmental Anomalies among Men and Women and its Psychological Effect in a Given Population. *Journal of Pharmaceutical Sciences and Research*; Cuddalore. 2017 Jun 20;9(6):869–73.
5. Suresh S, Prithiviraj E, Prakash S. Dose- and time-dependent effects of ethanolic extract of *Mucuna pruriens* Linn. seed on sexual behaviour of normal male rats. *J Ethnopharmacol.* 2009 Apr 21;122(3):497–501.
6. Sathyanarayana N, Pittala RK, Tripathi PK, Chopra R, Singh HR, Belamkar V, et al. Transcriptomic resources for the medicinal legume *Mucuna pruriens*: de novo transcriptome

- assembly, annotation, identification and validation of EST-SSR markers [Internet]. Vol. 18, BMC Genomics. 2017. Available from: <http://dx.doi.org/10.1186/s12864-017-3780-9>
7. Nagar L, Department of Botany, Ch. Charan Singh University, Meerut, UP, India, et al. Sequence to Structure Analysis of DOPA Protein from *Mucuna pruriens*: A Computational Biology Approach [Internet]. International journal of Emerging Trends in Science and Technology. 2015. Available from: <http://dx.doi.org/10.18535/ijetst/v2i8.12>
 8. Ozcinar B, Balci G, Buyukkaya R, Haslak A. Appendix Inflammation with Varicella Zoster Infection [Internet]. Vol. 12, Istanbul Medical Journal. 2011. p. 196–8. Available from: <http://dx.doi.org/10.5505/1304.8503.2011.30502>
 9. Hussian G, Manyam BV. *Mucuna pruriens* proves more effective than L-DOPA in Parkinson's disease animal model [Internet]. Vol. 11, Phytotherapy Research. 1997. p. 419–23. Available from: [http://dx.doi.org/10.1002/\(sici\)1099-1573\(199709\)11:6<419::aid-ptr120>3.0.co;2-q](http://dx.doi.org/10.1002/(sici)1099-1573(199709)11:6<419::aid-ptr120>3.0.co;2-q)
 10. Timothy CN, Samyuktha PS, Brundha MP. Dental pulp Stem Cells in Regenerative Medicine--A Literature Review. Research Journal of Pharmacy and Technology. 2019;12(8):4052–6.
 11. Medzhitov R. Origin and physiological roles of inflammation [Internet]. Vol. 454, Nature. 2008. p. 428–35. Available from: <http://dx.doi.org/10.1038/nature07201>
 12. Rane M, Suryawanshi S, Patil R, Aware C, Jadhav R, Gaikwad S, et al. Exploring the proximate composition, antioxidant, anti-Parkinson's and anti-inflammatory potential of two neglected and underutilized *Mucuna* species from India [Internet]. Vol. 124, South African Journal of Botany. 2019. p. 304–10. Available from: <http://dx.doi.org/10.1016/j.sajb.2019.04.030>
 13. Iauk L, Galati EM, Kirjavainen S, Forestieri AM, Trovato A. Analgesic and Antipyretic Effects of *Mucuna Pruriens*. Pharm Biol. 2008 Sep 27;31(3):213–6.
 14. Hannah R, Ramani P, Brundha MP, Sherlin HJ, Ranjith G, Ramasubramanian A, et al. Liquid Paraffin as a Rehydrant for Air Dried Buccal Smear. Research Journal of Pharmacy

and Technology. 2019;12(3):1197–200.

15. Avoseh ON, Ogunwande IA, Ojenike GO, Mtunzi FM. Volatile Composition, Toxicity, Analgesic, and Anti-Inflammatory Activities of *Mucuna pruriens*. *Nat Prod Commun*. 2020 Jul 1;15(7):1934578X2093232.
16. Upadhyay P. Phytochemistry and pharmacological activity of *Mucuna pruriens*: A review. *International Journal of Green Pharmacy (IJGP)* [Internet]. 2017 May 16 [cited 2021 Feb 18];11(02). Available from:
<https://www.greenpharmacy.info/index.php/ijgp/article/view/916>
17. Jayaseelan VP, Paramasivam A. Emerging role of NET inhibitors in cardiovascular diseases. *Hypertens Res*. 2020 Dec;43(12):1459–61.
18. Sivakumar S, Smiline Girija AS, Vijayashree Priyadharsini J. Evaluation of the inhibitory effect of caffeic acid and gallic acid on tetR and tetM efflux pumps mediating tetracycline resistance in *Streptococcus* sp., using computational approach. *Journal of King Saud University - Science*. 2020 Jan 1;32(1):904–9.
19. Smiline Girija AS. Delineating the Immuno-Dominant Antigenic Vaccine Peptides Against gacS-Sensor Kinase in *Acinetobacter baumannii*: An in silico Investigational Approach. *Front Microbiol*. 2020 Sep 8;11:2078.
20. Iswarya Jaisankar A, Smiline Girija AS, Gunasekaran S, Vijayashree Priyadharsini J. Molecular characterisation of csgA gene among ESBL strains of *A. baumannii* and targeting with essential oil compounds from *Azadirachta indica*. *Journal of King Saud University - Science*. 2020 Dec 1;32(8):3380–7.
21. Girija ASS. Fox3+ CD25+ CD4+ T-regulatory cells may transform the nCoV's final destiny to CNS! *J Med Virol* [Internet]. 2020 Sep 3; Available from:
<http://dx.doi.org/10.1002/jmv.26482>
22. Jayaseelan VP, Ramesh A, Arumugam P. Breast cancer and DDT: putative interactions, associated gene alterations, and molecular pathways. *Environ Sci Pollut Res Int*. 2021 Jun;28(21):27162–73.

23. Arumugam P, George R, Jayaseelan VP. Aberrations of m6A regulators are associated with tumorigenesis and metastasis in head and neck squamous cell carcinoma. *Arch Oral Biol.* 2021 Feb;122:105030.
24. Kumar SP, Girija ASS, Priyadharsini JV. Targeting NM23-H1-mediated inhibition of tumour metastasis in viral hepatitis with bioactive compounds from *Ganoderma lucidum*: A computational study. *pharmaceutical-sciences* [Internet]. 2020;82(2). Available from: <https://www.ijpsonline.com/articles/targeting-nm23h1-mediated-inhibition-of-tumour-metastasis-in-viral-hepatitis-with-bioactive-compounds-from-ganoderma-lucidum-a-comp-3883.html>
25. Girija SA, Priyadharsini JV, Paramasivam A. Prevalence of carbapenem-hydrolyzing OXA-type β -lactamases among *Acinetobacter baumannii* in patients with severe urinary tract infection. *Acta Microbiol Immunol Hung.* 2019 Dec 9;67(1):49–55.
26. Priyadharsini JV, Paramasivam A. RNA editors: key regulators of viral response in cancer patients. *Epigenomics.* 2021 Feb;13(3):165–7.
27. Mathivadani V, Smiline AS, Priyadharsini JV. Targeting Epstein-Barr virus nuclear antigen 1 (EBNA-1) with *Murraya koengii* bio-compounds: An in-silico approach. *Acta Virol.* 2020;64(1):93–9.
28. Girija As S, Priyadharsini J V, A P. Prevalence of Acb and non-Acb complex in elderly population with urinary tract infection (UTI). *Acta Clin Belg.* 2021 Apr;76(2):106–12.
29. Anchana SR, Girija SAS, Gunasekaran S, Priyadharsini VJ. Detection of *csgA* gene in carbapenem-resistant *Acinetobacter baumannii* strains and targeting with *Ocimum sanctum* biocompounds. *Iran J Basic Med Sci.* 2021 May;24(5):690–8.
30. Girija ASS, Shoba G, Priyadharsini JV. Accessing the T-Cell and B-Cell Immuno-Dominant Peptides from *A.baumannii* Biofilm Associated Protein (bap) as Vaccine Candidates: A Computational Approach. *Int J Pept Res Ther.* 2021 Mar 1;27(1):37–45.
31. Anita R, Paramasivam A, Priyadharsini JV, Chitra S. The m6A readers YTHDF1 and YTHDF3 aberrations associated with metastasis and predict poor prognosis in breast cancer

- patients. *Am J Cancer Res.* 2020 Aug 1;10(8):2546–54.
32. Varghese SS, Ramesh A, Veeraiyan DN. Blended Module-Based Teaching in Biostatistics and Research Methodology: A Retrospective Study with Postgraduate Dental Students. *J Dent Educ.* 2019 Apr;83(4):445–50.
 33. Arvind P TR, Jain RK. Skeletally anchored forsus fatigue resistant device for correction of Class II malocclusions-A systematic review and meta-analysis. *Orthod Craniofac Res.* 2021 Feb;24(1):52–61.
 34. Ramadurai N, Gurunathan D, Samuel AV, Subramanian E, Rodrigues SJL. Effectiveness of 2% Articaine as an anesthetic agent in children: randomized controlled trial. *Clin Oral Investig.* 2019 Sep;23(9):3543–50.
 35. Venugopal A, Vaid N, Bowman SJ. Outstanding, yet redundant? After all, you may be another Choluteca Bridge! *Semin Orthod.* 2021 Mar 1;27(1):53–6.
 36. Mathew MG, Samuel SR, Soni AJ, Roopa KB. Evaluation of adhesion of *Streptococcus mutans*, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary molars: randomized controlled trial [Internet]. Vol. 24, *Clinical Oral Investigations*. 2020. p. 3275–80. Available from: <http://dx.doi.org/10.1007/s00784-020-03204-9>
 37. Ram AJ, Roy A, Rajeshkumar S, Lakshmi T. ANTI INFLAMMATORY ACTIVITY OF CORIANDER OLEORESIN MEDIATED SELENIUM NANOPARTICLES. *PLANT CELL BIOTECHNOLOGY AND MOLECULAR BIOLOGY.* 2020 Nov 6;106–11.
 38. Vaishnavi Devi B, Rajasekar A, Rajeshkumar S. ANTIINFLAMMATORY ACTIVITY OF ZINC OXIDE NANOPARTICLES SYNTHESISED USING GRAPE SEED EXTRACT: AN *in vitro* STUDY. *PLANT CELL BIOTECHNOLOGY AND MOLECULAR BIOLOGY.* 2020 Aug 26;6–16.
 39. Kumaran H, Gajendran PL, Arjunkumar R, Rajesh S. ANTI INFLAMMATORY ACTIVITY OF LYCOPENE AND ITS BASED CHITOSAN NANOFORMULATION-AN *In vitro* STUDY. *PLANT CELL BIOTECHNOLOGY AND MOLECULAR BIOLOGY.*

2020 Nov 10;91–100.

40. Adam AA. EFFECT OF AQUEOUS EXTRACT OF MUCUNA PRURIENS LEAVES ON LIVER AND KIDNEY FUNCTION MARKERS IN WISTAR ALBINO RATS. 2016 Feb 15 [cited 2021 Feb 18]; Available from:
https://www.researchgate.net/publication/294554442_EFFECT_OF_AQUEOUS_EXTRACT_OF_MUCUNA_PRURIENS_LEAVES_ON_LIVER_AND_KIDNEY_FUNCTION_MARKERS_IN_WISTAR_ALBINO_RATS
41. Agarwal H, Shanmugam V. A review on anti-inflammatory activity of green synthesized zinc oxide nanoparticle: Mechanism-based approach. *Bioorg Chem.* 2020 Jan;94:103423.
42. Chel-Guerrero L, Galicia-Martínez S, Acevedo-Fernández JJ, Santaolalla-Tapia J, Betancur-Ancona D. Evaluation of Hypotensive and Antihypertensive Effects of Velvet Bean (*Mucuna pruriens* L.) Hydrolysates. *J Med Food.* 2017 Jan;20(1):37–45.
43. Katzenschlager R, Evans A, Manson A, Patsalos PN, Ratnaraj N, Watt H, et al. *Mucuna pruriens* in Parkinson's disease: a double blind clinical and pharmacological study. *J Neurol Neurosurg Psychiatry.* 2004 Dec;75(12):1672–7.
44. Modi KP, Patel NM, Goyal RK. Estimation of L-Dopa from *Mucuna pruriens* LINN and Formulations Containing *M. pruriens* by HPTLC Method [Internet]. Vol. 56, *CHEMICAL & PHARMACEUTICAL BULLETIN.* 2008. p. 357–9. Available from:
<http://dx.doi.org/10.1248/cpb.56.357>
45. Ghosal S, Singh S, Bhattacharya S. ALKALOIDS OF *MUCUNA PRURIENS* CHEMISTRY AND PHARMACOLOGY [Internet]. Vol. 19, *Planta Medica.* 1971. p. 279–84. Available from: <http://dx.doi.org/10.1055/s-0028-1099642>
46. Barma MD, Kannan SD, Indiran MA, Rajeshkumar S, Pradeep Kumar R. Antibacterial Activity of Mouthwash Incorporated with Silica Nanoparticles against *S. aureus*, *S. mutans*, *E. faecalis*: An in-vitro Study. *Journal of Pharmaceutical Research International.* 2020 Aug 24;25–33.