

Percussion in Cardiovascular Examination of a Tombstone

Abstract

The growing abundance of technology in medical field has led to doctors' sensory de-skilling. Technology is seen replacing the clinical diagnosis based on sensory perception. Percussion as a skill has lost from clinical practice. Though percussion disappeared from wards, it is still taught in medical colleges. Percussion is a very vital component in bedside examination of the patient. It was Auenbrugger who first thought of tapping the chest of patients to determine the cardiac size, borders, and presence of pericardial fluid, its extent and related pathologies. Percussion used to be a critical component of clinical heart disease evaluation. There was rapid advancement of technology in last century and with the introduction of technology in medical field, the diagnosis and treatment of disorders and diseases has improved to a great extent. Technology has a very huge impact on medical field, now there is hardly any major disease which can be diagnosed with clinical skills alone and without involving technology for confirmation. With the development of new imaging methods such as X-ray and echocardiography, percussion has been lost. Percussion can be a valuable bedside aid in determining the cardiac size, borders and to find the presence and extent of pericardial fluid. There is a possibility of misunderstanding if used alone, but when combined with other clinical characteristics, it appears to be it proves to be a powerful bedside tool in distinguishing tamponed from acute massive pulmonary embolism until echocardiography confirms the diagnosis. Cardiac percussion is not just tapping the chest and hearing the sound produced, it is also feeling the touch and observing the vibrations produced noting the tone, thus percussion takes a considerable amount of time, practice and a mindset to master as a clinical skill. Thus teaching the correct way to percuss right from the initial years in the medical school is extremely necessary. With proper knowledge and experience cardiac percussion can prove to be a very helpful tool in certain cardiac pathologies.

Keywords: Auenbrugger, Laennec, Chest percussion, Auscultation

Introduction

There was a time when percussion was a very important component in clinical assessment of heart disease. Percussion has been lost with the introduction of new imaging techniques like X-ray and echocardiography, both of which are more precise in determining the cardiac size, borders and to find the presence and extent of pericardial fluid.

In this rapid world of technology, do cardiologists spend time percussing the chest, to detect cardiomegaly and fluid in pericardium, when in minutes they can have a more precise and specific diagnosis with echocardiography? Honestly they don't. However, with the skill of percussion they can get quick information regarding pericardial effusion, pending verification with echocardiography. If used alone, there are chances of error but when used along with other clinical features it proves to be vital bedside tool in differentiating tamponed from acute massive pulmonary embolism until verification from echocardiography.

Should we still practice percussion? Obviously yes. Cardiac percussion is an art and a science and medical students must learn it. Also, the practicing cardiologists should use this tool in their clinical practice as it is of vital importance in certain cardiac conditions. ^[1]

History of Percussion. Leopold Auenbrugger, native of Austria, was the inventor of percussion

Leopold Auenbrugger (1722-1809) who invented percussion and René Laennec who is known as the father of modern physical examination. The inventor of percussion joined father of modern physical examination. When there was an event of the two-hundredth anniversary of the development of the stethoscope in the year 1816, I proceeded to Austria, Auenbrugger's hometown to further investigate more on material effects. This resulted in clearing out my vision though a bit translucent and also to a bit. Apparently, succeeding generations have not gratified to this incredible pioneer of clinical science. All the more reason for knowing and protecting what is left of him: buildings, monuments, portraits... Anyway, every time a doctor performs the chest percussion, Leopold Auenbrugger is called to the mind and tribute today, as he was in the past, (that is billions of times each year).^[2]

Let's start by recalling the traditional story. Leopold von Auenbrugger was son of owner of an inn, he became experienced observing different sounds of wine and beer barrels in his dad's vault. When he grew up he studied medicine and he used to treat his patients just like beer barrels. As a result the incredible thoracic percussion was discovered. This resulted to his great discovery of thoracic percussion, which he stated as void patients make the most clamor. His companions shut their eyes to his work and refused his correlation of patient and brew vessel. This antique transformation was in twilight until it got saved by Napoleon's physician named Baron Corvisart. He practiced and taught percussion. Because of the huge broadcasting, the skill of percussion got eternally loaded on medical students. ^[3]

It's not surprising that a musician's psyche led to the development of percussion. Leopold Auenbrugger was a physician, who composed some music for a queen of Austria. The music and medicine met in mind of this man, has its roots since he saw his dad tapping on wine barrels to decide its contents.

He studied at Leyden Medical School and graduated in 1752. He was a bright student and he wrote on aphorisms of Hippocrates for his thesis. At Spanish Hospital he worked without receiving any payment and then he got promoted to a higher post that is of an attending physician. He decided to drop off from working in hospitals in 1762. He explained the chest percussion in a text called "Inventum Novum" published in 1761.

He defined different lungs' sounds in different cases, but rather than understanding his approach, he was held accountable for piracy! He was charged that he copied from Hippocrates, since Hippocrates explained the succussion splash, the pleural rub,. But this was not true as Auenbugger pioneered chest percussion in normal and abnormal variants and in physical diagnosis.

He wrote, that what he has written, he has verified repeatedly by witnessing through his own senses and surrounded by strenuous, exhausting toilsome exertions; though observing, on each instant, against the alluring impact of confidence." He was fortunate that, Napoleon's physician, Corvisart, unearthed his work and immortalized it by publishing it, otherwise his innovation might have lost in darkness. ^[4]

According to Dr. Anna Harris of Maastricht University

One fine morning I received an email message from a professor of pulmonary medicine, it expresses that ‘percussion note is right in certain conditions and as he would see it is a kind of sound’. I began to pay more attention to percussion, particularly the respiratory examination.^[5] However, it wasn't long before other doctors stated that the method was all about touch, not sound. One professor of internal medicine stated, reclining in his office seat, that he could perform percussion even if he was unable to hear or deaf as it is about the sensation of percussion more than the sound itself and he knew it was contradictory to my findings.’ I was told by plenty of educators and learners that percussion was more about experiencing the vibrations than listening to the music, which was particularly true in loud hospital rooms. Not only was the sound created by touch, but the discoveries were also sensed by touch, via the listening fingers. There was no agreement on whether percussion was about sound or touch, and it didn't appear to issue eventually: phrases like “You are listening now”, "You are watching out at this point. Now you are thinking, 'Everything's incorporated!' a professor from Melbourne asserted.

Auenbrugger's instructions are supported by a strong correlation between sound and perception through touch in practice of percussion. The mixed character of the sensory perception of percussion is frequently emphasized in clinical examination texts. The Oxford Handbook of Clinical Medicine, for example, gives the following directions for percussion in a small yellow book placed on top of any decent medical student's heap of ipads, clipboards, and papers:

All locales, including the axillae, clavicles, and supraclavicular regions, ought to be percussed. Tune in and feel for the sound's inclination and evenness.

The practice of percussion is instructed as a listening or touch strategy, yet in addition as a mix of the two. Other textbooks back this up: Talley and O'Connor write, " The percussion note's vibe and the sound both are equally essential" (1992: 111). In the Essentials of Respiratory Medicine, Cade and Pain (1988) explain that ‘percussion brings about a perplexing arrangement of signs giving the analyst both the hear-able data of pitch and the material data of vibration or obstruction, and mirrors the proportion of air to solid structures under the space being percussed.'

Many historical, sociological, and anthropological studies of medical practise treat touching and hearing differently, assuming the faculties as free elements with interrelated tangible organs; the ears hear, the eyes see, the fingers contact, etc. Over time, these tidy divisions have become problematic. For example, David Howes (2003) discusses occurrences of intersense, a word that preserves a sense of sensory separation. Although Aristotle is frequently described as considering the five senses to be distinct, his effort on common sense shows an endeavor to look past the five classifications, to explore an intervening sense which helps us in feeling. Heller-Roazen finishes good judgment later philosophical work, expressing that the conventional issue of the center sense got new earnestness in later advancement, recast as an internal touch, a vibe of one's crucial movements, and one's experience of being alive. This was a sensation that couldn't be readily localised or defined; for some, it was skin, for others viscera, muscles, or blood, and for still others, it was everything at once (Heller-Roazen, 2007: 251).^[6]

The essay written by Corvisart has been broken down into six sections. The pericardium, heart muscle, tendinous components, varia and aorta are the five sites of organic change. The final section covers aetiology, diagnosis, prognosis, and therapy options. The discussion on pathological lesions is illustrated by case studies. Corvisart reported clinical entities that reflect the symptoms currently linked with coronary artery disease, but he seems to have been ignorant of, or hesitant to acknowledge, the ideologies on this ailment advanced over the Channel, probably due to his political context. He linked mitral valve stenosis to a perceptible "thrill," or it can be understood like some pulsations can be sensed by the fingertips on the chest.^[7] He was especially keen on cardiovascular enlargement, what was labelled by him as "aneurysm," and classified it into "active" (thickened muscle) and "passive" (non-thickened muscle) categories (thin muscle). Corvisart attempted to identify aneurysm in the clinical environment by placing his hand on chest of the patient or percussing the area of precordium to evaluate the amount of dullness in tone cardiac, and he felt positive about his capacities to figure out active from passive aneurysm.^[8] After two years, in his interpretation of Auenbrugger's *Inventum novum*, he emphasised on the importance of percussion.^[9] Corvisart used the face's tinge and presentation, the condition of the pulse, the presence of edema, difficulty in breathing, hepatomegaly, thrill that can be palpated, predictions of lung congestion, pleural effusion, or cardiomegaly as obtained by percussion to define the clinical indications of cardiac illness. There were numerous

symptoms, but even Corvisart admitted that it was often difficult to tell the difference between heart and lung problems. ^[10]

Laennec shared Corvisart's primary interest in myocardial volume alterations, but he was not blind to other elements of heart disease. He was able to distinguish between two types of congenital cardiac illness: patent ductus arteriosus and septal defects. He also left a unique depiction of alteration of valves, dubbed "globular excrescences," whose origin is still unknown.

^[11] He reported his first publication, a clinico-pathological case of a 22-year-old male who had hydrothorax and distention of right ventricle was anticipated by Corvisart, by the help of carcass percussion immediately before dissection, in 1802, at the age of twenty-one.

Corvisart had also suggested that "organic lesions" might be detected on the left side of the heart, and there was a possibility of discovering principal cause of the disease. ^[12] The autopsy found ossification of the mitral valve, verifying the accuracy of his prognosis regarding the lung and right-side ventricle.

According to P Rayer, his capacity to give a right analysis dependent on an association between the clinical viewpoint and pathological anatomy frequently dumbfounded his audience members.

^[13] He was a "pathological anatomy pioneer" who "autopsied every patient dying on his wards." During clinical examinations, Corvisart often placed his hand over his patient's chest. He came up with a very defining information by using this technique: the "murmur" of the heart. ^[14] After palpation, he also performed percussion, which Auenbrugger had envisioned forty years before.

^[15]

Despite his cautious wording, the last argument of Laennec, that disproved the vegetations' venereal etiology, was persuasive. In 1832, Laennec's observations were confirmed by James Hope. ^[16]

As a result, until 1830, there was no actual creation of a clinical entity. Furthermore, the claim which suggests the vegetative involvement of valves was related to syphilis got disparaged. 'Cardiac workings' gets affected by a few bruises, particularly specific indications or "noises," were noticed. Palpitations, respiratory difficulties, swelling of lower limbs, rheumatic pain, cough, pyrexia, tiredness, and dyspnea were all reported quite correctly, albeit vegetation of valves was not related to them. During autopsy, "vegetations," "concretions," or "outgrowths" on the cardiac valves or its chambers did not produced the diagnosis of a particular illness. As a result, it was often impossible to tell what was there from the start, what arrived subsequently,

and even what may have appeared after death in an illness that got increasingly convoluted on the long run. Further developed clinical perception procedures, as well as data gleaned from percussion as well as auscultation, helped to construct manifestations that matched to postmortem lesions. The hunt for a way to identify these sores in a living patient began then. In Paris, in particular, a genuine "research spirit" transformed the hospital into a type of laboratory as well as a teaching institution. ^[17]

The aye-aye (*Daubentonia madagascariensis*), a monkey discovered uniquely on Madagascar's island, is absolutely and unconditionally reliant on percussion to search its supper, that contains creepy crawly hatchlings burrowing underneath tree husk. The creature expands a long, spindly third digit into the tunnel and gathers up its feast in the wake of finding it with percussion (Short, 1995). ^[18]

Francis Henry Williams contrasted sizes of heart estimated by percussion with results on chest radiography, a novel method at the time, and heart weight recorded in 546 autopsy in 1899. He ruled out cases of emphysema," that puts percussion on loss." He discovered where the cardiac size was close to regular, percussion errors were minimal. Error was particularly common where cardiac dimensions were lower than regular. Percussive errors were less common but larger in size when the heart was enlarged (Jarcho, 1969). ^[19]

Cardiac Percussion Technique Reliability vs. Cardiothoracic Ratio and Transthoracic Echocardiography in Determining Left Ventricular Size

Background

Cardiac percussion (CP) is a bedside method for determining the cardiac boundaries, chamber size, and pericardial fluid extent. Despite the fact that new cardiac imaging tools have changed the procedures for detecting enlarged heart, CP as a skill is chosen most often, particularly in emergency cases. Notwithstanding the way that transthoracic echocardiography (TTE) is acknowledged more precise for deciding size of the left ventricle, Cardiac percussion can appraise size of the left ventricle with equivalent precision in a proper clinical setting. We will likely research the relationship of left ventricular size utilizing the Cardiac percussion method and contrast it with standard transthoracic echocardiography and the cardiothoracic proportion (CTR) on chest roentgenography.

Methodology

There was a research conducted in order to find out the correlation between the findings of Cardiac Percussion which we get easily bedside and the results of transthoracic echocardiography along with reference from roentgenography. The Telemetry unit of BronxCare Health System was the subject of this descriptive research. The sick person was kept at a Gatch angle of 30 to 45 degrees as part of the physical examination in this bedside method. Along midpoint of clavicle is percussed to the gladiolus of the sternum was used to measure cardiac dullness. The size of left ventricle (in centimetres) and the point of reference were then measured using a ruler (middle left parasternal border). Following that, during the same hospitalisation, we linked findings with transthoracic echocardiography dimensions of left ventricular end-diastolic diameter (LVEDD). We also used chest roentgenography to measure the CTR (CXR). The clinical history and imaging findings were hidden from the percussors. The left ventricular (LV) volumes were calculated using the Teicholz technique. The variability in each estimate was estimated using the coefficient of variation (COV). Pitman's test was utilized to investigate the distinction in change between any pair of measurements, and Bland-Altman plots were created to look at any estimations.

Results

As the above procedure was carried out the following results were obtained. Our 200 patients were on average 63(15.81) years old, with female accounting for 51 percent of the total. The average BMI was 28.5 KG/M2, and 33% of the population had a normal BMI. A total of 22% of the participants were found to have a diagnosis of heart failure. The left ventricular diameter evaluated with the Cardiac percussion technique was analogous to the left ventricular end-diastolic diameter obtained using transthoracic echocardiography (Mean 4.65cm 0.67 vs 4.69cm 0.76, COV 7.6%). The left ventricular end-diastolic diameter was shown to have no association with the LV diameter measured by CXR (5.79cm1.21, COV 14.3%, $r=-0.475$, $p0.001$). The left ventricular volumes determined using the Cardiac Percussion method were found to be in agreement with the left ventricular volumes assessed using transthoracic echocardiography. (14.9 percent coefficient of variation)

Conclusions

The CP technique is a simple bedside approach for determining the size and volume of the left ventricle. It has been found to have a strong relationship with LVEDD as assessed by TTE. In the estimate of LV size, CXR was shown to have no connection. The research underlines the need of utilising a physical examination approach to estimate LV size at the bedside. Our findings will need to be confirmed in future investigations. ^[20-29]

The developing plenitude of innovation in clinical field has prompted specialists' tangible de-skilling. Innovation is seen supplanting the clinical determination dependent on tactile insight. Percussion as an expertise has lost from clinical practice. Percussion is an extremely essential part in bedside assessment of the patient. It was Auenbrugger who initially considered tapping the chest of patients to decide the heart size, boundaries, and presence of pericardial liquid, its degree and related pathologies. Percussion used to be a basic part of clinical coronary illness assessment. There was quick progression of innovation in last century and with the presentation of innovation in clinical field, the analysis and treatment of problems and infections has improved by and large. Percussion can be a significant bedside help in deciding the cardiovascular size, borders and to track down the presence and degree of pericardial liquid. There is plausible of misconception whenever utilized alone, yet when joined with other clinical

attributes, it has all the earmarks of being it ends up being an amazing bedside practice in recognizing tamponed from acute massive pulmonary embolism until echocardiography affirms the determination.

Hence, this winds up to a conclusion that cardiac percussion should be learned by medical students and cardiac percussion should be used by practicing cardiologists and they should utilize this tool in their clinical practice as it proves to be of vital significance in specific heart conditions.

UNDER PEER REVIEW

References:

1. Hajar R. Cardiac percussion: a lost art?. *Heart Views*. 2011; 12(2):82. doi:10.4103/1995-705X.86022
2. Borghi L. Tapping on the chest of history. Lost and found memories of Leopold Auenbrugger, inventor of percussion, in Austria and beyond. *Acta Med Hist Adriat*. 2018 Jun;16(1):127-144.
3. LAENNEC, R. T. H. *Traiti de l'Auscultation Midiate*. Second edition (1826), reprinted. Paris, Asselin, 1879, P. 5.
4. Cummins, S. L. (1945). Auenbrugger and Laennec: The Discoverers of Percussion and Auscultation. *Proceedings of the Royal Society of Medicine*, 38(8), 409–412.
5. Leopold Auenbrugger's *Inventum Novum*. (1923). *JAMA: The Journal of the American Medical Association*, 80(8), 576. doi:10.1001/jama.1923.02640350058042
6. Smith, J. J. (1962). The *Inventum Novum* of Joseph Leopold Auenbrugger. *Bulletin of the New York Academy of Medicine*, 38(10), 691–701.
7. Percussion is also conducted in the cardiac and abdominal examinations of the body; however, it performs only a minor role in these examinations and I focus here on the respiratory examination where it is more prominent. My focus during fieldwork was on respiratory medicine.
8. Medicine has also been described as a craft by others, such as Hirschauer (1991), who likens medicine to building and carpentry, tailoring, sailors' knots and the butcher's trade. More generally Latour and Woolgar (1979) also discuss craftwork in science, Fujimura (1992) examines crafting of science, and Kondo (1990) the crafting of selves in the Japanese workplace.
9. *Ibid.*, p. 185. Support for this statement is contained in a manuscript consultation published by P. Soulie, 'Corvisart et le diagnostic clinique du retrecissement mitrale', *Hist. de la Med.*, 1955, no. 7, pp. 55-64.
10. Corvisart, *op. cit*, note 18 above, pp. 126-8.
11. Leopold Auenbrugger, *Nouvelle mehode pour reconnaitre les maladies de la poitrine*, trans. J. N. Corvisart, Paris, Migneret, 1806, pp. 420-32.

12. Corvisart, op. cit, note 18 above, pp. 315-30.
13. Laennec, op. cit., note 3 above, 1826, vol. 2, pp. 630-51. See also Frank D. Mann and Ruth J. Mann, 'Laennec as a critical pathologist', *J. Hist. Med.*, 1981, 36: 446-54, esp. pp. 451-2. The valvular changes may be due to disseminated intravascular coagulation. Joel D. Howell has suggested that the entity is probably that of marantic endocarditis (personal communication).
14. Rene Laennec, 'Observation d'une maladie du coeur', *J. de Medecine*, An x (1802), 4: 265-307.
15. Pierre Rayer, *Sommaire d'une histoire abregee de l'anatomie pathologique*, Paris, 1818, p. 131.
16. Erwin H Ackerknecht, *Medicine at the Paris Hospital, 1794-1845*, Baltimore, Johns Hopkins Press, 1967, pp. 84; *La medecine hospitaliere a Paris*, Payot, 1986, pp. 109-12.
17. Corvisart, op. cit., note 18, above, pp. 182-5: "By observing the disorder of the circulation, the practitioner may find, in the living man, I should presume, certain signs of this species of affection.... Beside the general signs of the diseases of the heart,
18. James Hope, *A treatise on the diseases of the heart and great vessels*, London, W Kidd, 1832, p. 506.
19. Some of the classic works on this subject are: Erwin Ackerknecht, op. cit., note 20 above; Michel Foucault, *The birth of the clinic*, transl. A M Sheridan, London, Tavistock, 1973; French title, *Naissance de la clinique*, Paris, PUF, 1963; Toby Gelfand, *Professionalizing modern medicine: Paris surgeons and medical science and institutions in the 18th century*, Westport, Conn., Greenwood Press, 1980; Russell Maulitz, *Morbid appearances: the anatomy of pathology in the early nineteenth century*, Cambridge University Press, 1987.
20. Short RV. Percussion as a way of life. *Lancet* 1995;346:67-68.
21. Jarcho S. Percussion of the heart contrasted with roentgen examination (Williams, 1899). *Am J Cardiol* 1969;23:845-849.
22. Syed MS, Zaidi AA, Bhandari M, Bella JN, Vittorio TJ. Reliability of Cardiac Percussion Technique to Determine Left Ventricular Size in Comparison with Cardiothoracic Ratio and Transthoracic Echocardiography. *Journal of Cardiac Failure*. 2019;(25):S38

23. Bittner, Vera A., Michael Szarek, Philip E. Aylward, Deepak L. Bhatt, Rafael Diaz, Jay M. Edelberg, Zlatko Fras, et al. "Effect of Alirocumab on Lipoprotein(a) and Cardiovascular Risk After Acute Coronary Syndrome." *JOURNAL OF THE AMERICAN COLLEGE OF CARDIOLOGY* 75, no. 2 (January 21, 2020): 133–44. <https://doi.org/10.1016/j.jacc.2019.10.057>.
24. Garg, Mayank, and Sandip Mohale. "Prevalence of Metabolic Obesity Normal Weight (MONW) in Cardiovascular Disease Patients - A Hospital-Based Case Control Study." *JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS* 9, no. 34 (August 24, 2020): 2427–31. <https://doi.org/10.14260/jemds/2020/528>.
25. Kamble, T. K., Ankita Kapse, Sunil Kumar, Sourya Acharya, and Aiswarya Ghule. "Study of Myocardial Performance Index in Prediabetes and Its Correlation with Other Cardiovascular Risk Factors." *JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS* 9, no. 10 (March 9, 2020): 721–25. <https://doi.org/10.14260/jemds/2020/157>.
26. Acharya, Sourya, Samarth Shukla, and Anil Wanjari. "Subclinical Risk Markers for Cardiovascular Disease (CVD) in Metabolically Healthy Obese (MHO) Subjects." *JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH* 13, no. 6 (June 2019): OC1–6. <https://doi.org/10.7860/JCDR/2019/41317.12890>.
27. Goodman, Shaun G., Philip E. Aylward, Michael Szarek, Vakhtang Chumburidze, Deepak L. Bhatt, Vera A. Bittner, Rafael Diaz, et al. "Effects of Alirocumab on Cardiovascular Events After Coronary Bypass Surgery." *JOURNAL OF THE AMERICAN COLLEGE OF CARDIOLOGY* 74, no. 9 (September 3, 2019): 1177–86. <https://doi.org/10.1016/j.jacc.2019.07.015>.
28. Ray, Kausik K., Helen M. Colhoun, Michael Szarek, Marie Baccara-Dinet, Deepak L. Bhatt, Vera A. Bittner, Andrzej J. Budaj, et al. "Effects of Alirocumab on Cardiovascular and Metabolic Outcomes after Acute Coronary Syndrome in Patients with or without Diabetes: A Prespecified Analysis of the ODYSSEY OUTCOMES Randomised Controlled Trial." *LANCET DIABETES & ENDOCRINOLOGY* 7, no. 8 (August 2019): 618–28. [https://doi.org/10.1016/S2213-8587\(19\)30158-5](https://doi.org/10.1016/S2213-8587(19)30158-5).
29. Schwartz, G. G., P. G. Steg, M. Szarek, D. L. Bhatt, V. A. Bittner, R. Diaz, J. M. Edelberg, et al. "Alirocumab and Cardiovascular Outcomes after Acute Coronary Syndrome." *NEW ENGLAND JOURNAL OF MEDICINE* 379, no. 22 (November 29, 2018): 2097–2107. <https://doi.org/10.1056/NEJMoa1801174>.