

Knowledge, attitude and practice analysis of Antibiotics use and misuse in tertiary care rural hospital patients

ABSTRACT:

INTRODUCTION: In last decades, the increase in bacterial resistance to against all major antibiotics is a growing worldwide problem and is as significant threat to global public health in the 21century.

OBJECTIVES-To assess the knowledge, attitude and practice of antibiotics use and misuse among patients of tertiary care rural hospital.

METHODS- this is cross-sectional analysis study ,was conducted at tertiary care teaching rural hospital in India between December2020 to April 2021. A self-administrated questionnaire was distributed to the patients in outpatient departments in hospital. The questionnaire consisted of 3 parts, first part consist of demographic variables, second and third part regarding knowledge, attitude and practice of patients towards antibiotic use and misuse. SPSS 21 was used for data analysis.

Results: Out of the 110 participants, 56(51.5%) were males and 54(48.5%) were females. 55(50%) patients gets antibiotics from friends, relative and pharmacist without medical doctor checkup.38(35%) patients start antibiotics as preventive medication for any infection,73 (66%) patients stopped medicine early if they feels better.52(47%) patient expect antibiotics from doctor for common cold. 42% patient obtained antibiotics without valid prescription from relatives, pharmacist or self prescription. In knowledge domain while majority of respondent aware that antibiotics is hazardous during pregnancy and lactation, can leads to severe allergic reactions. But very few respondent know about resistance can transfer from animal to human and human to human, also very few respondent know about antibiotics can alter human body bacterial flora.

Conclusion: There is lack of knowledge regarding antibiotics use and resistance among general population. Strict strategies need to employ like antibiotics prescription audit, antibiotics dispensing monitoring, effective public health education, and promoting health worker and patient communication.

Keywords: Antibiotics, Misuse, antibiotic resistance, health education, communication

Introduction:

In last decades, the increase in bacterial resistance to against all major antibiotics is a growing worldwide problem and is as significant threat to global public health in the 21century[1]. This spread of resistance directly affect not only therapeutic effectiveness of drugs but also increase in treatment failure episodes and costs and mortality rates among patients.[2]Improper and indiscriminate use of antibiotics are important reasons for antibiotic resistance in our country.[3–5]. Misuse of antibiotics occurs due to many reasons like physicians knowledge and experience, diagnostic uncertainty, while general public knowledge, practice, beliefs and attitudes towards antibiotic use, self-administration of antibiotics, patient's expectations from doctors, and patients experience with antibiotics, insufficient patient education by physicians, poor healthcare worker and patients communication are important key factors which directly influence antibiotics use and misuses [6–8]. There is a common preoccupation among common public that the use of antibiotics can help in early recovery and

sometimes prevention of further development of illness. There is a major gap in understanding about antibiotics use among general population, also very few studies in Indian rural population regarding antibiotics utilization(9,10) are available, hence current study was planned to assess knowledge, attitudes and practice regarding antibiotics misuse in rural population.

Materials and Methods:

Study design and population: This is cross-sectional analysis study was conducted at tertiary care rural hospital in India between December 2020 to April 2021 and consist of patients who are suffering from various illness and attended Medicine, ENT, Ophthalmology and General surgery departments. Patients aged 18 years or above were included in study. Well-structured English and local language (Marathi) questionnaires were made. The questionnaire was formed based on the questions taken from various previous studies and edited to suit the objectives.(24, 25,26).A literature review of past similar studies of antibiotics misuse was done to identify potential domains for the study instrument and changes made to suit the local population. A pilot study was also conducted to test the validity of the questionnaire for content, readability, design and comprehension on 20 local peoples and necessary modifications were made so that the questionnaire was simple for understanding and answering, still gave accurate data.

The final version of the questionnaire consist of 24 questions divided into four parts. The first part was about demographic variables of local populations. The second part consist of questions regarding knowledge and belief of participant regarding antibiotic use and antibiotic resistance with Yes, No and don't know options. Third part consist of questions regarding attitude of respondent towards antibiotics use consist of 5 statements with yes – no options. Fourth part consist of 4 multiple choice questions with suitable options.

A written consent was taken from all respondent before participation in study, Participants' confidentiality was assured by assigning each respondent a unique a code number for the purpose of analysis only. No any specific incentives or rewards were given to participants. Study was approved by the institutional review committee.

Data analysis was done using SPSS 21. The prevalence of antibiotic use, misuse and its awareness were determined by comparing outcome measures. Numerical variables were reported as percentage.

Results-Of the 130 questionnaires distributed, 110 participant (95.24%) completed questionnaires and were included in study, while the remaining 20 (4.76%) were excluded due to incomplete, invalid filling or double answers. Among 110 participants, 56 (51%) were males and 54 (49%) were females.

Respondents were compared according to their demographic parameters: age, gender, marital status, educational level, antibiotics usage. Among the respondents, 21(19%) were in the age group of 18-35, 46(42%) were in 35-50 years age group and 26(24%) were in 50 to 65 year age group, while 17(15%) are in more than 65 year age group. 83(75%) were married, 75(68%) were having education more than high school.

Table 1 - Demographic characteristics of respondent .

Characteristics		N	%
Gender	Male	56	51%
	Female	54	49%
Age in years	18-35	21	19%
	35-50	46	42%
	50-65	26	24%
	>65	17	15%
Education status	Uneducated	12	11%
	Less than high school	23	21%
	High school	39	35%
	Graduation	22	20%
	Postgraduation	14	13%
Marital status	Married	83	75%
	Non married	27	25%

Table 2 - Knowledge of respondents regarding antibiotics resistance and safety.

Sr no	Items		N	%
1	Respondents aware of the meaning of antibiotic resistance		44	40%
2	Awareness regarding reasons for antibiotic resistance			
	A) Using antibiotics when it is not needed		22	20%
	B) Not completing the full course of Antibiotics		14	13%
	C) I don't know		74	67%
3	Antibiotics are safe during pregnancy	Y	5	5%
		N	88	80%
		DK	17	15%
4	Antibiotics might develop allergy leading to death	Y	74	67%
		N	6	5%
		DK	30	27%
5	Are antibiotics are safe during breast feeding	Y	2	2%
		N	88	80%
		DK	20	18%
6	Excessive animal use of antibiotics can reduced its effect in	Y	23	21%

	human being	N	30	27%
		DK	57	52%
7	You stopped antibiotic course prematurely if symptoms are improving	Y	28	25%
		N	36	33%
		DK	46	42%
8	Antibiotics resistance can spread from human to human	Y	12	11%
		N	14	13%
		DK	84	76%
9	Antibiotics resistance can spread from animal to humans.	Y	6	5%
		N	14	13%
		DK	90	82%
10	Antibiotics can negatively alter our antibiotics flora	Y	16	15%
		N	12	11%
		DK	82	75%

Y-YES, N- NO, DK- DON'T KNOW

Table 3–Attitude domain questions towards antibiotics use among respondents.

sr no	ATTITUDE ITEMS	N	%
1	Keeping antibiotics at home as an emergency	48	44%
2	Taking antibiotics as preventive treatment before any symptoms	38	35%
3	Procured antibiotics from relative or friends with medical consultation	55	50%
4	Premature stopping antibiotics full course when feeling better	73	66%
5	Expect antibiotics from physician for common cold and every fever illness.	52	47%
6	Forcing doctor to prescribe antibiotics if not prescribed	15	14%

Table 4 – Practice domain question towards antibiotics use among respondents

SR NO	PRACTICE CHARACTERISTICS	N	%
1	Source of antibiotics		
	Doctors' prescription	64	58%
	Pharmacist	20	18%
	Relatives	12	11%
	Self-prescription	14	13%
2	<i>Reasons for use of antibiotics</i>		
	Fever	44	40%
	Cold	12	11%
	Cough	16	15%
	Sore throat	25	23%
	Toothache	37	34%
	Pain in any part of the body	31	28%
	Generally tired	12	11%
	Others	10	9%
3	<i>Reason for Stopping antibiotics</i>		
	After finishing the antibiotics course	40	36%
	When patient feel better	66	60%
4	I fully trust my doctor if he not prescribe me any antibiotics	55	50%

From assessment of knowledge domain it was found that only 44 % of respondent were aware about of antibiotic resistance, while 68% were not know reasons for antibiotic resistance, and 22% respondent thing about antibiotic resistance due to improper use and 14% know that prior stopping antibiotic could leads to antibiotic resistance. 88% were know about antibiotics are risky while pregnancy similarly 88% agree that antibiotic are not safe during breast feeding, 74% respondent were agree that antibiotics could cause allergic reactions. 57% respondents were unaware about antibiotics use in animal can cause human antibiotic resistance, in fact 30% denied that animal antibiotic use leads to human antibiotic resistance. Only 36% agree that one should not stopped taking antibiotics once patients start feeling better , while 28% responded disagree to it , while 46 % responded were not know relation between full course of antibiotics and antibiotics resistance. While only 12% respondents were aware that resistance can spread from human to human, and only 6% respondent aware that resistance can spread from animal to human, only 16% respondent aware about fact that antibiotics can cause negative effects on the body's own bacterial flora.

While analyzing attitude domain of questionnaire it was found that, 44% peoples are keeping antibiotics at home for any emergency use. While 35% respondent takes antibiotics as preventive treatment before any start of symptoms, 50% respondent procured antibiotics from friends and relatives without prior consultation with medical doctor. While 66% patient stop taking antibiotics before completing the course of treatment, 47% respondent expect antibiotics should be prescribed by

doctor for common cold symptoms and for every fever illness and 14% peoples force doctor to prescribe antibiotics if not prescribed.

68% respondent use antibiotics only after doctors prescription while 18% respondent use antibiotics on pharmacist advise, 11% on relatives advise and 13% on self prescription. 40% use antibiotics for fever, while 28% uses antibiotics for pain in any part of body and 11% use if they feel tired, 34% use for toothache like reason. 38% respondent stopped using antibiotics if they feel better, while 68% patient completed full course, 50% respondent trust doctor if he not prescribe antibiotics.

Discussion

According to WHO, Antibiotic resistance is a one of the top 10 global public health problem humanity is facing (1). Antibiotics resistance causing multiple problem any country. In addition to increase in mortality and morbidity, prolonged hospital stay, rise in demand of more expensive antibiotics and financial burden on individuals it causes significantly increase in financial burden on economy. (11,12) Prevalence of antibiotic resistance become major health problem in world specially in developing countries like India.(13). It is not difficult in India to obtain any antibiotic without prescription over the counter; hence problem of antibiotic resistance is increasing day by day. Major reasons for antibiotic resistance are misuse of antibiotics like indiscriminate and inadequate use of antibiotics (14), misuse of antibiotics in animals etc. (15). Resistance to ABs has been linked to levels of consumption (16,17) with evidence of a cause effect relationship (18). Our study is one of few attempts in literature to analyses the knowledge, beliefs, attitude and practice of antibiotics use by general public of Indian rural area.

On positive note we found that majority of population are agree that antibiotics hazardous for pregnancy and breastfeeding, also majority of population knows antibiotics can cause severe allergic reactions, which is comparable with other similar studies(19). While as much as about 40 % population aware of term antibiotic resistance considering rural area which interesting for us. Also majority of population are (58%) are still using antibiotic only after medical Doctors advice. This finding may be attributed to increase in educational level among population.

But still some major misbelief about antibiotic use and resistance are present among general population which leads to misuse of antibiotics. Around 47% population expect antibiotics for common cold, this is because there is lack of understanding that common cold is caused by viruses and not by bacteria and antibiotic are used against only for bacterial diseases. From this study nearly 40% population still believe that keeping antibiotics for emergency use is a good habit, while 50% population think that it is good to procured antibiotics from friends, relatives and friends and pharmacist without any proper medical consultation and doctor's prescription. Various past studies have confirmed that using leftover mediations results in poor patients compliance with antibiotics therapy (20) and this result in increases in antibiotics resistance.(20)

About 25% respondents disagree for need to complete full course of antibiotics and 66% respondent are stopped taking antibiotics if they feels better. Its very important to complete full course on antibiotics to restrict antibiotics resistance, According to the WHO, when a patient stops antibiotics course prematurely it results in promotes growth of Natural resistant strains and therefore it is advisable for patients to always complete full course of antibiotics prescribed tothem by a certified health professional.(21).Furthermore, patients need to educate regarding antibiotics resistance as 72% and 82% unaware about fact that antibiotics resistance can spread from person to person and animal to person, respectively. Also about 75% respondents are not aware that antibiotics can negatively

alter our own natural bacterial flora. This finding are correlates with various past studies studies.(19,24,25,26)

Subsequently, the general public needs to be educated on the fact that, the antibiotics efficacy will be preserved only when they are used only as per doctors guidelines and when the full course is completed [20,21]. Patients and healthcare workers communication can play important role to minimize antibiotics resistance. Healthcare worker can convey appropriate usage of antibiotics to general public through effective communication. This has proved in various studies also (22,-25). But in same study it is noted that healthcare workers and doctors not give enough time to communicate how to use antibiotics, this should be changed (26-28). A number of related studies were reported (29-33). Some of the key studies on antibiotics resistance and related issues were reviewed(34-38). On contradictory to other studies (26-28) , in our study level of education and misuse of antibiotics not relate each other significantly, as even respondent with higher education have deficit of basic antibiotics usage knowledge.

Conclusion :

Considering rapid growth of antibiotic resistance in India, and related gross effects due to antibiotics resistance, current study demonstrated important lacunae regarding antibiotic use in general public. There is lack knowledge and understanding regarding the safe use and consumption of antibiotics. There is need to established effective multifaceted intervention to improve appropriate use of antibiotics, which should include following: 1) need to audit antibiotic prescription from healthcare facility. 2) Effective public health education program using all media 3) strategies need to employ for promoting effective communication between healthcare workers specifically pharmacist and doctors with general public regarding safe and appropriate antibiotics usage.4) Need monitoring sources antibiotics through enforcing strict regulations.

References:

1. World Health Organization (2007) The world health report 2007. A safer future: global public health security in the 21st century. Available: <http://www.who.int/whr/2007>. Accessed 1 July 2021
2. World Health Organization (2001) WHO Global Strategy for Containment of Antimicrobial Resistance.WHO/CDS/CSR/DRS/2001.2. Available: http://www.who.int/csr/resources/publications/drugresist/en/EGlobal_Strat.pdf. Accessed 1 July 2021
3. Costelloe C, Metcalfe C, Lovering A, Mant D, Hay AD (2010) Effect of antibiotic prescribing in primary care on antimicrobial resistance in individual patients: systematic review and meta-analysis. *BMJ* 340: c2096. doi: 10.1136/bmj.c2096 PMID: 20483949
4. Vaananen MH, Pietila K, Airaksinen M (2006) Self-medication with antibiotics—does it really happen in Europe? *Health policy* 77: 166–171. PMID: 16095749
5. You JH, Yau B, Choi KC, Chau CT, Huang QR, et al. (2008) Public knowledge, attitudes and behavior on antibiotic use: a telephone survey in Hong Kong. *Infection* 36: 153–157. doi: 10.1007/s15010-0077214-51 PMID: 18231717.
6. Davey P, Pagliari C, Hayes A (2002) The patient's role in the spread and control of bacterial resistance to antibiotics. *ClinMicrobiol Infect* 8: 43–68. PMID: 12427207.

7. Franco BE, Altagracia Martinez M, Sanchez Rodriguez MA, Wertheimer AI (2009) The determinants of the antibiotic resistance process. *Infect Drug Resist* 2: 1–11. PMID: 21694883
8. Hulscher ME, van der Meer JW, Grol RP (2010) Antibiotic use: how to improve it? *Int J Med Microbiol* 300: 351–356. doi: 10.1016/j.ijmm.2010.04.003 PMID: 20434950
9. Purohit MR, Chandran S, Shah H, Diwan V, Tamhankar AJ, StålsbyLundborg C. Antibiotic Resistance in an Indian Rural Community: A 'One-Health' Observational Study on Commensal Coliform from Humans, Animals, and Water. *Int J Environ Res Public Health*. 2017 Apr 6;14(4):386. doi: 10.3390/ijerph14040386. PMID: 28383517; PMCID: PMC5409587.
10. Alvarez-Uria G, Zachariah S, Thomas D. High prescription of antimicrobials in a rural district hospital in India. *Pharm Pract (Granada)*. 2014 Apr;12(2):384. doi: 10.4321/s1886-36552014000200003. Epub 2014 Mar 15. PMID: 25035714; PMCID: PMC4100948.
11. Ott E, Bange FC, Reichardt C, Graf K, Eckstein M, Schwab F, et al. Costs of nosocomial pneumonia caused by methicillin-resistant *Staphylococcus aureus*. *J Hosp Infect*. 2010;76:300–3.
12. Raupach-Rosin H, Rübsamen N, Szkopek S, Schmalz O, Karch A, Mikolajczyk R, Castell S. Care for MRSA carriers in the outpatient sector: a survey among MRSA carriers and physicians in two regions in Germany. *BMC Infect Dis*. 2016;16(1):184.
13. Kumar SG, Adithan C, Harish BN, Sujatha S, Roy G, Malini A. Antimicrobial resistance in India: A review. *J Nat Sci Biol Med*. 2013 Jul;4(2):286-91. doi: 10.4103/0976-9668.116970. PMID: 24082718; PMCID: PMC3783766.
14. Tenover FC, McGowan JE Jr. Reasons for the emergence of antibiotic resistance. *Am J Med Sci*. 1996 Jan;311(1):9-16. doi: 10.1097/00000441-199601000-00003. PMID: 8571988
15. van den Bogaard AE, Stobberingh EE. Antibiotic usage in animals: impact on bacterial resistance and public health. *Drugs*. 1999 Oct;58(4):589-607. doi: 10.2165/00003495-199958040-00002. PMID: 10551432.
16. Bell BG, Schellevis F, Stobberingh E, Goossens H, Pringle M. A systematic review and meta-analysis of the effects of antibiotic consumption on antibiotic resistance. *BMC Infect Dis* 2014;14: 13.
17. Musher DM, Thorner AR. Community-acquired pneumonia. *New Eng J Med* 2014; 371: 1619-1628.
18. Bell BG, Schellevis F, Stobberingh E, Goossens H, Pringle M. A systematic review and meta-analysis of the effects of antibiotic consumption on antibiotic resistance. *BMC Infect Dis* 2014;14: 13.
19. Al-Shibani N, Hamed A, Labban N, Al-Kattan R, Al-Otaibi H, Alfadda S. Knowledge, attitude and practice of antibiotic use and misuse among adults in Riyadh, Saudi Arabia. *Saudi Med J*. 2017 Oct;38(10):1038-1044. doi: 10.15537/smj.2017.10.19887. PMID: 28917069; PMCID: PMC5694638.
20. Kardas P, Devine S, Golembesky A, Roberts C. A systematic review and meta-analysis of misuse of antibiotic therapies in the community. *Int J Antimicrob Agents*. 2005;26(2):106–13.
21. Ventola CL. The antibiotic resistance crisis: part 1: causes and threats. *P T*. 2015 Apr;40(4):277-83. PMID: 25859123; PMCID: PMC4378521.
22. Lee CR, Lee JH, Kang LW, Jeong BC, Lee SH. Educational effectiveness, target, and content for prudent antibiotic use. *Biomed Res Int*. 2015;2015:214021. doi: 10.1155/2015/214021. Epub 2015 Apr 5. PMID: 25945327; PMCID: PMC4402196.

23. Lee CR, Cho IH, Jeong BC, Lee SH. Strategies to minimize antibiotic resistance. *Int J Environ Res Public Health*. 2013 Sep 12;10(9):4274-305. doi: 10.3390/ijerph10094274. PMID: 24036486; PMCID: PMC3799537.
24. Effah CY, Amoah AN, Liu H, Agboyibor C, Miao L, Wang J, Wu Y. A population-base survey on knowledge, attitude and awareness of the general public on antibiotic use and resistance. *Antimicrob Resist Infect Control*. 2020 Jul 11;9(1):105. doi: 10.1186/s13756-020-00768-9. PMID: 32653034; PMCID: PMC7353772.
25. Awad AI, Aboud EA. Knowledge, attitude and practice towards antibiotic use among the public in Kuwait. *PLoS One*. 2015 Feb 12;10(2):e0117910. doi: 10.1371/journal.pone.0117910. PMID: 25675405; PMCID: PMC4326422.
26. Ling Oh A, Hassali MA, Al-Haddad MS, Syed Sulaiman SA, Shafie AA, Awaisu A. Public knowledge and attitudes towards antibiotic usage: a cross-sectional study among the general public in the state of Penang, Malaysia. *J Infect Dev Ctries*. 2011 May 28;5(5):338-47. doi: 10.3855/jidc.1502. PMID: 21628809.
27. Barker, A., Brown, K., Ahsan, M. *et al.* Social determinants of antibiotic misuse: a qualitative study of community members in Haryana, India. *BMC Public Health* **17**, 333 (2017). <https://doi.org/10.1186/s12889-017-4261-4>.
28. Ahmed SA, Kumar A, Sethi P, Kapil A, Pandey RM, Wig N. Effectiveness of education and antibiotic control programme at All India Institute of Medical Sciences, New Delhi. *Natl Med J India*. 2018 Sep-Oct;31(5):262-267. doi: 10.4103/0970-258X.261176. PMID: 31267989.
29. Sontakke, Yogesh A., and R. R. Fulzele. "Cytogenetic Study on Genotoxicity of Antitumor-Antibiotic Mitomycin C." *BIOMEDICAL RESEARCH-INDIA* 20, no. 1 (April 2009): 40–44.
30. Walia, Inderdeep Singh, Rajiv M. Borle, D. Mehendiratta, and Abhilasha O. Yadav. "Microbiology and Antibiotic Sensitivity of Head and Neck Space Infections of Odontogenic Origin." *JOURNAL OF MAXILLOFACIAL & ORAL SURGERY* 13, no. 1 (March 2014): 16–21. <https://doi.org/10.1007/s12663-012-0455-6>.
31. Aghi, Taru, Sanjiv Choudhary, Shilpi Basak, and Adarshlata Singh. "Clinico-Bacteriological Study of Pyoderma with Special Reference to Antibiotic Sensitivity in a Tertiary Care Rural Hospital." *RESEARCH JOURNAL OF PHARMACEUTICAL BIOLOGICAL AND CHEMICAL SCIENCES* 6, no. 2 (April 2015): 1701–7.
32. Pradhan, Swetalina, Bhushan Madke, Poonam Kabra, and Adarsh Lata Singh. "Anti-Inflammatory and Immunomodulatory Effects of Antibiotics and Their Use in Dermatology." *INDIAN JOURNAL OF DERMATOLOGY* 61, no. 5 (October 2016): 469–81. <https://doi.org/10.4103/0019-5154.190105>.
33. Dahake, Prasanna T., Sudhindra M. Baliga, Vijay M. Kumbar, and Kishore G. Bhat. "Cytotoxicity of Novel Polymeric Gel Matrix Triple Antibiotic Paste-an In Vitro Study." *REGENERATIVE ENGINEERING AND TRANSLATIONAL MEDICINE*, n.d. <https://doi.org/10.1007/s40883-020-00191-x>.
34. Rana, Deeksha, and Suvarna Sande. "Study of Prevalence and Antimicrobial Susceptibility Pattern of Enterococci Isolated from Clinically Relevant Samples with Special Reference to High Level Aminoglycoside Resistance (HLAR) in a Rural Tertiary Care Hospital." *JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS* 9, no. 34 (August 24, 2020): 2472–78. <https://doi.org/10.14260/jemds/2020/537>.
35. Dangre, G., N. Tankhiwale, and A. Mudey. "Antimicrobial Resistance in Neonatal Septicaemia: A Therapeutic Challenge to Pediatricians of Rural India." *INTERNATIONAL*

JOURNAL OF INFECTIOUS DISEASES 16, no. 1 (June 2012): E412.
<https://doi.org/10.1016/j.ijid.2012.05.564>.

36. Mallick, S. K., S. Basak, and S. Bose. "Inducible Clindamycin Resistance in Staphylococcus Aureus-A Therapeutic Challenge." JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH 3, no. 3 (June 2009): 1513–18.
37. Raut, U., S. Rantai, P. Narang, D. S. Chauhan, M. Chahar, R. Narang, and D. K. Mendiratta. "Comparison of the 3-(4,5-Dimethylthiazole-2-Yl)-2,5-Diphenyltetrazolium Bromide Tube Method with the Conventional Method and Real-Time Polymerase Chain Reaction for the Detection of Rifampicin Resistance in Mycobacterium Tuberculosis." INDIAN JOURNAL OF MEDICAL MICROBIOLOGY 30, no. 1 (March 2012): 81–84.
<https://doi.org/10.4103/0255-0857.93047>.
38. Sharma, S. K., J. Chaubey, B. K. Singh, R. Sharma, A. Mittel, and A. Sharma. "Drug Resistance Patterns among Extra-Pulmonary Tuberculosis Cases in a Tertiary Care Centre in North India." INTERNATIONAL JOURNAL OF TUBERCULOSIS AND LUNG DISEASE 21, no. 10 (October 2017): 1112–17. <https://doi.org/10.5588/ijtld.16.0939>.

UNDER PEER REVIEW