

# Original Research Article

## **Folic acid supplementation diminishes diabetes induced neural tube defects by recovering impaired embryo gene expression through its antioxidant activity**

### **Abstract**

**Aims:** This study aimed to determine whether folic acid supplementation could decrease the neural tube defects (NTDs) in embryos caused by hyperglycemia and the mechanism is associated with its antioxidative activity.

**Study design:** Diabetic pregnancy model were made and folic acid, Vitamin E and control were given to mice. Then the neural tube defects rate, oxidative stress markers and embryos size were measured and were analyzed.

**Place and Duration of Study:** Department of Development, College of Life Science, Northwest University, between Jan 2019 and Oct 2021.

**Methodology:** We injected folic acid to diabetic mice to score NTDs rate, measured embryos size, assayed ROS scavenging enzymes and Pax3 expression and oxidative stress marker including malondialdehyde and H<sub>2</sub>O<sub>2</sub> as well as the vitamin E controls.

**Results:** Injection of folic acid during gestation could diminish neural tube defect rate of diabetic mice and recover the Pax3 expression inhibition which is similar to vitamin E, but no effect was observed on nondiabetic mice. Folic acid supplementation also normalized the embryos size, decreased oxidative stress level including malondialdehyde and H<sub>2</sub>O<sub>2</sub> in diabetic mice and decreased the intracellular ROS of embryonic cells on day 8.5 induced by high glucose incubation in a dose-dependent manner. No effects were observed on ROS scavenging enzymes activity by folic acid supplementation.

**Conclusion:** Overall, the conclusion is that folic acid supplementation could diminish NTDs induced by hyperglycemia, the mechanism is associated with its antioxidant activity which can reduce the oxidative stress and recover the inhibition of Pax3 expression.

**Keywords:** Diabetic embryopathy, Embryo, Folic acid, Neural tube defect, Oxidative stress

### **Abbreviations**

MDA malondiadehyde

NTDs neural tube defects

### **1. Introduction**

Diabetic Embryopathy is a complication of diabetes in which the embryo of a diabetic pregnancy develops congenital malformations. The incidence rate of congenital malformations in diabetic pregnancy is 6-9% compared to 2-3% in normal pregnancy[1]. Among the various defects caused by the hyperglycemia, neural tube and heart are two most common malformations[2-3]. Neural tube defects are severe, irreversible malformations including anencephaly and spina bifida. The rate of diabetic malformation could be not brought to the nondiabetic rate even after the intensive insulin therapy. So it should be necessary to examine the exact molecular mechanism behind diabetic embryopathy which has not been fully understood. Pax3, a gene encodes a transcription factor is essential for neural tube development. Homozygous Splotch embryos (Sp/Sp) which carry loss of

functional Pax3 alleles develop the neural tube defects, neural crest, and skeletal muscle with 100% penetrance[4-7]. Previous studies have indicated that maternal diabetes induced by streptozotocin (STZ), or transient induction of hyperglycemia ( $\geq 13.9$  mM ) by glucose injection on embryonic day 7.5 leads to decreased expression of Pax3 on day 8.5 and increased incidence of neural tube defect (NTD), which can be recognized by day 10.5[8-9]. Studies suggested that diabetic pregnancy is associated with oxidative stress which caused by maternal diabetes has been demonstrated to be the reason of the inhibition of Pax3 expression and increases NTDs [10]. Moreover, administration of antioxidants such as vitamin E or Glutathione (GSH) ethyl ester can prevent hyperglycemia-induced developmental defects through the decreasing oxidative stress, recover the inhibition of Pax3[11]. Supplementary folic acid intake during human pregnancy can diminish fetal rate of defects [12-18] . However, the antiteratogenic mechanism of folic acid is still unclear. The studies results show that folate could increase the production of methionine from homocysteine, there by preventing the accumulation of homocysteine which is an oxidant[19-20]. In addition, folic acid has antioxidant properties of its own[21]. Thus it is possible that folic acid supplementation could diminish the neural tube defects caused by the hyperglycemia through reducing the oxidative stress in embryo, at least in part, diminish the inhibition of Pax3.

In this study, we tested the hypothesis that folic acid supplementation could block the inhibition of Pax3 expression and the NTDs caused by the hyperglycemia, and whether this is associated with its antioxidative activity to decrease oxidative stress.

## **2. Material and methods / experimental details / methodology**

### **2.1. Animal**

ICR mice used in the experiments were purchased from animal center of Xi'an Jiaotong University of China. All ICR mice were fed with commercial food and sterile water. Male and female ICR mice were mated and checked the plug next morning. The day which copulation plug was found is considered the gestational 0.5 day. Mice were killed to recover embryos for assay of markers of oxidative stress MDA and H<sub>2</sub>O<sub>2</sub> on day 7.5, for assay of genes expression on day 8.5, for score the NTD on day 10.5. We also measure the crown-rump length, biaparietal diameter, abdominal anteroposterior diameter and abdominal transverse diameter of embryos on day 10.5.

### **2.2. Induction of diabetes**

Diabetes was induced in 5 to 7 weeks old female ICR mice with 100 mg/Kg streptozotocin (STZ Sigma) dissolved in 10 mM sodium citrate (pH 4.5). Blood glucose was monitored daily with ACCU-CHEK (Roche). STZ-diabetic mice maintained euglycemia with insulin pellets before pregnancy but developed hyperglycemia beginning on day 4.5 of pregnancy, thereby exposing the embryo to hyperglycemia during the entire post implantation period. Nondiabetic mice were injected in parallel with PBS.

### ***2.3. Injection of folic acid***

Control and diabetic pregnant ICR mice were given 10 mg/Kg folic acid (Sigma) by daily subcutaneous injections. The injections commenced on gestational day 0.5 and continued up to termination of pregnancy on gestational day 7.5, 8.5 or 10.5. Controls were injected in parallel with PBS. Vitamin E-treated animals were supplemented with 0.5% (w/w) vitamin E succinate beginning on day 0.5 of pregnancy continued to the animals were killed.

### ***2.4. Preparation of total RNA and cDNA***

The total RNA from 50 embryos (n=5) of each group were extracted with a total RNA isolation kit TRIzol reagent (Invitrogen) according to the manufacture's instruction. The reaction mixture containing 2 µg of RNA, 2.5 µM random primer, 200 U of molony murine leukemia virus reverse transcriptase (MMLV, Promega), 2 mM of each dNTPs, 5 U of RNasin in a total volume of 25 µl was incubated for 1 h at 42°C to synthesize cDNA.

### ***2.5. Analysis of SOD, CAT, GPX and Pax3 mRNA levels***

Specific primers encoding CuZnSOD, MnSOD, CAT, GPX and Pax3 were used (Table 1). Each 25 µl of Realtime-PCR reaction (Takara) volume included 25 ng of cDNA 2µl, 12.5 µl of 2× SYBR Green Premix Ex Taq buffer, 1µl forward and reverse primer and 8.5 µl H<sub>2</sub>O. PCR cycle parameters

were 95 °C for 30 s followed by 40 cycles at 95 °C for 5 s and 60 °C for 30 s.

### ***2.6. Preparation of embryonic cell suspensions and assessment of intracellular ROS production***

Embryos on day 8.5 of gestation were dissected from uteri and extraembryonic structure in PBS and placed in DMEM medium. Whole embryos were minced then by passing through cell strainer with 40 $\mu$ M nylon mesh (BD Falcon TM), to generate single cell suspensions. The suspension was prepared at a density of  $2 \times 10^7$  cells/ml. The embryonic cells were cultured in DMEM medium under different conditions for 24 hours: 0 mmol/L glucose medium, 30mmol/L glucose medium, 30mmol/L with 1 mmol/L and 3 mmol/L folic acid. Cells were then washed twice with PBS. Intracellular ROS production was measured with 2',7'-dichlorodihydrofluorescein diacetate (DCF-DA) (Molecular Probes). Embryonic cells were incubated with DCF-DA (5  $\mu$ mol/L) in serum-free medium at 37 °C for 30 minutes and then washed with PBS. DCF fluorescence was excited at 488 nm, and emission at 530nm was measured on Gemini EM Reader (Molecular Devices). Fluorescence intensity values were presented as the percentage of the control value, after subtraction of background fluorescence (Fig 3).

### ***2.7. Assay of Malondialdehyde and H<sub>2</sub>O<sub>2</sub>***

Embryos were recovered on day 7.5 for the assay of malondialdehyde (MDA) and H<sub>2</sub>O<sub>2</sub>. MDA was assayed spectrophotometrically using the thiobarbituric acid (TBA) test, taking purified MDA (Sigma) to generate a standard curve. H<sub>2</sub>O<sub>2</sub> was assayed using kits of Cayman Chemicals according to the instructions. MDA and H<sub>2</sub>O<sub>2</sub> were normalized to protein concentrations in cell extracts by using Bio-Rad protein reagent (Bio-Rad).

### ***2.8. Statistical Analysis***

Data were analyzed by SPSS software. The results are expressed as mean  $\pm$  SD. Student's t-test was used to compare difference between two groups. A p value less than 0.05 was considered to be statistically significant.

### 3 Results

#### *3.1. Folic acid diminished the inhibition of Pax3 expression and NTD induced by hyperglycemia in vivo*

Diabetic mice display serious NTD malformation. Among the defects of diabetic mice, 53.6% of defects are single exencephaly located at forebrain, midbrain or hindbrain. 21.4% defects are combined exencephaly. 25% defects are spina bifida. Subcutaneous injection of folic acid during gestation could diminish the NTD rate of diabetic mice. Injection of folic acid lowered the NTD rate from 44% to 17% similar to vitamin E supplementation which decreased to 16% (Fig 1A). The malformation in folic acid injection group have similar formation to diabetic group( 55.6% for single exencephaly, 22.2% for combined exencephaly and 22.2% for spina bifida) (Table 2). Injection of folic

acid could recover the Pax3 inhibition by high glucose and it has similar effect to vitamin E which had been proved to prevent the NTD from diabetic mice through the decreasing of oxidative stress induced by hyperglycemia (Fig 1B). For both NTD incidence and Pax3 expression level, folic acid injection has no effects on nondiabetic mice. There is no effect for folic acid to change the mice glucose level (Table 3 and Table 4).

### ***3.2. Folic acid could normalize the fetal size in diabetic mice***

We measured the size of embryos day 10.5 including crown-rump length (CRL), biaparietal diameter (BPD), abdominal anteroposterior diameter (APD) and abdominal transverse diameter (ATD) under different treatment conditions. We found the decreased size including CRL, BPD, APD and ATD in diabetic mice embryos compared with nondiabetic mice ( $p < 0.01$ ). Injection of folic acid could normalize the decreased size in diabetic mice significantly compared with the diabetic mice ( $p < 0.01$ ). There are still difference about the CRL, BPD and APD between folic acid group and nondiabetic mice. But there is no difference on ATD of D/FA compared with ND ATD (Table 5). There is also no difference on somite number between each group.

### ***3.3. Folic acid decreased the intracellular ROS caused by hyperglycemia in embryonic cells In situ***

ROS production of embryonic cells on day 8.5 were measured by DCF fluorescence, there was significantly increased ROS after 24 hours incubation with 30 mmol/L glucose medium. Embryonic cells on day 8.5 cultured in 30 mmol/L glucose medium increased intracellular ROS production by  $142 \pm 11.2\%$  compared with control which is 0 mmol/L glucose medium (100%;  $p < 0.01$ ;  $n = 6$ ). Addition of 1mmol/L and 3 mmol/L folic acid to 30 mmol/L glucose medium could decrease the ROS production compared with 0 mmol/L folic acid in a dose-dependent manner (decreased from  $142 \pm 11.2\%$  to  $130 \pm 7.84\%$  and  $118 \pm 6.68\%$  versus control 100%  $n=6$  Fig 2 ).

### ***3.4. Folic acid reduced the oxidative stress marker level***

To further test whether the effect of reducing neural tube defect and inhibition of Pax3 expression by the supplementation of folic acid is the result of blocking of oxidative stress induced by the hyperglycemia, we assayed the oxidative stress markers of MDA and H<sub>2</sub>O<sub>2</sub> with and without the supplementation of folic acid. As shown in Fig 4, MDA, a marker of lipid peroxidation, was significantly increased by hyperglycemia. H<sub>2</sub>O<sub>2</sub> was also increased. On the other hand, there was no effect of folic acid on MDA in nondiabetic pregnancies, but the increase in MDA observed in diabetic mice was significantly reduced by folic acid, although it was still significantly increased over MDA in nondiabetic pregnancies. Similarly, there was no effect of folic acid on H<sub>2</sub>O<sub>2</sub> of nondiabetic pregnancies. The increase in H<sub>2</sub>O<sub>2</sub> observed in diabetic mice was significantly suppressed by supplementation of folic acid, although it remained significantly increased compared with nondiabetic mice (Fig 3).

### ***3.5. Folic acid didn't affect the ROS scavenging enzymes activity in diabetic embryos***

We measured the ROS scavenging enzymes activity in diabetic embryos, diabetes didn't affect embryos ROS scavenging enzymes on day 8.5, we didn't find any significant effect of folic acid on CuZnSOD, MnSOD, CAT, GPX expression level on day 8.5. There are no significant difference between the diabetic embryos and normal embryos on enzymes activity ( $p > 0.05$ ). There is no difference in diabetic embryos with or without folic acid treatment ( $p > 0.05$ ) (Fig 4).

#### 4 Discussion

To investigate whether folic acid could prevent the NTD caused by the hyperglycemia and its molecular mechanism, we treated diabetic mice model with folic acid in vivo. The data reported here support the hypothesis that folic acid could diminish the NTD induced by high glucose and the mechanism is associated with its antioxidant activity which can decrease the oxidative stress and reduce the inhibition of Pax3 induced by hyperglycemia.

Folic acid supplementation could reduce the incidence of NTDs in many case, it has been applied as an effective preventive teratogenic method for many years[17]. Many studies have shown that 50-70% of NTDs can be prevented by folic acid supplementation before and during pregnancy[22-23]. The question is how about the preventive effect of folic acid against NTDs in diabetic pregnancies which has a higher NTDs incidence rate compared with nondiabetic pregnancy. Folic acid injection to diabetic mice indicates that folic acid supplementation could reduce the NTDs caused by hyperglycemia (Fig 1A, Fig5). Folic acid supplementation not only can diminish the malformation rate in diabetic mice but also can normalize the smaller size of embryos caused by high glucose (Table 5). So folic acid addition could defeat glucose effect on embryo development. But the molecular mechanism is still unclear. Some studies show that addition of extra folic acid corrects subtle alterations in the folic acid metabolism, but folic acid could not reduce the NTD rate to zero no matter in clinic or in the animal model[24-25]. At the same time, maternal folate levels in most affected pregnancies are within the normal range[26]. The abnormal folate metabolism is another point involved in the mechanism of folic acid against malformation. But there is no correlation between folic acid levels in the mother and incidence of NTDs in infants and no detectable abnormal metabolism of folate in diabetic pregnancy women compared with normal pregnant women[27-30]. The study using

the rat diabetic pregnancy model and embryos culturing shows that diabetic environment induces a state of functional folic acid deficiency in the embryo. This deficiency is only partly a transport and concentration deficiency, as the high glucose cultured embryos did not show decreased levels of folic acid despite the presence of developmental malformation [11]. So it seems there must be some other folic acid beneficial activity involved in its supplementation effect.

More and more evidences suggest that the gene-environment interactions are aetiology of NTDs. There are more than 100 single-gene defects could cause NTDs in mouse. Folate status is a key environmental factor. Genes whose expression is altered in diabetes-exposed embryos might represent excellent candidates for folate-responsive genes, and may mediate the beneficial effect of folate in the prevention of neural tube and other developmental defects [31-33]. Study shows that folate deficiency does not cause NTDs in wild-type mice, but caused a significant increase in cranial NTDs among  $Sp^{2H}$  embryos which carry a mutation in Pax3, which is crucial for the neural tube closure[26]. So it appears reasonable to conclude that Pax3 which was inhibited by the hyperglycemia might be a target for folic acid effect. In vivo experiments shows that subcutaneous injection of folic acid to the STZ treated diabetic mice recovered partly the Pax3 expression level compared with normal mice embryos. We also observed that the folic acid injection did not alter the malformation rate and Pax3 expression in the control mice (Fig 1 B). It seems that the folic acid will not affect the Pax3 expression in the normal condition, but it can correct the Pax3 expression in high glucose. From the data provided here it appears reasonable to conclude that the folic acid did not take effect on Pax3 directly. The effect must be taken by other mechanism.

Numerous studies have shown that oxidative stress plays a very important role in the development defects such as NTD caused by diabetic pregnancy[34-40]. Oxidative stress causes

diabetic pregnancy-induced NTD because of its disrupting expression of Pax3. Our previous results have shown that hypoxic stress in diabetic contributes to impaired Pax3 expression and defective development by inducing oxidative stress. The antioxidants such as Vitamin E could prevent the maternal diabetes defects by the recovering of inhibition of Pax3 decrease the oxidative stress in experimental animals[11]. In considering mechanisms by which folic acid might recover Pax3 expression and decrease NTD rate, we consider several mechanisms especially the antioxidant activity of folic acid. Although the studies results shown that folate could increase the production of methionine from homocysteine, there by preventing the accumulation of homocysteine which is an oxidant[41-44], another study result shown that the preventive effect of folic acid is independent from homocysteine metabolism [45]. So we looked forward to the own antioxidant properties of folic acid. To explore the effect of folic acid on the oxidative stress, we measured intracellular ROS in embryonic cells cultured in high glucose with or without folic acid firstly. Folic acid could decrease the increased ROS level caused by hyperglycemia and importantly it is in a dose-dependent manner (Fig 2). Some other oxidative stress markers were also measured. The data shows that the folic acid injection could decrease the MDA and H<sub>2</sub>O<sub>2</sub> in the diabetic embryos (Fig 3). The effect was similar to vitamin E which was a positive control. Combined with the findings that folic acid supplementation did not protect the fetuses from neural tube defects in splotch mice which with a defect Pax3 function and a high incidence of NTDs [46], it is further reconfirm that the mechanism which the folic acid recover the Pax3 expression and decrease NTD by at least partly decrease the oxidative stress. We also measure the ROS scavenging enzymes activity of embryos in diabetic and nondiabetic mice with or without folic acid treatment, no changes were found. So it seems that the antioxidant activity of folic acid in embryos doesn't come from the changing of scavenging enzymes activity.

## 5 Conclusion

In conclusion, folic acid could reduce the inhibition of Pax3 expression in diabetic pregnancy by decrease the oxidative stress and recover the NTDs. Is it a general model of folate-responsive genes whose expression is altered in diabetes-exposed embryos or specific for the Pax3 gene? On the other hand, the mechanism of the folic acid preventive effect to malformations for the normal pregnancy is the same as diabetic pregnancy or not? It needs further study to answer them.

## Ethical approval

All authors hereby declare that all experiments have been examined and approved by the ethics committee of Northwest University

## COMPETING INTERESTS DISCLAIMER:

**Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.**

## References

1Loeken MR. Current perspectives on the causes of neural tube defects resulting from diabetic pregnancy. American journal of medical genetics Part C, Seminars in medical

- genetics. 2005; 135C:77-87.
- 2 Eriksson UJ. Congenital anomalies in diabetic pregnancy. *Seminars in fetal & neonatal medicine*. 2009; 14:85-93.
- 3 Zabihi S, Loeken MR. Understanding diabetic teratogenesis: where are we now and where are we going? *Birth defects research Part A, Clinical and molecular teratology*. 2010; 88:779-790.
- 4 Epstein DJ, Vogan KJ, Trasler DG, Gros P. A mutation within intron 3 of the Pax-3 gene produces aberrantly spliced mRNA transcripts in the splotch (Sp) mouse mutant. *Proceedings of the National Academy of Sciences of the United States of America*. 1993; 90:532-536.
- 5 Copp AJ, Greene ND. Genetics and development of neural tube defects. *The Journal of Pathology*. 2010; 220:217-230.
- 6 Maczkowiak F, Mateos S, Wang E, Roche D, Harland R, Monsoro-Burq AH. The Pax3 and Pax7 paralogs cooperate in neural and neural crest patterning using distinct molecular mechanisms, in *Xenopus laevis* embryos. *Developmental biology*. 2010;340:381-396.
- 7 Nelms BL, Pfaltzgraff ER, Labosky PA. Functional interaction between Foxd3 and Pax3 in cardiac neural crest development. *Genesis*. 2011; 49:10-23.
- 8 Phelan SA, Ito M, Loeken MR. Neural tube defects in embryos of diabetic mice: role of the Pax-3 gene and apoptosis. *Diabetes*. 1997;46:1189-1197.
- 9 Fine EL, Horal M, Chang TI, Fortin G, Loeken MR. Evidence that elevated glucose causes altered gene expression, apoptosis, and neural tube defects in a mouse model of diabetic pregnancy. *Diabetes*. 1999;48:2454-2462.
- 10 Chang TI, Horal M, Jain SK, Wang F, Patel R, Loeken MR. Oxidant regulation of gene

- expression and neural tube development: Insights gained from diabetic pregnancy on molecular causes of neural tube defects. *Diabetologia*. 2003;46:538-545.
- 11 Li R, Chase M, Jung SK, Smith PJ, Loeken MR. Hypoxic stress in diabetic pregnancy contributes to impaired embryo gene expression and defective development by inducing oxidative stress. *American journal of physiology Endocrinology and metabolism*. 2005; 289:E591-599.
- 12 Hall JG. Folic acid: the opportunity that still exists; [comment]. *CMAJ : Canadian Medical Association journal-journal de l'Association medicale canadienne*.2000;162:1571-1572.
- 13 Botto LD. International retrospective cohort study of neural tube defects in relation to folic acid recommendations: are the recommendations working? *BMJ*. 2005; 330:571.
- 14 Cantfield MA, Collins JS, Botto LD, Williams LJ, Mai CT, Kirby RS, Pearson K, Devine O, Mulinare J, National Birth Defects Prevention N. Changes in the birth prevalence of selected birth defects after grain fortification with folic acid in the United States: findings from a multi-state population-based study. *Birth defects research Part A, Clinical and molecular teratology*. 2005;73:679-689.
- 15 Robbins JM, Cleves MA, Collins HB, Andrews N, Smith LN, Hobbs CA. Randomized trial of a physician-based intervention to increase the use of folic acid supplements among women. *American journal of obstetrics and gynecology* 2005;192:1126-1132
- 16 Ryan-Harshman M, Aldoori W. Folic acid and prevention of neural tube defects. *Canadian family physician Medecin de famille canadien*. 2008; 54:36-38.
- 17 Molloy AM, Kirke PN, Troendle JF, Burke H, Sutton M, Brody LC, Scott JM, Mills JL. Maternal vitamin B12 status and risk of neural tube defects in a population with high neural tube defect prevalence and no folic Acid fortification. *Pediatrics*. 2009;123:917-923.

- 18 Mosley BS, Cleves MA, Siega-Riz AM, Shaw GM, Canfield MA, Waller DK, Werler MM, Hobbs CA, National Birth Defects Prevention S. Neural tube defects and maternal folate intake among pregnancies conceived after folic acid fortification in the United States. *American journal of epidemiology*. 2009; 169:9-17.
- 19 Rosenquist TH, Ratashak SA, Selhub J. Homocysteine induces congenital defects of the heart and neural tube: effect of folic acid. *Proceedings of the National Academy of Sciences of the United States of America*. 1996; 93:15227-15232.
- 20 Koz ST, Gouwy NT, Demir N, Nedzvetsky VS, Etem E, Baydas G (2010) Effects of maternal hyperhomocysteinemia induced by methionine intake on oxidative stress and apoptosis in pup rat brain. *International journal of developmental neuroscience : the official journal of the International Society for Developmental Neuroscience*. 2010;28:325-329.
- 21 Nakano E, Higgins JA, Powers HJ. Folate protects against oxidative modification of human LDL. *The British journal of nutrition*. 2001;86:637-639.
- 22 Czeizel AE, Toth M, Rockenbauer M. Population-based case control study of folic acid supplementation during pregnancy. *Teratology*. 1996;53:345-351.
- 23 Mills JL, Signore C. Neural tube defect rates before and after food fortification with folic acid. *Birth defects research Part A, Clinical and molecular teratology*. 2004;70:844-845.
- 24 Lumley J, Watson L, Watson M, Bower C. Periconceptional supplementation with folate and/or multivitamins for preventing neural tube defects. *The Cochrane database of Systematic reviews*:CD001056 2001
- 25 Wald NJ, Law MR, Morris JK, Wald DS. Quantifying the effect of folic acid. *Lancet*.

2001;358:2069-2073.

26 Burren KA, Savery D, Massa V, Kok RM, Scott JM, Blom HJ, Copp AJ, Greene ND.

Gene-environment interactions in the causation of neural tube defects: folate deficiency increases susceptibility conferred by loss of Pax3 function. *Human molecular genetics*. 2008;17:3675-3685.

27 Kirke PN, Molloy AM, Daly LE, Burke H, Weir DG, Scott JM. Maternal plasma folate and vitamin B12 are independent risk factors for neural tube defects. *The Quarterly journal of Medicine*. 1993; 86:703-708.

28 Mooij PN, Steegers-Theunissen RP, Thomas CM, Doesburg WH, Eskes TK.

Periconceptional vitamin profiles are not suitable for identifying women at risk for neural tube defects. *The Journal of nutrition*. 1993;123:197-203.

29 Kaplan JS, Iqbal S, England BG, Zawacki CM, Herman WH. Is pregnancy in diabetic women associated with folate deficiency? *Diabetes care*. 1999; 22:1017-1021.

30 Scott JM. Folate and vitamin B12. *The Proceedings of the Nutrition Society*. 1999; 58:441-448.

31 Spiegelstein O, Cabrera RM, Bozinov D, Wlodarczyk B, Finnell RH. Folate-regulated changes in gene expression in the anterior neural tube of folate binding protein-1 (Folbp1)-deficient murine embryos. *Neurochemical research*. 2004; 29:1105-1112.

32 Pickett EA, Olsen GS, Tallquist MD. Disruption of PDGFR $\alpha$ -initiated PI3K activation and migration of somite derivatives leads to spina bifida. *Development*. 2008; 135:589-598.

33 Pavlinkova G, Salbaum JM, Kappen C. Maternal diabetes alters transcriptional programs in the developing embryo. *BMC genomics*. 2009; 10:274.

34 Eriksson UJ, Borg LA. Protection by free oxygen radical scavenging enzymes against

- glucose-induced embryonic malformations in vitro. *Diabetologia*. 1991;34:325-331.
- 35 Cederberg J, Eriksson UJ. Decreased catalase activity in malformation-prone embryos of diabetic rats. *Teratology*. 1997;56:350-357.
- 36 Sivan E, Lee YC, Wu YK, Reece EA. Free radical scavenging enzymes in fetal dysmorphogenesis among offspring of diabetic rats. *Teratology*. 1997; 56:343-349.
- 37 Eriksson UJ, Wentzel P, Minhas HS, Thornalley PJ. Teratogenicity of 3-deoxyglucosone and diabetic embryopathy. *Diabetes*. 1998;47:1960-1966.
- 38 Ornoy A, Zaken V, Kohen R. Role of reactive oxygen species (ROS) in the diabetes-induced anomalies in rat embryos in vitro: reduction in antioxidant enzymes and low-molecular-weight antioxidants (LMWA) may be the causative factor for increased anomalies. *Teratology*. 1999;60:376-386.
- 39 Cederberg J, Basu S, Eriksson UJ. Increased rate of lipid peroxidation and protein carbonylation in experimental diabetic pregnancy. *Diabetologia*. 2001; 44:766-774.
- 40 Wentzel P, Gareskog M, Eriksson UJ. Decreased cardiac glutathione peroxidase levels and enhanced mandibular apoptosis in malformed embryos of diabetic rats. *Diabetes*. 2008;57:3344-3352.
- 41 Xu D, Neville R, Finkel T. Homocysteine accelerates endothelial cell senescence. *FEBS letters* 2000; 470:20-24.
- 42 Chern CL, Huang RF, Chen YH, Cheng JT, Liu TZ. Folate deficiency-induced oxidative stress and apoptosis are mediated via homocysteine-dependent overproduction of hydrogen peroxide and enhanced activation of NF-kappaB in human Hep G2 cells. *Biomedicine & pharmacotherapy*. 2001; 55:434-442.

- 43 Green NS. Folic acid supplementation and prevention of birth defects. *The Journal of nutrition*. 2002;132:2356S-2360S.
- 44 Ryan-Harshman M, Aldoori W. Folic acid and prevention of neural tube defects. *Canadian family physician Medecin de famille canadien*. 2008; 54:36-38.
- 45 Oyama K, Sugimura Y, Murase T, Uchida A, Hayasaka S, Oiso Y, Murata Y. Folic acid prevents congenital malformations in the offspring of diabetic mice. *Endocrine journal*. 2009;56:29-37.
- 46 Gefrides LA, Bennett GD, Finnell RH. Effects of folate supplementation on the risk of spontaneous and induced neural tube defects in Splotch mice. *Teratology*. 2002;65:63-69.

Table 1 Nucleotide sequences of the primers used in the study

---

Primer name	Sequence (5'-3')
-------------	------------------

---

---

CuZnSOD forward <sup>[23]</sup>	AAGGCCGTGTGCGTGCTGAA
CuZnSOD reverse <sup>[23]</sup>	CAGGTCTCCAACATGCCTCT
MnSOD forward <sup>[23]</sup>	GCACATTAACGCGCAGTCA
MnSOD reverse <sup>[23]</sup>	AGCCTCCAGCAACTCTCCTT
CAT forward <sup>[23]</sup>	GCAGATACCTGTGAACTGTC
CAT reverse <sup>[23]</sup>	GTAGAATGTCCGCACCTGAG
GPX forward <sup>[23]</sup>	CCTTCAAGTACGTCCGACCTG
GPX reverse <sup>[23]</sup>	CAATGTCGTTGCGGCACACC
Pax3 forward <sup>[24]</sup>	CCA ACC ATA TCC GCC ACAA
Pax3 reverse <sup>[24]</sup>	TCT TAG AGA CGC AAC CAT GGG
GAPDH forward <sup>[24]</sup>	TGTGTCCGTCGTGGATCTGA
GAPDH reverse <sup>[24]</sup>	CCTGCTTCACCACCTTCT TGA

---

Table 2 Morphology of day 10.5 mice embryos

---

N	Normal	Exencephaly	Exencephaly	Spina bifida
---	--------	-------------	-------------	--------------

---

			(single)	(combined)	
ND	58	55	3	0	0
N/FA	56	53	3	0	0
D	62	34	15	6	7
D/FA	54	45	5	2	2
D/VitE	56	46	7	1	2

Morphology of day 10.5 embryos from normal (ND) and diabetic (D) mice, some of which were supplemented with folic acid (N/FA and D/FA) and vitamin E (D/VitE). Exencephaly (single): only one malformation located on forebrain, midbrain or hindbrain. Exencephaly (combined): more than two malformation in one embryo.

Table 3 Mean blood glucose levels of pregnancies on day 7.5 in Fig 1A

ND	ND/FA	ND/Vit E	D	D/FA	D/Vit E
----	-------	----------	---	------	---------

glucose, mM	7.1±0.07	6.8±0.03	7.3±0.08	21.5±0.22	23.4±0.19	20.2±0.18
-------------	----------	----------	----------	-----------	-----------	-----------

Value are mean ± SE. All D p < 0.001 vs ND n=6 ND: non diabetic pregnancy D: diabetic pregnancy

D/FA: diabetic pregnancy giving folic acid D/Vit E: diabetic pregnancy giving vitamin E ND/FA: non

diabetic pregnancy giving folic acid ND/Vit E: non diabetic pregnancy giving vitamin E

UNDER PEER REVIEW

Table 4 Mean blood glucose levels of pregnancies in Fig 2B.

ND	ND/FA	ND/Vit E	D	D/FA	D/Vit E
----	-------	----------	---	------	---------

---

glucose, mM	6.8±0.04	7.2±0.05	7.1±0.09	20.1±0.12	22.7±0.15	19.8±0.12
-------------	----------	----------	----------	-----------	-----------	-----------

---

Value are mean  $\pm$  SE. All D  $p < 0.001$  vs ND n=6 ND: non diabetic pregnancy D: diabetic pregnancy

D/FA: diabetic pregnancy giving folic acid D/Vit E: diabetic pregnancy giving vitamin E ND/FA: non

diabetic pregnancy giving folic acid ND/Vit E: non diabetic pregnancy giving vitamin E

UNDER PEER REVIEW

Table 5 Embryo size on gestational day 10.5

---

N	Somite number	CRL( $\mu$ m)	BPD( $\mu$ m)	APD( $\mu$ m)	ATD( $\mu$ m)
---	---------------	---------------	---------------	---------------	---------------

---

ND	58	36.2 ± 0.6	5038±258	2022±188	2366±148	1397±178
D	62	35.6 ± 0.5	4074±249	1494±172	1852±130	1207±127
D/FA	54	34.9 ± 0.8	4691±242*	1865±152**	1987±87*	1368±129***

Value are mean ± SE. All D p < 0.01 vs ND, All D/FA p < 0.01 vs D, For D/FA, \* p < 0.01 vs ND, \*\* p < 0.05 vs ND, \*\*\* p > 0.05 vs ND. CRL: crown-rump length BPD: biaparietal diameter APD: abdominal anteroposterior diameter ATD: abdominal transverse diameter ND: non diabetic pregnancy D: diabetic pregnancy D/FA: diabetic pregnancy giving folic acid

UNDER PEER REVIEW





Fig 1 Effects of folic acid or vitamin E on Pax3 expression or NTD induced by maternal hyperglycemia. ICR mice were induced diabetic mice (D) and nondiabetic mice (ND), they were given folic acid injection or vitamin E supplementation as described in MATERIALS AND METHODS. A: \*  $p < 0.01$  vs. ND; \*\*  $p < 0.05$  vs. all other treatment groups. B: \*  $p < 0.05$  vs. all other treatment groups; \*\*  $p < 0.05$  vs. all other treatment groups. n=6 ND: non diabetic pregnancy D: diabetic pregnancy ND/FA: non diabetic pregnancy giving folic acid D/FA: diabetic pregnancy giving folic acid ND/Vit E: diabetic pregnancy giving vitamin E




Fig 2 Effects of folic acid on the decreased ROS production on day 8.5 embryonic cells caused by hyperglycemia. Embryonic cells on day 8.5 were cultured in 30 mmol/L DMEM medium for 24 hours shows increased ROS production ( 142% vs 100%, \*  $p < 0.01$  n=6). Addition of 1 mmol/L and 3 mmol/L folic acid decreased the ROS significantly (G30F1 vs G30F0 \*\*  $p < 0.05$ ; G30F3 vs G30F0 \*\*\*  $p < 0.01$ ; G30F3 vs G30F1 \*\*\*  $p < 0.05$ ; G30F3 vs G0F0 \*\*\*  $p > 0.05$ ; n=5). G0F0: 0 mmol/L glucose medium with no folic acid G30F0: 30 mmol/L glucose medium with no folic acid G30F1: 30 mmol/L glucose medium with 1 mmol/L folic acid G30F3: 30 mmol/L glucose medium with 3 mmol/L folic acid







Fig 3 MDA and H<sub>2</sub>O<sub>2</sub> in diabetic mice and nondiabetic mice under different treatments. A: MDA \* p < 0.001 vs. ND; \*\* p < 0.05 vs. all other treatment groups. B: H<sub>2</sub>O<sub>2</sub> \* p < 0.01 vs. all other treatment groups; \*\* p < 0.05 vs. ND. n=6






Fig 4 Effects of folic acid on ROS scavenging enzymes activity expression in diabetic and nondiabetic mice. ICR mice were induced diabetic mice (D) and nondiabetic mice (ND)DS. There are not statistically difference ( $p > 0.05$   $n = 6$ ) between every two groups. ND: non diabetic pregnancy D: diabetic pregnancy ND/FA: non diabetic pregnancy giving folic acid D/FA: diabetic pregnancy giving folic acid.