

Review Article

Antibiotic resistance in *Clostridium difficile*: A rapidly evolving threat

ABSTRACT:

The bacteria, *Clostridium difficile* the main pathogen associated with nosocomial infections is a gram positive, anaerobic and forms spores. The host and pathogen reaction results in a wide array of manifestation of the disease which ranges at one end as asymptomatic carriage to the other end as severe; toxic megacolon and thus making the treatment with antibiotic a challenge. To add upon the *Clostridium difficile* shows antibiotic resistance making its eradication almost impossible. This antibiotic resistance of *Clostridium difficile* has made it a global challenge over the decades, making it impossible for us to come to a conventional antibiotic to treat infections by *Clostridium difficile*. Several new strains of *Clostridium difficile* have emerged following the unscrupulous use of antimicrobial agents and subsequent transfer of resistance-causing genes between virulent strains the subsequent generations. The resistance to antibiotics is still rampant and not in control, rather its on rise as the bacteria has greater access to new host due to its mobility and very infectious nature. With the use of extended antimicrobial therapy, an acute danger of creating and spreading new resistant and multi-drug resistant strains always looms. *Clostridium difficile*, during its life cycle, produces spores which give the organism the capacity to resist extremes of change in the environment. In a country such as India, with a high population density and thus forming a high potential platform for the spread of multidrug-resistant strains of *Clostridium difficile*. An understanding of the emergence of antibiotic resistance in *Clostridium difficile*, the mechanisms by which the *Clostridium difficile* acquires it, and sensitivity to antibiotics becomes very important.

Conclusion: Therefore, we present a review of the antibiotic resistance seen in *Clostridium difficile*. RT027 has emerged as the most virulent strain all the strain of *Clostridium difficile*. Drug resistance is via multiple pathways, including *erm*, *cfr*, *tet*, and *rpo B* genes, as well as *Gyr A* and *Gyr B*. Of these, *tet* is transferable to non-resistant strains. Thus, indiscriminate use of antibiotics has created multiple antibiotic resistant strains of *Clostridium difficile*. Judicious and planned antibiotic treatment is advisable.

Keywords- *Clostridium difficile*, multidrug resistance, hospital-acquired diarrhoea, fluoroquinolones, vancomycin, metronidazole, virulent strains

Background

Clostridium difficile which has been named as *Clostridioides difficile* recently, is a gram positive, anaerobic and forms spores. Even healthy infants harbour it in its gut flora. The organism spreads via the fecal-oral route and is amongst the most common fecal-oral bacteria that is associated with nosocomial infections. The principal cause of diarrhoea amongst the hospitalised patients has been attributed to *Clostridium difficile* infection(1). *Clostridium difficile* infection has varied manifestations in infected people , having on one hand asymptomatic carriage to another hand a very severe form ; toxic megacolon. This varied manifestation has been attributed to host and pathogen interactions (1).The management of the *Clostridium Difficile* has been a difficult task for the clinicians not only because of its various manifestations but also due to the rapid evolve of new strains of *Clostridium Difficile* who are resistant to the existing antibiotics regime (1).

*Clostridium difficile*infections are amongst the most common cause of diarrhoea in nosocomial diarrhoea in India, with more than one million cases recorded every year(1). The causes of the global resistance in *Clostridium difficile* are overcrowded population,increased movement across countries, over-prescription and self prescription of antibiotics in may countries among human and animal populations , wildlife spread, indiscriminate use of antibiotics without indications as well as poor system for proper sewage disposal and sanitation(2). Beside being asymptomatic *Clostridium difficile* infection in an individual can manifest as foul-smelling diarrhoea, with nausea and fever, three or more times a day for more than two days or toxic megacolon, specially in hospitalised patients (3). The treatment with antibiotic which may irradiate the *Clostridium difficile* is a herculean job as the molecular mechanism which makes the *Clostridium difficile* more susceptible to a particular antibiotic is still unravelled .The new strains which prop up in subsequent generations, which show resistant to the antibiotics forms a very crucial key to the basis of the steps in the diagnostic and management of the *Clostridium Difficile*(4). The molecular basis to identify *Clostridium Difficile* genotype includes a variety of molecular methods like ; PCR ribotyping, multilocus sequence typing (MLST), pulsed-field gel electrophoresis (PFGE), multilocus variable-number tandem-repeat analysis (MLVA), and sequencing of functional genes such as *slpA* and *tcdC* (5). Each of these typing methods has its own advantages and disadvantages, and may be applied in different occasions according to the study Table 1(3).A definitive diagnosis may be made through a polymerase chain reaction test for the presence of specific bacterial genes (6).

Table 1. Diagnostic Method for *Clostridium difficile* and target.

Testing Method	Target(s)	Notes	Advantages	Disadvantages	Sensitivity and Specificity percentage
Rapid Diagnostic Tests					
Real Time-PCR(7)	tcdB or tcdC genes	• for tcdA- / tcdB+ strains cause disease	Rapid and accurate		Sensitivity of 73.3% and a specificity of 99.2%.

EIA(8)	GDH Glutamate Dehydrogenase EIA	Cannot be used alone and has to be paired with another test for confirmation	Rapid highly sensitive	Not for use as a standard test since toxin must be confirmed	sensitivity and specificity of 93% and 89%, respectively.
EIA(9)	Toxins A or B	• Fast with Variable specificity and sensitivity	moderate specificity		50% to 90% and 70% to 95%, respectively
NAAT(10)		Highly sensitive and specific for Toxicogenic Clostridium Difficile but the test is costly	High sensitivity fast test	Besides infections may also detect colonisation of Clostridium Difficile	Over 90% or even close to 100%; these values depend on the test that was used as a reference test
LAMP(11)	<i>tcdA</i> or <i>tcdB</i> genes	Rapid	High Sensitive		
Routine Gold Standard Tests					
Toxigenic Culture(12 and 13)	Toxigenic Clostridium difficile	•Reference standard •Difficult to perform	High sensitivity; the organism recovered can be used for genotyping and thus facilitate in the antimicrobial susceptibility studies mechanism	• long time for the final result	sensitivity and specificity of 93% and 89%, respectively

Abbreviations: *CDI*, *Clostridium difficile* infection; EIA, enzyme immunoassay; GDH, glutamate dehydrogenase; **LAMP**, loop-mediated isothermal amplification; **NAAT**, nucleic acid amplification testing; **RT-PCR**, real-time polymerase chain reaction.

Clostridium difficile form spores which facilitate the spread of infection among the population. The spread of this aerobic bacteria is further facilitated by its production of toxins; enterotoxin, Toxin A and cytotoxin, Toxin B. *Clostridium difficile* is well-known for its resistance to antibiotics including broad spectrum. It is stated that, previous history of indiscriminate antibiotic use is a predisposing risk factor for antibiotic resistance (14). Fluoroquinolones and cephalosporins, in particular, which has been used in the treatment of *Clostridium difficile* infection has been associated with the evolution of antibiotic resistance in *Clostridium difficile*. Medicines like proton pump inhibitors like esomeprazole, and H2 receptor antagonist, famotidine or ranitidine increase the risk of resistance to multiple times (15). Infection with *Clostridium difficile* strains resistant to multiple antibiotics is associated with an increased burden on the health system, expenses on part of the patient, and poor prognosis leading to even death (16).

Rationale:

In a densely-populated country such as India, multidrug resistant pathogens can quickly cause an epidemic, since there is a significant potential for rapid spread among vulnerable populations. Therefore, it is empirical that a thorough knowledge of the biochemical mechanism by which the bacteria escapes the antibiotic treatment and develops resistance to an antibiotic must be understood. This mystery of antibiotic resistance unravelled in *Clostridium difficile* would be very useful in arresting the rise of resistant species through monitoring and modification of antibiotic usage policies. Therefore, we decided to put forth few literature put in by scientists and throw a light on the phenomenon of antibiotic resistance in *Clostridium difficile*.

Objective: The objective was to review state-of-the-art as well as classical literature available regarding the emergence and mechanisms by which the *Clostridium difficile* attains antibiotic resistance.

Methodology:

Search strategy ; study selection

The electronic online reference databases including PubMed, Scopus, Medline, Embase, web of sciences, cochrane library and Google Scholar were used to explore for literature in relation to antibiotic resistance in *Clostridium difficile*. Research articles using the keywords; *Clostridium difficile*, antibiotic, resistance, strain, ribotype, fluoroquinolone, and genes; along with combinations of the Boolean operators, “AND” and “OR”. Various combinations yielded 10000+ research papers, and further selection was based on content and completeness of data. Since this was a scoping review, literature that reported resistance to a few or specific antibiotics were also included. Preference was given to research papers which covered resistance study for at least nine antibiotics. The search was further refined for recency and relevance using the year of publication choice in Google Scholar, “From 2017-“. Research papers published in reputed journals and with greater number of citations were studied carefully. Finally, the data obtained were summarized to yield an understanding of the situation and arrive at a recommendation.

Epidemiology of antibiotic resistance of *Clostridium difficile*

Initially, in the late 1970's, Clindamycin was recognized as the single antibiotic that predisposed patients to persistent *Clostridium difficile* infections (17). Clindamycin – resistant *Clostridium difficile* outbreaks were repeatedly reported in the continents of United States of America, Canada and there after in Europe (18). As a result, the use of clindamycin was

curtailed in the United States and Europe, leading to a reduced risk of contracting infection by *Clostridium* in the next decade (19).

Cephalosporins were the next class of antibiotics that *Clostridium difficile* became resistant to, in the late 1990's and early 2000's (20,21).79% of recent virulent strains are resistant to first-generation cephalosporins, while about 35% are resistant to second-generation cephalosporins (22).

There was an epidemic of *Clostridium difficile* infection between 2000-2005, which led to the closer examination of the bacterium, since the financial burden and burden upon the health system was considerable high (23). *Clostridium difficile* – associated diarrhoea (CDAD) was attributed during this epidemic to long duration use of antibiotics and the subsequent emergence of highly virulent forms of *Clostridium difficile* .

Over a short period of years strains of *Clostridium difficile* with increased virulence have evolved across continents. Classification of these strains is based either on the PCR ribotype or on pulsed field gel electrophoresis results combined with specific restriction endonuclease subtypes (24). In particular, RT 027 evolved as a result of fluoroquinolone overuse. Other virulent strains include RT 001/072, RT 078, RT 126, and RT 014/020. The evolution of *Clostridium difficile* resistant has been extensively studied in Europe (25,26). Several ribotypes have emerged in Europe, with RT027 appearing most frequently. A greater variety of ribotypes were seen in the UK, and in Portugal, Poland, Austria, Switzerland, France, Netherlands, Greece, and Germany, while Slovakia, Cyprus, Denmark, and Hungary had comparatively fewer variants. RT027 was the prevailing variant in Hungary and Poland. All European varieties, are, however, sensitive to fidaxomicin, metronidazole and vancomycin(2).

Clostridium difficile has also been isolated from domestic animals in Europe, particularly from pigs (62 strains) and dogs (39 strains). The most common *Clostridium difficile* ribotypes found these cases were 078 and 010.Type 010, found in dogs, is non-toxicogenic. RT078, 150, 127, 014/020, 126, and 031/1 were antibiotic-resistant. Similarly, *Clostridium difficile* exhibited multiple ribotypes in United States of America, with increased resistance to metronidazole, but not vancomycin. Global studies have found an increasing resistance to metronidazole and vancomycin(27).

Table 2 :*Clostridium difficile* , mechanisms of resistance of few antibiotics.

Sr. No.	Antibiotics used in the treatment	Resistance developed bio-mechanism	Genes affected	Remarks
1.	Metronidazole (MTZ)(28)	Biofilm formation		
2.	Ciprofloxacin(29)	Quinolone resistance-determining regions (QRDRs) showing alterations in the target enzymes through stepwise mutations)	gyrA and gyrB,	Ciprofloxacin

3.	Vancomycin (24)	biofilm formation		mur G
4.	Chloramphenicol(30)	Acetyl transferase	Tn4453a and Tn4453b	cat D
5.	Rifamycins (31)	Altered target		rpo B
6.	Cephalosporins(30)	Antibiotic enzymatic destruction		Putative β -lactamases and PBPs
7.	Fluoroquinolones (30)	Altered target	gyr A/gyr B	
8.	Ciprofloxacin(6)	Mutational alterations in target enzymes through stepwise mutations in the quinolone resistance-determining regions (QRDRs)	gyrA and gyrB,	topoisomerase IV genes

The drugs of choice thus remain Metronidazole and Vancomycin. For example, cephalosporins are destroyed enzymatically via altered beta-lactamase action, while 23 S RNA methylases and RNA methyltransferases render Erythromycin ineffectual. Efflux mechanisms may or may not be involved in this resistance. *Clostridium difficile* presents an altered target to fluoroquinolones, with the gyr A/ gyr B locus involved (28). Rifamycins and Vancomycin also find their targets altered in resistant strains. Fidaxomicin and vancomycin target bacterial RNA polymerases, and thus in resistant varieties, rpoB or rpoC may be mutated (29). Additionally, altered peptidoglycan synthesis may also be involved in vancomycin resistance through mutations in the MurG gene (Figure 1, Table 2).

Multiple amino acid changes in bacterial proteins have been associated with antibiotic resistance. Particularly, Threonin at position 82 is a hotspot, with amino acids changing commonly to Isoleucine, alanine or valine. Multiple mechanisms thus exist in the antimicrobial resistance exhibited by *Clostridium difficile* . Usually, antibiotic resistance can be through reduced uptake, increased efflux, or enzymatic (30). The mechanisms employed by multiple-antibiotic-resistant *Clostridium difficile* are heterogenous. An important aspect is the mobility of resistance-conferring portions of the genome between bacteria(30). The ribotype variation seen to occur in *Clostridium difficile* is maintained via mobile genetic elements, particularly, transposons, which participate in transduction, transformation and conjugation. Mobile genetic elements almost constitute as much as 11% of the *Clostridium difficile* genome and this is the reason for the constantly changing face that the bacterium presents to antibiotics(31).Tn5398, Tn5215, and Tn6194 ribotypes together confer resistance to erythromycin and other MLSB (macrolide-lincosamide-streptogramin B) family

antibiotics. Transposons Tn5397, Tn916, and Tn6164 transfer tet genes to the recipient bacterium, conferring tetracycline resistance. It is possible that this element was acquired in *Clostridium difficile* from another bacterium, such as Bifidobacterium. Transposons Tn4453a and Tn4453b, make the catP gene mobile, conferring chloramphenicol resistance upon the recipient via enhancing acetyltransferase activity (27).

The resistance exhibited to metronidazole is of special interest, since metronidazole is popularly prescribed and is an economical alternative, compared to vancomycin. The response is of a heterogenous and inducible nature. One method employed by the bacterium is to produce a thick impenetrable biofilm matrix made of proteins, polysaccharides and DNA, and regulated by the Cwp84 cysteine protease component(32). Metabolic pathways involved in conferring metronidazole resistance are poorly understood, but are thought to involve nitroreductases and DNA repair mechanisms, and, additionally, iron uptake(24). GyrA and GyrB genes may also be involved in fluoroquinolone resistance. GyrA/GyrB represent two subunits of a DNA gyrase that prevents fluoroquinolone-induced damage in *Helicobacter pylori* as well as *Clostridium difficile* (33).

The variants seen across continents are of significance in understanding the epidemiology, behaviour, and spread of *Clostridium difficile* among various populations. Agar dilution tests, broth dilution tests, MIC gradient diffusion, and other antibiotic susceptibility tests help determine whether an emerging variant is one of concern. Additionally, susceptibility studies help understand the nature of variations and the possible mechanisms of transfer of antibiotic resistance in *Clostridium difficile* (1).

This rapid rate and variety of mutations conferring both increased virulence and antibiotic resistance is of concern, since serious infections with *Clostridium difficile* may lead to debilitation and death(34). A high economic cost is also involved with the administration of third generation antibiotics and specialized formulations. Mobile elements within the bacterial genome ensure the rapid emergence of resistant variants, while already acquired resistances are retained by selection against a background of indiscriminate antibiotic use(35-39). The increased morbidity and mortality associated with the newer, more virulent strains, coupled with the multi-drug resistance seen to evolve, underscore the need for understanding the molecular bases behind the mechanisms of harm avoidance employed by the numerous strains, as well as the development of newer antibiotics and alternate treatment strategies(14).




Figure 1: The resistance developed after the mutation of the RNA polymerase enzyme of the *Clostridium difficile*

Conclusion

The greatest burden of *Clostridium difficile* infections globally is its resistance to antibiotics which has become a matter of utmost importance since its major cause of nosocomial diarrhoea. To add, further, to the burden, new strains of the *Clostridium difficile* have evolved following unscrupulous use of antibiotics which over the years have increased in incidence and newer strains keep on emerging. The underlying biochemical mechanism inherent in *Clostridium difficile* is an interesting mystery to unravel which will certainly bring a drop in the incidence of this burden and help scientists in the development of new drugs against it. Finally, attention has to be drawn towards the clinicians and self-administered populations towards the blind use of antibiotics which will in return facilitate to decrease the occurrence as well as spread of the drug resistant *Clostridium difficile* infection.

UNDER PEER REVIEW

References:

1. Starr J. *Clostridium difficile* associated diarrhoea: diagnosis and treatment. *BMJ Br Med J*. 2005;331(7515):498.
2. Aslam B, Wang W, Arshad MI, Khurshid M, Muzammil S, Rasool MH, et al. Antibiotic resistance: a rundown of a global crisis. Vol. 11, *Infection and Drug Resistance*. 2018. p. 1645–58.
3. Switaj TL, Winter KJ, Christensen S. Diagnosis and Management of Foodborne Illness. *Am Fam Physician*. 2015;92(5):358–65.
4. Cheng J-W, Xiao M, Kudinha T, Kong F, Xu Z-P, Sun L-Y, et al. Molecular Epidemiology and Antimicrobial Susceptibility of *Clostridium difficile* Isolates from a University Teaching Hospital in China. *Front Microbiol*. 2016;0(OCT):1621.
5. Manzoor SE, Tanner HE, Marriott CL, Brazier JS, Hardy KJ, Platt S, et al. Extended Multilocus Variable-Number Tandem-Repeat Analysis of *Clostridium difficile* Correlates Exactly with Ribotyping and Enables Identification of Hospital Transmission. *J Clin Microbiol*. 2011;49(10):3523.
6. Sholeh M, Krutova M, Forouzesh M, Mironov S, Sadeghifard N, Molaeipour L, et al. Antimicrobial resistance in Clostridioides (*Clostridium*) *difficile* derived from humans: a systematic review and meta-analysis. *Antimicrob Resist Infect Control* 2020 91. 2020;9(1):1–11.
7. Song PH, Min JH, Kim YS, Jo SY, Kim EJ, Lee KJ, et al. Rapid and accurate diagnosis of *Clostridium difficile* infection by real-time polymerase chain reaction. *Intest Res*. 2018;16(1):109.
8. Novak-Weekley SM, Marlowe EM, Miller JM, Cumpio J, Nomura JH, Vance PH, et al. *Clostridium difficile* testing in the clinical laboratory by use of multiple testing algorithms. *J Clin Microbiol*. 2010;48(3):889–93.
9. Bella S Di, Ascenzi P, Siarakas S, Petrosillo N, Masi A di. *Clostridium difficile* Toxins A and B: Insights into Pathogenic Properties and Extraintestinal Effects. *Toxins (Basel)*. 2016;8(5).
10. Humphries RM, Uslan DZ, Rubin Z. Performance of *Clostridium difficile* Toxin Enzyme Immunoassay and Nucleic Acid Amplification Tests Stratified by Patient Disease Severity. *J Clin Microbiol*. 2013;51(3):869.
11. O'Horo JC, Jones A, Sternke M, Harper C, Safdar N. Molecular techniques for diagnosis of *Clostridium difficile* infection: Systematic review and meta-analysis. In: *Mayo Clinic Proceedings*. 2012. p. 643–51.
12. Mirzaei EZ, Rajabnia M, Sadeghi F, Ferdosi-Shahandashti E, Sadeghi-Haddad-Zavareh M, Khafri S, et al. Diagnosis of *Clostridium difficile* infection by toxigenic culture and PCR assay. *Iran J Microbiol*. 2018;10(5):287.
13. R C, DB L. The role of toxins in *Clostridium difficile* infection. *FEMS Microbiol Rev*. 2017;41(6):723–50.
14. Fair RJ, Tor Y. Antibiotics and Bacterial Resistance in the 21st Century. *Perspect Medicin Chem*. 2014;6(6):25.

15. Saha S, Kapoor S, Tariq R, Schuetz AN, Tosh PK, Pardi DS, et al. Increasing antibiotic resistance in *Clostridioides difficile*: A systematic review and meta-analysis. *Anaerobe*. 2019;58:35–46.
16. DePestel DD, Aronoff DM. Epidemiology of *Clostridium difficile* Infection. *J Pharm Pract*. 2013;26(5):464.
17. JG B. Historical perspectives on studies of *Clostridium difficile* and *C. difficile* infection. *Clin Infect Dis*. 2008;46 Suppl 1(SUPPL. 1).
18. S J, MH S, KA F, GE K, FC T, D L, et al. Epidemics of diarrhea caused by a clindamycin-resistant strain of *Clostridium difficile* in four hospitals. *N Engl J Med*. 1999;341(22):1645–51.
19. DN G. Clindamycin, cephalosporins, fluoroquinolones, and *Clostridium difficile* -associated diarrhea: this is an antimicrobial resistance problem. *Clin Infect Dis*. 2004;38(5):646–8.
20. R K, J C. Impact of changes in antibiotic policy on *Clostridium difficile* -associated diarrhoea (CDAD) over a five-year period in a district general hospital. *J Hosp Infect*. 2003;54(2):104–8.
21. Bignardi GE. Risk factors for *Clostridium difficile* infection. Vol. 40, *Journal of Hospital Infection*. 1998. p. 1–15.
22. Monot M, Eckert C, Lemire A, Hamiot A, Dubois T, Tessier C, et al. *Clostridium difficile* : New Insights into the Evolution of the Pathogenicity Locus. *Sci Rep*. 2015;5.
23. J P, L V, S G, S R, I B. Outcomes of *Clostridium difficile* -associated disease treated with metronidazole or vancomycin before and after the emergence of NAP1/027. *Am J Gastroenterol*. 2007;102(12):2781–8.
24. Z P, D J, HB K, CW S, B W, YW T, et al. Update on Antimicrobial Resistance in *Clostridium difficile* : Resistance Mechanisms and Antimicrobial Susceptibility Testing. *J Clin Microbiol*. 2017;55(7):1998–2008.
25. J F, J V, K M, S N, S T, C L, et al. Pan-European longitudinal surveillance of antibiotic resistance among prevalent *Clostridium difficile* ribotypes. *Clin Microbiol Infect*. 2015;21(3):248.e9-248.e16.
26. Kuijper EJ, Coignard B, Brazier JS, Suetens C, Drudy D, Wiuff C, et al. Update of *Clostridium difficile* -associated disease due to PCR ribotype 027 in Europe. *Euro Surveill*. 2007;12(6).
27. Spigaglia P, Drigo I, Barbanti F, Mastrantonio P, Bano L, Bacchin C, et al. Antibiotic resistance patterns and PCR-ribotyping of *Clostridium difficile* strains isolated from swine and dogs in Italy. *Anaerobe*. 2015;31:42–6.
28. IT B, EA Y, RMCP D, EO F. *Clostridium difficile* : a problem of concern in developed countries and still a mystery in Latin America. *J Med Microbiol*. 2012;61(Pt 2):169–79.
29. JA L, M S, S M, SW B, A R. In vitro selection, via serial passage, of *Clostridium difficile* mutants with reduced susceptibility to fidaxomicin or vancomycin. *J Antimicrob Chemother*. 2014;69(1):41–4.

30. J P, N S, MA C, ME A, MP C, S A, et al. Emergence of fluoroquinolones as the predominant risk factor for *Clostridium difficile* -associated diarrhea: a cohort study during an epidemic in Quebec. *Clin Infect Dis*. 2005;41(9):1254–60.
31. E V, MD C, BW W. The *Clostridium difficile* PCR ribotype 027 lineage: a pathogen on the move. *Clin Microbiol Infect*. 2014;20(5):396–404.
32. A S, W M, V L-B, S S, Y D, J S, et al. An adherent Gardnerella vaginalis biofilm persists on the vaginal epithelium after standard therapy with oral metronidazole. *Am J Obstet Gynecol*. 2008;198(1):97.e1-97.e6.
33. Lachowicz D, Pituch H, Wultańska D, Kuijper E, Obuch-Woszczatyński P. Surveillance of antimicrobial susceptibilities reveals high proportions of multidrug resistance in toxigenic *Clostridium difficile* strains in different areas of Poland. *Anaerobe*. 2020;62.
34. Smits WK, Lyras D, Lacy DB, Wilcox MH, Kuijper EJ. *Clostridium difficile* infection. *Nat Rev Dis Prim*. 2016;2:1–20.
35. Paterson DL. Resistance in Gram-Negative Bacteria: Enterobacteriaceae. *Am J Med*. 2006;119(6 SUPPL. 1).
36. Banawas SS. *Clostridium difficile* Infections: A Global Overview of Drug Sensitivity and Resistance Mechanisms. Vol. 2018, BioMed Research International. 2018.
37. Dahake, Prasanna T., Sudhindra M. Baliga, Vijay M. Kumbar, and Kishore G. Bhat. “Cytotoxicity of Novel Polymeric Gel Matrix Triple Antibiotic Paste-an In Vitro Study.” REGENERATIVE ENGINEERING AND TRANSLATIONAL MEDICINE, n.d. <https://doi.org/10.1007/s40883-020-00191-x>.
38. Abbafati, Cristiana, Kaja M. Abbas, Mohammad Abbasi, Mitra Abbasifard, Mohsen Abbasi-Kangevari, Hedayat Abbastabar, Foad Abd-Allah, et al. “Five Insights from the Global Burden of Disease Study 2019.” *LANCET* 396, no. 10258 (October 17, 2020): 1135–59.
39. Abbafati, Cristiana, Kaja M. Abbas, Mohammad Abbasi, Mitra Abbasifard, Mohsen Abbasi-Kangevari, Hedayat Abbastabar, Foad Abd-Allah, et al. “Global Burden of 369 Diseases and Injuries in 204 Countries and Territories, 1990-2019: A Systematic Analysis for the Global Burden of Disease Study 2019.” *LANCET* 396, no. 10258 (October 17, 2020): 1204–22.