

Original Research Article

Correlation of vitamin D levels with polycystic ovary syndrome: A case control study

Abstract

Objective: To compare vitamin D status in, women with PCOS & fertile women without PCOS and its subsequent evaluation.

Introduction: PCOS is an endocrine disorder of women in reproductive age, characterised by obesity, hyperandrogenemia and insulin resistance. Women with PCOS tend to be overweight and have increased risk of development of Type II Diabetes and cardiovascular disease. Exact Etiology of PCOS still remain an enigmatic dilemma however various studies conducted till date include diet and lifestyle modification as the key factor to promote health, BMI, reduced hyperinsulinemia and reduce the risk of development of PCOS.

Main aim of our study was to compare vitamin D status in women with PCOS, with fertile women in Uttarakhand.

Methodology: The conducted study was cross sectional, involving the enrolment of 100 women comprising of 50 women with PCOS and 50 fertile women without PCOS. The serum 25-hydroxy vitamin D and other metabolic markers were measured. Vitamin D deficiency was defined as serum 25 (OH) D concentrations less than 20 ng/ml.

Result: Serum 25-hydroxyvitamin D was significantly lower in women with PCOS compared to fertile controls ($p < 0.0001$), and the prevalence rates of 25(OH) D deficiency and insufficiency were higher in women with PCOS than in fertile women ($p < 0.0001$). The prevalence of 25 (OH) D deficiencies in PCOS women was significantly higher. Serum 25 (OH) D concentrations was significantly negatively correlated with body mass index (BMI), waist-to-hip ratio (WHR), fasting insulin, total cholesterol and low-density lipoprotein cholesterol (LDL-C), ($P < 0.05$). In comparison, serum 25 (OH) D concentrations were significantly positively correlated with high-density lipoprotein cholesterol (HDL-C) ($P < 0.05$). Increased BMI and WHR, high levels of fasting insulin, total cholesterol and LDL-C were regarded as risk factors, but high level of HDL-C was considered to be protective factor of vitamin D deficiency in PCOS women.

Conclusion: The study demonstrated that women with PCOS have a significantly lower 25(OH) D compared to fertile controls. A compromised vitamin D status in PCOS women is associated with a higher prevalence and metabolic risk of PCOS in women.

Introduction

PCOS is an endocrine disorder of women in reproductive age and is the main cause of anovulatory infertility (1). The prevalence of PCOS range from 9 to 18% (2,3). Characteristic feature of PCOS is presence of polycystic ovaries and menstrual dysfunction leading to

infertility along with biochemical abnormality. PCOS manifest clinically as hirsutism, Hyperandrogenism and acne (4). The most prevalent characteristics of PCOS is Obesity, Hyperandrogenemia and insulin resistance (5). The prevalence of PCOS will rise in near future (6). The potential impact will be negatively on population growth and cardiovascular morbidity and mortality, and consequently, it will lead to major public health concern. The development of PCOS is influenced by Vitamin D and hormonal modulation influence insulin metabolism and fertility regulation (7,8). Several studies have reported low levels of vitamin D in women with PCOS, with average 25-hydroxy vitamin D (25OHD) levels between 11 and 31 ng/ml,(7,9,10,8,11-17) with the majority having values <20 ng/ml (67–85%).(9,18,19,8,13,14) The functioning of all systems of the body is disrupted by vitamin D deficiency, which increases the risk of chronic disease including physical diseases such as cancer, cardiovascular, autoimmune and infectious diseases; and psychological disorders such as depression and chronic pain (20). Vitamin D3 is obtained from the diet or synthesised endogenously through sunlight- induced photochemical conversion of cholesterol in the skin and subsequently hydroxylation in the liver and kidney. In the skin, 7-dehydrocholesterol undergoes ultraviolet photolysis to form vitamin D3. Vitamin D3 then undergoes two successive hydroxylations, the first of which takes place in the liver and is catalysed by vitamin D-25 hydroxylase to form 25OHD. The second hydroxylation step is regulated by parathyroid hormone (PTH) and mediated by 25-hydroxyvitamin D3 1 α -hydroxylase and occurs predominately in the kidney. This second hydroxylation produces the final active metabolite of vitamin D3, which is 1,25-dihydroxyvitamin D3. 1, 25-dihydroxyvitaminD3 circulates bound to vitamin D-binding protein until it reaches its target tissue where it binds to vitamin D receptors to initiate its effect.

It has been very controversial regarding the difference between vitamin D levels between PCOS women and healthy women, and the relationship between vitamin D and metabolic factors in PCOS women. Certain studies have shown that PCOS women had lower serum 25 (OH) D concentrations than healthy women, and vitamin D deficiency.

Vitamin D deficiency and PCOS are associated with metabolic disorders, but little is known about vitamin D status in women with PCOS in Dehra Dun, Uttarakhand. Therefore, this study aims to investigate vitamin D status and analyse the relationship between vitamin D deficiency and metabolic risk factors in women with PCOS.

Methodology

It was an observational, case control study conducted from July 2019 to January 2020. The sample size was n= 100 out of which 50 were control and 50 patients belonged to PCOS group. All women fulfilling the inclusion criteria were included after informed consent. Non probability sampling technique was used for the purpose of data collection. All the young women of age (18-40), and those with polycystic ovary syndrome for more than or equal to one year duration were included. PCOS women can be diagnosed by two of the three criteria; : Oligoovulation or anovulation, clinical or biochemical evidence of hyperandrogenism and polycystic ovaries (12 follicles of size 5-7mm). Vitamin D deficiency was defined as 25-hydroxyvitamin D [25OHD] level < 20 ng/ml. Women with thyroid disease, congenital adrenal hyperplasia or autoimmune diseases were excluded. The protocol was approved by the Institutional Ethical committee of the hospital.

Statistical Analysis

Data in this study was analyzed using SPSS version 19.0. The continuous variables are presented as mean \pm SD, which were performed by Student's t-test or variance analysis. The categorical parameters are displayed as numbers (%), which were analyzed by chi-square test.

Linear regression analysis was used to analyze the correlation of 25 (OH) D concentrations with metabolic parameters. Post-stratification was done by applying Chi-square test. $P \leq 0.05$ was taken as significant.

Result

The numerical range for age in our study was from 18 to 45 years with mean age of control 24.02 ± 4.3 and PCOS group 23.16 ± 5.0 years, mean duration of PCO was 2.5 ± 1.2 years and mean BMI for control 24.48 ± 5.72 and PCOS 25.90 ± 5.16 . Overall, 29(38.16%) control and 47(61.84%) PCOS group of women had vitamin D deficiency which was statistically significant ($p=0.0001$). (Table1) When outcome variable was stratified with respect to age, BMI, waist hip ratio, systolic and diastolic blood pressure, it showed no significant difference as p-values were 0.3577, 0.1942, 0.4663, 0.1547 and 0.2098.752 respectively. Table 2 shows the clinical and biochemical profiles of the subjects. Women with PCOS showed increased serum levels of FSH, LH, TSH and T3 compared with matched controls, but they showed no differences in the levels of T4. There was marked difference in Vitamin D levels or prevalence of vitamin D deficiency (< 20 ng/ mL) (Table 2). In addition, we found correlations between serum vitamin D level and clinical or metabolic profiles in both PCOS patients and controls (Table 2). Vitamin D insufficiency was observed in the majority of the subjects (61.84% of patients and 38.16% of the controls Table 2).

Basic Characteristic of the Studied Groups

Table 1 illustrates the baseline characteristics of women in the two groups.

Table 1: Baseline characteristics of women between the two groups.			
	Control N=50	PCOS Group N=50	P-value
Age(year)	24.02 ± 4.3	23.16 ± 5.0	0.3577
BMI	24.48 ± 5.72	25.90 ± 5.16	0.1942
Waist circumference	102.86 ± 15.35	94.58 ± 12.78	0.0042
Hip circumference	0.95 ± 0.06	0.94 ± 0.06	0.0031
Waist hip ratio	0.95 ± 0.06	0.94 ± 0.06	0.4663
Blood Pressure			
Systolic	122.02 ± 7.41	124.28 ± 8.3	0.1547
Diastolic	84.8 ± 5.79	83.2 ± 6.8	0.2098

Clinical and Biochemical profiles of the subjects

Table 2: illustrates the clinical and biochemical profiles of women in the two groups.

Table 2: The Clinical and Biochemical profiles of women in the two groups.
--

	Control N=50	PCOS Group N=50	P-value
FSH	4.44±5.22	4.51±5.84	0.4586
LH	2.89±7.22	3.6±7.76	0.9231
TSH	1.76±2.19	1.79±2.56	0.1192
T ₃	81.0±89.26	84.2±89.12	0.2220
T ₄	5.98±6.21	5.98±7.63	0.0729
Triglyceride	130.31±17.89	124.13±29.25	0.2059
Total Cholesterol	166.08±25.05	151.29±30.76	0.0098
LDL	83.78±28.03	75.62±23.88	0.1207
Vitamin D ₃	21.09±18.07	11.91±10.57	0.0001

Table 3: illustrate the Vitamin D Outcome in PCOS patients and Control group

Table 3: The Vitamin D Outcome in PCOS patients and Control group			
Variable	Control N=50	Case N=50	p- value*
Vitamin D3	18.2 (6-66.5) 21.9±13.4	11.2 (5.23-31.2) 11.9 ±4.7	<0.0001
Sufficient (>=20.00)	21 (42)	3 (6)	1 (ref)
Low Vitamin D3 (<20.00)	29 (58)	47 (94)	11.3 (3.1-41.4)#

Data expressed as Mean±Sd and Median (Min-Max) , f(%). *Ranksum test, # Odd's ratio (95%CI)

It was observed that the mean vitamin D3 level among cases was less as compared to controls, and it was found statistically significant. The prevalence of vitamin D deficiency was high in the study population (76%) however the odds of having disease was 11 times more in Vitamin D deficient as compared to normal.

No significant difference was found when comparing other baseline characteristics between the two groups (P > 0.05).

Vitamin D Status between PCOS Women and Controls

The serum 25 (OH) D concentrations were significantly lower in PCOS women than in controls (11.91± 10.57 vs. 21.09 ± 18.07 ng/mL, P < 0.001). In addition, the serum 25 (OH) D concentration deficiencies were significantly high in women with PCOS than in controls (61.84% vs. 38.16%, P < 0.0001). Furthermore, the prevalence of normal 25 (OH) D statuses

in women with PCOS was significantly lower than that in controls (12.5% vs. 87.5%, $P < 0.0001$). Table 2 shows the Clinical and Biochemical profile of patients in two groups. There were statistically significant differences in HDL and Abdomen girth among the two groups ($P < 0.05$). No significant difference was found when comparing FSH, LH, TSH, T3, and T4, total cholesterol, LDL and triglycerides among two groups ($P > 0.05$).

Discussion:

In women with PCOS, low vitamin D levels are associated with obesity, metabolic and endocrine disturbances and vitamin D supplementation might improve menstrual frequency and metabolic disturbances in those women.[21] Vitamin D deficiency is still considered a problem of the past by health care professionals and the public. Populations at risk include infants, children, pregnant and postmenopausal women.

This study focuses on the serum levels of vitamin D and the prevalence of vitamin D deficiency in patients with PCOS. Our study found significant differences in the absolute level of serum vitamin D or prevalence of vitamin D deficiency between women with PCOS and control group. Additionally, we found correlations between serum vitamin D and hormonal or metabolic profiles in either PCOS patients or controls. Our findings suggest that there is role of vitamin D in the pathogenesis of PCOS. Similar to our study Wehr et al. [20] reported lower serum vitamin D levels in a large number of women with PCOS ($n = 545$) compared to controls ($n = 145$) (25.7 vs. 32.0 ng/mL, respectively), a substantial number of studies suggest that serum vitamin D levels are similar in women with and without PCOS [12,17,30]. In previous studies, average serum vitamin D levels in women with PCOS were reported to be between 11 ng/mL and 31 ng/mL, with the majority having mean values < 20 ng/mL [12,13,15-17,20,22,29, 31-33]. In our study, the mean 25-(OH) D3 level in women with PCOS was also < 20 ng/mL (11.91 ± 10.57 ng/mL), and vitamin D deficiency (lower than 20 ng/mL) was observed in 61.84% of patients. However, vitamin D deficiency is also common in the control group, with 60% of adults having values lower than 20 ng/mL. The control subjects in the current study also showed a high prevalence of vitamin D deficiency (38.16%), with a mean level of 21.09 ng/ mL. Many studies have investigated an association between vitamin D status and hormonal or metabolic features in PCOS. In women with PCOS, a low vitamin D level is thought to be related to metabolic risk factors such as insulin resistance, high total cholesterol, blood pressure, glucose, C-reactive protein, triglycerides, and low high-density lipoprotein (HDL) cholesterol [14,17]. In addition, vitamin D replacement therapy may have a beneficial effect on insulin resistance or fasting and on stimulated glucose and triglycerides levels in women with PCOS [16, 33]. Furthermore, several studies have identified relationships between low vitamin D status and measures of hyperandrogenism such as SHBG, the degree of hirsutism, FAI, total T and dehydroepiandrosterone sulphate [13, 14, 17, and 29]. Although our study found significant differences in the absolute level of serum vitamin D or prevalence of vitamin D deficiency between PCOS women and matched controls, the results need to be interpreted with caution. First, vitamin D deficiency may be a universal phenomenon across PCOS patients and controls. Second, inverse associations between obesity (BMI, body fat and waist measurements) and serum vitamin D levels have been reported in many studies [12-14, 17, 22, and 29]. As vitamin D is fat soluble, a higher proportion of vitamin D may be sequestered in adipose tissue in obese individuals, which might lower serum levels. In our study, patients and controls were matched by BMI, thus there were no differences in BMI, WC, or body and visceral fat masses between the two groups. Third, a potential limitation of the present study is the modest sample size in the PCOS group, which precludes drawing strong conclusions. Finally, we did not evaluate the presence of other potential confounding factors, such as outdoor times or dietary patterns which could affect the serum vitamin D levels. In summary,

we found significant difference in the absolute level of serum vitamin D or prevalence of vitamin D deficiency between women with PCOS and matched controls. The women with PCOS are at 11.34% at risk of developing vitamin D deficiency. Additionally, we did not find any correlation between serum vitamin D and hormonal or metabolic profiles in either PCOS patients or controls. Although our findings suggest that the role of vitamin D in the pathogenesis of PCOS is clear, vitamin D deficiency is a common finding among PCOS patients and controls. Finally, the potential relationship between vitamin D and PCOS requires further investigation, since vitamin D deficiency has been continuously proposed to increase the risk of insulin resistance and T2DM, which is also a core pathophysiology of PCOS.

References:

1. Azziz R, Woods KS, Reyna R, Key TJ, Knochenhauer ES, Yildiz BO. The prevalence and features of the polycystic ovary syndrome in an unselected population. *J Clin Endocrinol Metabol.* (2004) 89:2745–9.
2. March WA, Moore VM, Willson KJ, Phillips DI, Norman RJ, Davies MJ. The prevalence of polycystic ovary syndrome in a community sample assessed under contrasting diagnostic criteria. *Hum Reprod.* (2010) 25:544–51.
3. The Amsterdam ESHRE/ASRM-Sponsored 3rd PCOS Consensus Workshop Group. Consensus on women's health aspects of polycystic ovary syndrome (PCOS). *Hum Reprod.* (2012) 27:14–24.
4. Dunaif, A. (1997) Insulin resistance and the polycystic ovary syndrome: mechanism and implications for pathogenesis. *Endocrine Reviews*, 18, 774–800.
5. Lim SS, Davies MJ, Norman RJ, Moran LJ. Overweight, obesity and central obesity in women with polycystic ovary syndrome: a systematic review and meta-analysis. *Hum Reprod Update.* (2012)18: 618–37.
6. Pasquali, R., Gambineri, A. & Pagotto, U. (2006) The impact of obesity on reproduction in women with polycystic ovary syndrome. *BJOG: An International Journal of Obstetrics and Gynaecology*, 113, 1148.
7. Panidis, D., Balaris, C., Farmakiotis, D. et al. (2005) Serum parathyroid hormone concentrations are increased in women with polycystic ovary syndrome. *Clinical Chemistry*, 51, 1691–1697.
8. Li, H.W.R., Brereton, R.E., Anderson, R.A. et al. (2011) Vitamin D deficiency is common and associated with metabolic risk factors in patients with polycystic ovary syndrome. *Metabolism: Clinical and Experimental*, 60, 1475–1481.
9. Hahn, S., Haselhorst, U., Tan, S. et al. (2006) Low serum 25-hydroxyvitamin D concentrations are associated with insulin resistance and obesity in women with polycystic ovary syndrome. *Experimental and Clinical Endocrinology and Diabetes*, 114, 577–583.

10. Yildizhan, R., Kurdoglu, M., Adali, E. et al. (2009) Serum 25-hydroxyvitamin D concentrations in obese and non-obese women with polycystic ovary syndrome. *Archives of Gynecology and Obstetrics*, 280, 559–563.
11. Wehr, E., Trummer, O., Giuliani, A. et al. (2011) Vitamin D associated polymorphisms are related to insulin resistance and vitamin D deficiency in polycystic ovary syndrome. *European Journal of Endocrinology*, 164, 741–749.
12. Mahmoudi, T., Gourabi, H., Ashrafi, M. et al. (2010) Calcitropic hormones, insulin resistance, and the polycystic ovary syndrome. *Fertility and Sterility*, 93, 1208–1214.
13. Thys-Jacobs, S., Donovan, D., Papadopoulos, A. et al. (1999) Vitamin D and calcium dysregulation in the polycystic ovarian syndrome. *Steroids*, 64, 430–435.
14. Selimoglu, H., Duran, C., Kiyici, S. et al. (2010) The effect of vitamin D replacement therapy on insulin resistance and androgen levels in women with polycystic ovary syndrome. *Journal of Endocrinological Investigation*, 33, 234–238.
15. Kotsa, K., Yavropoulou, M.P., Anastasiou, O. et al. (2009) Role of vitamin D treatment in glucose metabolism in polycystic ovary syndrome. *Fertility and Sterility*, 92, 1053–1058.
16. Wehr, E., Pieber, T.R. & Obermayer-Pietsch, B. (2011) Effect of vitamin D3 treatment on glucose metabolism and menstrual frequency in PCOS women—a pilot study. *Journal of Endocrinological Investigation*, 34, 757–763.
17. Muscogiuri, G., Policola, C., Prioletta, A. et al. (2012) Low levels of 25(OH)D and insulin-resistance: 2 unrelated features or a cause-effect in PCOS? *Clinical Nutrition*, PMID:22260937. [Epub ahead of print].
18. Yildizhan, R., Kurdoglu, M., Adali, E. et al. (2009) Serum 25-hydroxyvitamin D concentrations in obese and non-obese women with polycystic ovary syndrome. *Archives of Gynecology and Obstetrics*, 280, 559–563.
19. Wehr, E., Pilz, S., Schweighofer, N. et al. (2009) Association of hypovitaminosis D with metabolic disturbances in polycystic ovary syndrome. *European Journal of Endocrinology*, 0, EJE-9– EJE0432
20. Holick, M.F. (2007) Vitamin D Deficiency. *New England Journal of Medicine*, 357, 266–281.
21. Reis JP, von Muhlen D, Kritz-Silverstein D, Wingard DL, BarrettConnor E. Vitamin D, parathyroid hormone levels, and the prevalence of metabolic syndrome in community-dwelling older adults. *Diabetes Care* 2007; 30:1549-55
22. Wehr E, Trummer O, Giuliani A, Gruber HJ, Pieber TR, ObermayerPietsch B. Vitamin D-associated polymorphisms are related to insulin resistance and vitamin D deficiency in polycystic ovary syndrome. *Eur J Endocrinol* 2011; 164:741-9

23. Panidis D, Balaris C, Farmakiotis D, Rousso D, Kourtis A, Balaris V, et al. Serum parathyroid hormone concentrations are increased in women with polycystic ovary syndrome. *Clin Chem* 2005;51: 1691-7
24. Li HW, Brereton RE, Anderson RA, Wallace AM, Ho CK. Vitamin D deficiency is common and associated with metabolic risk factors in patients with polycystic ovary syndrome. *Metabolism* 2011; 60:1475-81.
25. Thomson RL, Spedding S, Buckley JD. Vitamin D in the aetiology and management of polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 2012; 77:343-50.
26. Mahmoudi T, Gourabi H, Ashrafi M, Yazdi RS, Ezabadi Z. Calcitropic hormones, insulin resistance, and the polycystic ovary syndrome. *Fertil Steril* 2010; 93:1208-14.
27. Selimoglu H, Duran C, Kiyici S, Ersoy C, Guclu M, Ozkaya G, et al. The effect of vitamin D replacement therapy on insulin resistance and androgen levels in women with polycystic ovary syndrome. *J Endocrinol Invest* 2010; 33:234-8.
28. Muscogiuri G, Policola C, Priolella A, Sorice G, Mezza T, Lassandro A, et al. Low levels of 25(OH)D and insulin-resistance: 2 unrelated features or a cause-effect in PCOS? *Clin Nutr* 2012; 31:476-80
29. Yildizhan R, Kurdoglu M, Adali E, Kolusari A, Yildizhan B, Sahin HG, et al. Serum 25-hydroxyvitamin D concentrations in obese and non-obese women with polycystic ovary syndrome. *Arch Gynecol Obstet* 2009; 280:559-63.
30. Thys-Jacobs S, Donovan D, Papadopoulos A, Sarrel P, Bilezikian JP. Vitamin D and calcium dysregulation in the polycystic ovarian syndrome. *Steroids* 1999; 64:430-5.
31. Kotsa K, Yavropoulou MP, Anastasiou O, Yovos JG. Role of vitamin D treatment in glucose metabolism in polycystic ovary syndrome. *Fertil Steril* 2009; 92:1053-8.
32. Wehr E, Pieber TR, Obermayer-Pietsch B. Effect of vitamin D3 treatment on glucose metabolism and menstrual frequency in polycystic ovary syndrome women: a pilot study. *J Endocrinol Invest* 2011; 34:757-63
33. Wehr E, Pilz S, Schweighofer N, Giuliani A, Kopera D, Pieber TR, et al. Association of hypovitaminosis D with metabolic disturbances in polycystic ovary syndrome. *Eur J Endocrinol* 2009; 161:575- 82
34. Hahn S, Haselhorst U, Tan S, Quadbeck B, Schmidt M, Roesler S, et al. Low serum 25-hydroxyvitamin D concentrations are associated with insulin resistance and obesity in women with polycystic ovary syndrome. *Exp Clin Endocrinol Diabetes* 2006;114: 577-83