

## Review Article

### Effectiveness Of Immunotherapy on Covid-19

#### Abstract:

COVID-19 is an infection which is triggered by a recently discovered corona virus, SARS-CoV-2. COVID-19 patients experience some respiratory illness which includes difficulty in breathing, pneumonia, cough etc. People also suffer from body-ache and extreme weakness. SpO<sub>2</sub> level of the patient becomes very less leading to worsening of the disease. Many people have lost their life due to corona and has become a pandemic. Recently after many investigations vaccine against this virus has been discovered which is considered to reduce the infection up to some extent. Still vaccinated people also have chances of getting infection, but the severity of symptoms gets reduced to much extent. It was found that novel vaccine technology used in COVID-19 vaccine can prevent upto 95% infection. Immunotherapy plays major role in fighting against the infection. According to studies made it is found that hematology patients are more prone to suffer from infection. The macrophages, neutrophils and dendritic cells helps in reducing the infection. There are mainly two types of vaccine discovered against coronavirus disease –

- Covaxin
- Covishield

Covaxin is 78% - 100% effective whereas Covishield is 70% - 90% effective.

Clinical studies are used to assess vaccine effectiveness. After receiving the corona disease vaccine, several negative side effects may arise. Vaccines rescue the breathes of millions of beings who are diseased with viruses and bacteria. We celebrate the advances made and the highpoint challenges ahead in vaccinations and the science underlying them as the world waits for safe and effective COVID-19 vaccines. We outlined the current prospective therapeutic methods for disorders caused by COVID-19 infection, as well as their mechanisms of action, safety, and efficacy, in this study.

#### INTRODUCTION:

This is a type of review article. The information for the article is taken from PUBMED, WHO AND GOOGLE. The objective of this short appraisal is to underline the effectiveness of COVID-19 vaccination, threat for COVID-19 extremity, and the repercussions in sufferers with haematological illnesses exacerbated by COVID-19. It also aims to access the threat linked with SARS-CoV-2 contagion in older people. It also highlights the importance of vaccines to cure the covid infection. This article also reflects the existing experience and evidence related to immunotherapy for COVID-19.

## 1. COVID- 19

COVID-19 is a infectious illness which is triggered due to newly discovered corona virus virus called SARS-CoV -2. It causes respiratory illness and lead to pneumonia. The maximum not unusualplace signs of COVID -19 are fever, dry cough, fatigue, and frame aches. SARS-CoV-2 can transmit through through many modes like contact and droplet transimission, air borne transmission etc. Some studies have detected RNA of SARS-CoV-2 in plasma and serum cells. COVID-19 can also be transmitted through secretions like saliva, respiratory secretions & droplets such as coughing, sneezing. SARS-CoV-2 RNA has additionally been determined in various biological models, including some sufferers' faeces and urine. Coronavirus disorder 2019 (COVID-19) is a new, hastily spreading viral infection resulting from contamination with the coronavirus 2 that reasons extreme acute respiration syndrome (SARS-CoV-2). The continuing coronavirus disease 2019 (COVID-2019) pandemic has expanded across 213 countries and infected approximately 1,870,000 people since April 15, 2020, posing an unprecedented threat to global health and the economy. The knowledge gained from previous administrations of respiratory viral diseases has aided in the treatment of SARS-CoV-2. Only a few of the new therapies that have been explored in clinical settings are supportive intervention, immunomodulatory medicines, antiviral therapy, and convalescent plasma transfusion. Several of these medications have been demonstrated to be particularly successful in treating people infected with the corona virus. Furthermore, intensive research and clinical studies are being conducted to assess the effectiveness of current drugs and identify potential therapeutic targets for the development of innovative COVID-19 therapeutics.

### **Immunotherapy and its importance**

Immunotherapy is the treatment for certain disease by activating or suppressing the immune system. They increases the immune response of the body . It is of utmost importance as it

plays role in host response for viral clearance in severe covid 19. Since, the epidemic of covid 19 there is need for control of this pandemic using several options such as immunotherapy. It includes various types of vaccines , monoclonal candidates etc. COVID 19 infection leads to formation of anti-SARS-CoV-2 antibodies. The antibody decreases the replication of virus through neutralization and also take part in covid 19 pathogenesis through antibody dependent process.

The recent vaccine against COVID-19 has marked the beginning of national and international debates on feasibility and safety for all the individuals. Therefore there is a urgent need to determine the effectiveness of vaccine in various types of individuals, i.e., individuals suffering from various types of disorders and diseases.

When the number of cases rises and transmission speeds up, there's a greater potential that new dangerous and more transmissible variations emerge, which can spread more quickly and cause more severe sickness.

Vaccines are proven effective against existing variations, according to the evidence we've gathered so far, notably in terms of reducing the risk of life-threatening disease, hospitalisation, and mortality. However, some variations have a minor impact on vaccinations' capacity to protect against mild illness and infection.

Vaccines are expected to remain effective against variants because to the broad immune response they elicit, which implies that even if the virus changes or mutations occur, vaccines are unlikely to become completely useless.

WORLD HEALTH ORGANISATION continues to constantly review the evidence and will update its guidance as we find out more.?’

Continuing to use tried-and-true public health strategies and immunizations is one of the best methods to protect against new variations. All vaccinations against COVID-19 that have been sanctioned by the World Health Organization for emergency use have been thoroughly evaluated and demonstrated to give a high level of protection against serious disease and death. As more powerful virus strains arise, it's critical to get your vaccine when it's time.

**RESULTS / FINDINGS:**

Some reports of SARS and MERS-COV have shown that poorly neutralized , already existing antibodies are developed as an outcome of vaccine or infection increase the chances of subsequent infection which is called as antibody-dependent enhancement phenomenon (ADE). (1) COVID-19 related morbidity and mortality is more prone in patients of Haematology who undergo immunotherapy.(2) 60 years or above age people are more affected to SARS-CoV-2. Preexisting morbidities **results in multiplied COVID-19 related mortality** in geriatric population. (3) Cancer patients infected with COVID who undergone anti-cancer treatments did not observe high risk of mortality and exacerbation. Surgery, chemotherapy, immunotherapy **aren't associated with multiplied threat** of death . However, 28 days chemotherapy treatment leads to increase risk of mortality. (4)

COVID-19 vaccine uses cutting-edge vaccination technology to prevent illness in up to 95% of people. On the other hand, The uncertainty of vaccination **protection in sufferers** with autoimmune inflammatory rheumatic disease (AIIRD) **who're immunocompromised because of immunological disorder** and immunosuppressive **remedy offers** unambiguous advice. According to Korean College of Rheumatology, the currently available vaccine for COVID-19 is safe and effective.(5)

The host's immunological reaction to COVID-19 might result in an abnormal inflammatory response or "cytokine storm," which worsens the patient's condition. Patients who were exposed to COVID-19 developed macrophage activation syndrome (MAS), which was characterised by reduced human leucocyte antigen D related expression and a decrease in lymphocytes and natural killer cells. (6)

According to recent study, it was concluded that the igG antibody concentration in mildly infected and **recuperating sufferers confirmed no distinction among male and lady sufferers**. However, in acute conditions it was observed that female patients have relatively high serum SARS-CoV-2 igG antibody concentration when compared with male.(7)

According to the recent study made, it was found that after taking second dose of vaccine, general diseased and certain age people showed oral symptoms. Facial symptoms were observed after administration of first dose of vaccine in general diseased subjects. However, subject with autoimmune pathologies showed probability of facial symptoms after taking both first and second dose of vaccine. As a result, it was determined that there is no link between COVID-19 vaccination injection and face and oral symptoms.(8)

SARS-CoV-2 mRNA immunisation induces significant humoral immunity in nursing and

pregnant women, according to a recent study. There was also indication of antibodies being transferred to the infant. However, further research into the effects of the covid-19 vaccine on maternal and foetal outcomes is still required.(9)

In human therapeutics, IgG plays a critical role. Maximum mammals employ immunoglobulin as a means of passive immunity. IgG is a key component of the immune system and is found in milk and colostrum. Microfiltered immunological raw milk, commonly known as colostrum from cows vaccinated against SARSCoV2, is believed to provide short-term protection against coronavirus infection in humans.(10)

Since antiviral therapy is not available, thus stimulating immunity by COVID-19 vaccines is the best option to prevent future infections. Vaccines have potential to control COVID-19 infections (11)

A body temperature (fever) test is the primary test used to detect infection at some countries' borders, and concerns about its efficacy have been raised. A recent study found that such screening approaches are ineffective among hospitalised patients; however, data for young adults, who frequently come with moderate or asymptomatic illness, is limited. Importantly, this is the segment of the population that is deemed highly infectious. They are also the demographic group most likely to travel with temperature measurements being implemented at airports around the world.(12)

The World Health Organization (WHO) has declared the 2019 COVID-19 pandemic a public health emergency of international concern. Previous studies of the 1918 influenza pandemic hypothesized that vitamin D had a nonclassical role in reducing severe pneumonia and mortality. Recent clinical trials have also revealed that vitamin D administration reduces the development of acute respiratory infections as well as the persistence of respiratory tract illnesses in both adults and children. We'll focus on three critical issues here: 1. Vitamin D may assist in the prevention of SARS-CoV-2 infection: Vitamin D metabolism and regulation in the kidneys and extra-renal systems Vitamin D: An overview of molecular processes and multifunctional roles beyond skeletal homeostasis. A review of vitamin D-mediated local immunomodulation in human infectious diseases. Infection caused by an antiviral agent. There are anti-malaria and anti-systemic lupus erythematosus drugs on the market (SLE). 2. Vitamin D, which acts as a potent immunosuppressant, may diminish the cytokine release syndrome in COVID-19: Vitamin D suppresses critical pro-inflammatory pathways such as nuclear factor kappa B (NF-kB), interleukin-6 (IL-6) and tumour necrosis factor (TNF-) (TNF). 3. Vitamin D may protect COVID-19 neurons from harm by increasing the expression

of neurotrophins such as Nerve Growth Factor (NGF): Vitamin D induces the production of a significant neurotrophic factor.(13)

Outbound mobility was positively associated with COVID-19 infection at lag levels of 1, 2, and 3 weeks, but solar exposure was negatively associated. The coefficients of the interaction items suggested that sun radiation reduced the relationship between outward mobility and the number of daily new confirmed cases at 2- and 3-week lag levels. The moderating impact, however, was minor, and the favourable effect of outward mobility on COVID-19 infection was not eradicated. As a consequence, our data demonstrated that solar radiation only significantly attenuated the relationship between human mobility and COVID-19 infection, emphasising that movement should still be restricted on sunny days during the COVID-19 pandemic. Some observational research and clinical trials revealed that vitamin D supplementation reduced the risk of influenza, whereas others did not. The outbreak occurred during the winter, when 25-hydroxyvitamin D (25(OH)D) concentrations are at their lowest; the number of cases in the Southern Hemisphere near the end of summer is low; vitamin D deficiency has been linked to acute respiratory distress syndrome; and case-fatality rates rise with age and chronic disease comorbidity, both of which are linked to COVID-19. People at risk of influenza and/or COVID-19 might consider taking 10,000 IU/d vitamin D3 for a few weeks to rapidly raise 25(OH)D levels, followed by 5000 IU/d to reduce the risk of infection. The objective is to raise 25(OH)D levels to 40-60 ng/mL (100-150 nmol/L). Increased vitamin D3 dosages may be effective in the treatment of COVID-19 infections. Randomized controlled trials and large population studies should be carried out to examine these recommendations. (14)

Asthmatics are considered a high-risk category for severe disease from the fatal corona virus, according to current WHO guidelines. Viruses are a significant cause of asthma flare-ups, and the current corona virus epidemic has raised various questions about the best therapy strategies. The debate over whether systemic corticosteroids should be used in the normal therapy of COVID-19-related asthma exacerbations is addressed here. According to recent WHO investigations, corticosteroids should not be used if COVID-19 is suspected due to distresses since these drugs may reduce protective innate antiviral immune responses. This may not be acceptable in the event of asthma exacerbation, which is a state linked with heightened type 2 inflammation, a disease characteristic known to directly decrease antiviral immunity. Corticosteroids are expected to restore compromised antiviral immunity in

asthmatics through their suppressive effects on type 2 inflammation, and, in contrast to non-asthmatic patients, show beneficial clinical effects in the context of SARS-CoV-2 infection.

CTLs (cytotoxic T-lymphocytes) play an important role in developing protective immunity against infections and enmities. As a vaccine delivery platform, moderately oxidised acetalated dextran nanoparticles (Ox-AcDEX NPs) with an average diameter of 100 nm are developed for the first time. Ox-AcDEX NPs are imine bond formed with a representative CTL peptide epitope (CTLp) from human mucin-1 (MUC1) with the sequence TSAPDTRPAP (referred to as Mp1) and an immune-increasing adjuvant R837 (referred to as R) to yield AcDEX-(imine)-Mp1-R NPs that can be used to develop effective anticancer vaccines. AcDEX (immigrant) Mp1R NPs induce potent and sustained immune responses against MUC1 CTLs in mice to improve tumor prophylaxis. To demonstrate its versatility, this nanoplatform is also being used to deliver the Coronavirus 2 Severe Acute Respiratory Syndrome (SARSCoV2) epitope to prevent 2019 (COVID19) coronavirus disease. AcDEX (immigrant) SpR NPs for SARSCoV2 vaccination candidates are generated by splicing OxAcDEX NPs to a putative SARSCoV2 CTL epitope (designated Sp) and R837.. Several epitopes are being researched and addressed as possible factors to the production of strong and protective anti-SARS-CoV-2 CTL responses. These findings provide insight on the extensive usage of Ox-AcDEX NPs for the delivery of tumor- and virus-related epitopes. (15-19)

#### DISCUSSION:

Term 'Vaccine effectiveness' refers to a measure of how vaccine works in real world. Vaccines with two doses provide only partial protection after the first dosage, and the second dose boosts that protection.

It was found that haematological patients are more prone to COVID-19 infection in terms of mortality and morbidity. Also it was found that COVID-19 cancer infected patients who had undergone anti-cancer treatments had low risk of mortality and exacerbation. Novel vaccine technology used in covid vaccine has potential to cure 95% of the infection. A survey concluded that in normal conditions of infection, there is no difference in concentration of IgG antibodies in males and females. Effectiveness of COVID-19 vaccine depends upon how subject's body respond.

Vaccines are the best possible options at present. It has the potential to fight COVID-19 infection.

In this review, we move one step closer to this goal by elucidating the function of extrarenal vitamin D metabolism in the prevention and treatment of SARSCoV2 infection. By higher information the protecting antibody reputation process, researchers can create vaccines and pills to deal with SARSCoV2. We offer 91030, a monoclonal antibody that goals the SARSCoV2 receptor binding web website online for ACE2 as a part of a publicly to be had antibody reaction encoded via way of means of the IGHV353/IGHV366 gene. The reputation houses of 91030 magnificence and comparable antibodies have been sequenced and structurally analyzed to decide how they relate to neutralization of SARSCoV2. We generated a moderate/mild collection signature and determined that antibody precursors have been found in 1 in 44,000 human B cells, constant with trendy antibody availability in lots of COVID19 patients. These magnificence signatures offer genetic, structural and practical immunological factors which can assist boost up SARSCoV2 antibody therapy.

All COVID-19 vaccines permitted for emergency use via way of means of the World Health Organization (WHO) have passed through randomized medical trials to make certain quality, protection and efficacy. A vaccine have to be as excessive as 50% or extra powerful to be permitted. Once permitted, they're monitored to make certain they're secure and powerful for life.

#### CONCLUSION:

There is want for using intravenous immunoglobulin (IVIG) and passive immunotherapy in big managed scientific trials.

Cancer patients who had gone anticancer treatment using various therapies like immunotherapy, surgery are not associated with COVID-19 infection. The role of anticancer therapy in COVID-19 patients need further study and findings.

Immunotherapy is effective in curing COVID-19 infections on large extent. But in some cases it was found that immunotherapy had certain ill effects, especially in elderly patients.

Vaccines are the best possible immunotherapy available and has potential to fight COVID-19.

**COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

#### REFERENCES:

1. Karthik K, Senthilkumar TMA, Udhayavel S, Raj GD. Role of antibody-dependent enhancement (ADE) in the virulence of SARS-CoV-2 and its mitigation strategies for the development of vaccines and immunotherapies to counter COVID-19. *Hum Vaccin Immunother.* 2020 Dec 1;16(12):3055–60.
2. Fox TA, Troy-Barnes E, Kirkwood AA, Chan WY, Day JW, Chavda SJ, et al. Clinical outcomes and risk factors for severe COVID-19 in patients with haematological disorders receiving chemo- or immunotherapy. *Br J Haematol.* 2020 Oct;191(2):194–206.
3. Dhama K, Patel SK, Natesan S, Vora KS, Iqbal Yattoo M, Tiwari R, et al. COVID-19 in the elderly people and advances in vaccination approaches. *Hum Vaccin Immunother.* 2020 Dec 1;16(12):2938–43.
4. Wang B, Huang Y. Immunotherapy or other anti-cancer treatments and risk of exacerbation and mortality in cancer patients with COVID-19: a systematic review and meta-analysis. *Oncoimmunology.* 2020 Sep 22;9(1):1824646.
5. COVID-19 Vaccination in Patients with Autoimmune Inflammatory Rheumatic Diseases: Clinical Guidance of the Korean College of Rheumatology - PubMed [Internet]. [cited 2021 Oct 27]. Available from: <https://pubmed.ncbi.nlm.nih.gov/33783147/>
6. Immune reactivity during COVID-19: Implications for treatment - PubMed [Internet]. [cited 2021 Oct 27]. Available from: <https://pubmed.ncbi.nlm.nih.gov/33421440/>
7. A comparison study of SARS-CoV-2 IgG antibody between male and female COVID-19 patients: A possible reason underlying different outcome between sex - PubMed

- [Internet]. [cited 2021 Oct 27]. Available from: <https://pubmed.ncbi.nlm.nih.gov/32383183/>
8. Facial and Oral Manifestations Following COVID-19 Vaccination: A Survey-Based Study and a First Perspective - PubMed [Internet]. [cited 2021 Oct 27]. Available from: <https://pubmed.ncbi.nlm.nih.gov/34066995/>
  9. Vaccination against COVID-19 infection: the need of evidence for diabetic and obese pregnant women - PubMed [Internet]. [cited 2021 Oct 27]. Available from: <https://pubmed.ncbi.nlm.nih.gov/34181082/>
  10. Jawhara S. Can Drinking Microfiltered Raw Immune Milk From Cows Immunized Against SARS-CoV-2 Provide Short-Term Protection Against COVID-19? *Front Immunol.* 2020;11:1888.
  11. Recent advances in vaccine and immunotherapy for COVID-19 - PubMed [Internet]. [cited 2021 Oct 27]. Available from: <https://pubmed.ncbi.nlm.nih.gov/33156739/>
  12. Mitra B, Luckhoff C, Mitchell RD, O'Reilly GM, Smit DV, Cameron PA. Temperature screening has negligible value for control of COVID - 19. *Emergency Medicine Australasia.* 2020 Aug 17;32(5):867–9.
  13. The importance of vitamin d metabolism as a potential prophylactic, immunoregulatory and neuroprotective treatment for COVID-19 - PubMed [Internet]. [cited 2021 Oct 27]. Available from: <https://pubmed.ncbi.nlm.nih.gov/32847594/>
  14. Evidence that Vitamin D Supplementation Could Reduce Risk of Influenza and COVID-19 Infections and Deaths - PubMed [Internet]. [cited 2021 Oct 27]. Available from: <https://pubmed.ncbi.nlm.nih.gov/32252338/>
  15. Developing Acid-Responsive Glyco-Nanoplatform Based Vaccines for Enhanced Cytotoxic T-lymphocyte Responses Against Cancer and SARS-CoV-2 - PubMed [Internet]. [cited 2021 Oct 27]. Available from: <https://pubmed.ncbi.nlm.nih.gov/34512228/>
  16. Burhani, Tasneem Sajjad, and Waqar M. Naqvi. "Telehealth - A Boon in the Time of COVID 19 Outbreak." *JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS* 9, no. 29 (July 20, 2020): 2081–84. <https://doi.org/10.14260/jemds/2020/454>.

17. Gawai, Jaya Pranoykumar, Seema Singh, Vaishali Deoraaji Taksande, Tessy Sebastian, Pooja Kasturkar, and Ruchira Shrikant Ankar. "Critical Review on Impact of COVID 19 and Mental Health." JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS 9, no. 30 (July 27, 2020): 2158–63. <https://doi.org/10.14260/jemds/2020/470>.
18. Naqvi, Waqar M., and Arti Sahu. "Paradigmatic Shift in the Education System in a Time of COVID 19." JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS 9, no. 27 (July 6, 2020): 1974–76. <https://doi.org/10.14260/jemds/2020/430>.
19. Wanjari, A. K., Ayush Dubey, Sourav Chaturvedi, and Sunil Kumar. "Young COVID 19 Presenting as Fatal Subarachnoid Hemorrhage: Association or Chance?" MEDICAL SCIENCE 24, no. 104 (August 2020): 2712–15.

UNDER PEER REVIEW