

### **Review on Moderate to Severe Asthma in Primary Care, Review Article**

#### **Abstract:**

Asthma is a chronic condition characterized by wheezing, coughing, and shortness of breath due to airway inflammation and hyper-responsiveness. Severe asthma accounts for a considerable amount of asthma-related costs, although being less common than milder asthma. According to a review of US studies, the expenses for people with severe asthma are roughly 1.7- to 5-fold higher than for people with mild asthma. A subspecialized severe asthma services provides the benefit of an organized, variety of approaches to validate the diagnosis, asthma severity and phenotype, and risk factors and comorbidities management. This multimodal approach frequently comprises a team of respiratory physicians, nurses, and support health specialists, such as physiotherapists, speech therapists, nutritionists, and clinical psychologists. In addition to evaluation and monitoring for common comorbidities, they provide physician review, lung function testing, blood tests, inhaler optimization, and general illness awareness. This review aims to overview approach to diagnosis and management of moderate to severe asthma in primary care settings.

#### **Introduction**

Asthma is a chronic condition characterized by wheezing, coughing, and shortness of breath due to airway inflammation and hyper-responsiveness. The term "asthma" refers to a group of illnesses that have similar symptoms but differ in their underlying etiology and prognosis. Depending on the age of onset, clinical signs might indicate one of several phenotypes of the disease, each with its own set of diagnostic, management, and therapy problems. It is hoped that a better understanding of the various subtypes of asthma will help us better evaluate, manage, and cure the symptoms.[1-3]

Asthma is often assumed to be an illness that starts in childhood. Despite the fact that asthma is most typically diagnosed in children, it can manifest clinically at any age. Indeed, according to a national survey, 3.1 percent each year who diagnosed with asthma are over 65 years old which is not significantly different from those between

the ages of 18 and 34 where 4.0 percent are diagnosed per year. Asthma occurrence in adults over 65 years of age is reported to be 7%, which is comparable to the total prevalence. Prematurity, early lung infections, sinusitis, tobacco consumption, and obesity are linked to diagnosis of asthma in adults. As a result, an older patient's beginning of chronic cough should not prevent the physician from examining asthma.[4-7]

According to the National Health Interview Survey–2012, roughly 40 million people in the USA have had asthma at some point in their lives, this comprises about 13 % of the total population .and 8% are currently having asthma which includes about 26 million patient . Incidences for young adults aged 18-24 years are higher (10.3 percent) when compared to older people. 5% to 10% of the entire adult asthma populations are diagnosed with severe asthma.[8,9] Severe asthma accounts for a considerable amount of asthma-related costs, although being less common than milder asthma. According to a review of US studies, the expenses for people with severe asthma are roughly 1.7- to 5-fold higher than for people with mild asthma.[10-12]. Asthma is substantially more common in women (10.4%) than in men (6.2%), in those living in poverty (11.8%), and in those who identify as an ethnic or racial minority, particularly black race (10.2%) and Puerto Rican Hispanic ethnicity (14.9 percent ). The incidence varies greatly per state, ranging from 4.9 percent to 12.7 percent. Despite the availability of a wide range of treatment options, over half of individuals with asthma report having one or more attacks in the previous year, emphasising the significance of symptom management and disease control.[13]

Severe asthma is a diverse condition that is described as a treatment-resistant disease characterized by persistent symptoms or recurrent acute episodes despite the use of standard medications. It means patients with "difficult-to-treat" asthma, in which a variety of factors contribute to poor management of the disease , as well as patients with "biologically severe" asthma, who remain to have poor control yet after alternative diagnoses have been ruled out and contributory factors have been optimized. There are now successful, focused biological medicines aimed at targeting the underlying pathogenic pathways for some people with physiologically severe asthma. Many individuals with difficult-to-treat asthma, on the other hand, have risk factors that exacerbate symptoms, and by carefully managing these variables, they can achieve significant enhancement. As a result, it's critical to establish care models that best address the complex and varied needs of people with severe asthma, with the overall goal of enhancing patient outcomes for all.[14-19]

The vast number of asthma patients is treated by primary care physicians, placing them at the forefront of asthma management, allowing them to identify at-risk patients and give appropriate medication and awareness. In a 2012 survey, reported that just 22% of asthma patients were managed by a specialist on a regular basis, and 48% of patients had never seen a specialist. As a result, it's critical for healthcare professionals to conduct routine asthma management evaluations and ask patients the right questions in order to keep track of asthma severity. Patients with asthma should be referred to specialists by primary care providers who do not do pulmonary function testing on a regular basis.[20-23] According to an Australian survey, up to 45 percent of community members had uncontrolled asthma, resulting in frequent hospitalizations and unscheduled medical visits. However, just half of the patients had seen a general practitioner in the previous year, and only 10% had had a specialist evaluation.[24]

A subspecialized severe asthma services provides the benefit of an organized, variety of approaches to validate the diagnosis, asthma severity and phenotype, and risk factors and comorbidities management. This multimodal approach frequently comprises a team of respiratory physicians, nurses, and support health specialists, such as physiotherapists, speech therapists, nutritionists, and clinical psychologists. In addition to evaluation and monitoring for common comorbidities, they provide physician review, lung function testing, blood tests, inhaler optimization, and general illness awareness. It expands educational and self-management opportunities, as well as access to modern biological medicines. These services are now encouraged as they showed positive results on improving the symptoms of the disease, quality of life and decreased incidences of disease exaggeration [25-29]

### **Definition and pathogenesis of severe asthma**

Historically, asthma was thought to be a condition caused by increase in the activity of the airway smooth muscle, or bronchoconstriction, in reaction to environmental stimuli. Even in people with just occasional symptoms or new-onset asthma, chronic airway inflammation is now regarded a characteristic of the condition. The incidence of symptoms, the extent of functional disability, and the rate of asthma attacks were all used to determine the severity of asthma. More recent practice guidelines reflect increased knowledge of the basic pathophysiology of asthma, so the previous factors are considered only as elements of asthma management, and the severity of asthma disease is categorized by the type and dosage of medicines a patient needs to

maintain proper disease control. As a result, symptom control is evaluated separately from and then incorporated into the criteria of asthma severity.[30,31].

The International European Respiratory Society (ERS) and the American Thoracic Society (ATS) updated the concept of severe asthma in 2014. Severe asthma in people aged 6 and up is defined by the ERS/ATS guidelines as either (1) asthma that needs treatment with medium- to high-dose inhaled corticosteroids (ICS) and one or more additional medicines (eg, long-acting 2-agonist, , leukotriene modifier, and theophylline) or systemic corticosteroids for minimum 6 months According to these criteria, uncontrolled asthma requires at least one of the following symptoms: unsatisfactory symptom control, recurrent acute exacerbations, major exacerbation (requiring hospitalization), or airflow limitation.[32] .Although asthma has a bigger influence on physical than mental functioning, poorly controlled asthma is linked to poor school performance, attention, and focus, as well as an increase in depression and anxiety symptoms. Furthermore, airway remodeling can result in airway functional impairment and the evolution of chronic obstructive pulmonary disease (COPD). Taking oral corticosteroids to treat severe uncontrolled asthma has its own set of side effects, including osteoporosis, slowed growth in children, and increased risk of cataract and inflammation [30,31,33,34]

## **Assessment and evaluation of Asthma**

The combination of patient symptoms and respiratory function testing is required for a reliable diagnosis of asthma. As the symptoms of asthma are sometimes ambiguous and can be triggered by other diseases, it's crucial to rule out disorders that resemble asthma, especially in older people who are more subjected to have other illnesses.

### **History:**

Asthma symptoms include wheezing, tightness of the chest, and coughing. These symptoms are frequently recurrent in nature and can vary in strength. Changes in stimulators such as allergens, irritants, or respiratory illnesses are related to changes in symptoms. Bronchoconstriction in reaction to exercise can cause symptoms of asthma. At rest, people with uncontrolled illness may experience persistent symptoms. The presence of chest tightness as the primary presenting symptom should raise suspicions of heart illness. Asthmatics frequently suffer from rhinitis and sinusitis, where allergic rhinitis is a predisposing factor for developing asthma. Upper airway symptoms should be assessed, as they are regarded to represent diverse presentations of a shared allergic pathogenesis.[35,36]

Any possible exposures that aggravate the patient's respiratory problems should be tested by physician. The presence of identifiable triggers raises the likelihood of underlying asthma. Allergens are well-known triggers that might be seasonal (usually outdoors) or persistent (typically indoor). Outdoor allergen exposure is widely different across the United States in terms of distribution and timing. Outdoor allergies are primarily pollen-based and can come from a variety of sources, including trees, and weeds, and can be detected at different periods throughout the year. Dust mites, cockroaches, molds, and animal dander from pets are examples of perennial allergies. Cigarette smoking, secondhand smoke exposure, perfumes or strong scents, excess heat or cold, exercise, or psychosocial strain are among non-allergic triggers. Occupational exposures can cause the onset of asthma or exacerbate the symptoms of asthma that already exist. It's crucial to ask about employment exposures in relation to asthma diagnosis and symptom variation by asking about changes in symptoms between weekdays and weekends. Environmental sensitivity is suggested by changes in respiratory symptoms while travel, which supports an asthma diagnosis. [37-40]

### **Physical Examination**

The physical examination is most helpful in determining whether or not there are any concomitant or mimicking conditions. A pulmonary examination in an asthmatic patient is frequently unremarkable. Expiratory wheezing is possible; however it is neither sensitive nor unique to this disease. Inspiratory wheeze is unusual and could indicate a different or additional condition. Crackles, on the other hand, should prompt you to examine other possibilities. Patients may exhibit symptoms of rhinitis or postnasal drip. Eczema can be discovered with a skin examination. Finally, to check for evidence of heart failure, a cardiac examination should be conducted.

### **Tests**

Spirometry is an essential diagnostic tool used when both the history of the patient and its physical evaluation point to asthma as a potential diagnosis. Spirometry should be performed before and after administration of bronchodilator to check for the two major criteria for asthma diagnosis: expiratory airflow obstruction and airflow variability.

Spirometry that shows both airflow obstruction and complete reversal of airflow obstruction after bronchodilator delivery confirms an asthma diagnosis. Spirometry is typically normal when asthma is adequately treated since variability in symptoms and airflow limitation is a significant aspect of asthma. Additionally, due to increased disease symptoms during presentation, some individuals with asthma who present with expiratory airflow obstruction may not entirely reverse following bronchodilator administration. These other illnesses should be examined because partial reversibility is also a characteristic of COPD or asthma-COPD overlap. In such cases, spirometry is unable to reliably differentiate between asthma and COPD.

A ratio of fractional exhaled volume in the first second (FEV<sub>1</sub>) to total volume forcefully expelled (FVC) less than the lower limit of normal (LLN) indicates the presence of airflow blockage. Because the LLN compensates for the projected fall in FEV<sub>1</sub>/FVC that happens

with ageing, it is preferable over adopting a set threshold (e.g., 70%). The term "post-bronchodilator response" refers to an increase in FEV1 or FVC of more than 12% and more than 200 mL after using a bronchodilator. These limits are set by guidelines that emphasize that the clinical situation should be considered when evaluating test results. For example, a patient who improves their FEV1 by 10% after a bronchodilator and also improves their FEV1 before the bronchodilator following a trial of inhaled corticosteroids (ICS) would have had a clinically meaningful response that is highly compatible with a diagnosis of asthma.[41-43]

When spirometry indicates normal results but physician still consider the diagnosis of asthma, spirometry must be repeated at a later time point, because minimally single episode of obstruction must be documented to confirm the conclusion of asthma. If reversibility on spirometry cannot be determined, sequential testing with a peak flow meter can be used to establish variability in airflow limitation. Over the course of two weeks, the patient is told to record the best of three attempted peak flows twice daily (usually in the morning and late evening) or more frequently during periods of respiratory discomfort. Variable airflow limitation can be diagnosed by a 7-day average of each day's largest to smallest recorded value, divided by the day's average.[42]

Bronchoprovocation tests, such as a methacholine challenge, for measuring the hyper-responsiveness of airways has a poor sensitivity for asthma, as other respiratory disorders, such as COPD, and the general population, where the prevalence has ranged from 4% to 37%, have been linked to airway hyper-responsiveness. As a result, this test is no longer used to verify asthma, although it does have a function in excluding asthma in some groups due to its high negative predictive value, which has been observed to approach 100%.[44-46]

A progressive diagnostic strategy involving spirometry or peak flow testing to determine blockage and variable airflow limitation is used mainly unless the patient shows exaggerated symptoms on administration. When patients are already using asthma drugs, identifying asthma-related blockage might be difficult. According to a recent Canadian study, up to one-third of those who have been diagnosed with asthma by a doctor do not actually have it. These patients were less likely to have received formal testing for airflow limitation, and the outcomes of the study highlight the importance of objective testing in asthma diagnosis.[47]

In most cases, radiographic studies and blood tests are not required in the diagnostic process until there is a suspicion of a different diagnosis. Elevations in eosinophils, immunoglobulin E (IgE), or allergen-specific IgE, while useful in detecting allergic illness and evaluating advanced therapy, are neither sensitive nor specific for the initial diagnosis of asthma. Although fractional exhaled nitric oxide (FeNO) is a sign of eosinophilic airway inflammation, it is rarely used in asthma diagnosis.

## **Management:**

To enhance disease control and address underlying inflammation while reducing side effects from prescribed drugs, successful care necessitates ongoing review over time. Symptoms, risk of worsening, drug tolerance and adherence, and comorbidities should all be assessed at each appointment.

Respiratory function, symptom recurrence, and frequency of exacerbations can all be used to determine the severity of asthma before starting medical treatment. Severity is defined as frequent, mild - to - moderate persistent or severe persistent, and can guide first therapy decisions. Mild intermittent symptomatology is no longer advised since it suggests that other severities cannot be symptomatic. Importantly, asthma severity is not a static attribute and should be reclassified at each visit based on the quantity of medication required to manage or relieve asthma symptoms. Changes in asthma severity might signal new environmental exposures, comorbidities, or disease progression.

### **Non-pharmacological Managements:**

Asthma care relies heavily on avoiding triggers. Common triggers and techniques for dealing with them are listed below:

- Ambient air pollution :Remain indoors during poor air quality days
- Certain foods :Test for food-specific allergies, avoidance
- Cigarette smoke :Smoking cessation assistance, home smoking ban
- Cockroaches: Sweep and vacuum regularly, use roach traps.

During periods of high pollen counts or poor outdoor air quality, patients with sensitivity to air pollution are advised to stay home. Pollen avoidance should be timed according to each patient's sensitivity profile, which should be established if the patient has a strong history of seasonal allergies and necessitates a referral to an allergist. The importance of avoiding secondhand smoke and quitting smoking is underlined. Indoor allergen avoidance, focusing on specific allergens to which the patient is allergic, is recommended but difficult to put into practice due to the difficulty of attaining total allergy remediation and the high cost of therapies.[48-50]. Overweight or obese patients should be counseled on weight loss as part of general health maintenance, and all patients should be urged to eat a balanced diet. Weight loss of 10 to 15 kgs has been shown to enhance asthma severity.[51]

### **Pharmacological Managements:**

### **Medication for controlling asthma:**

The main goal of such medications is to block the inflammatory process that causes asthma and thus preventing any non-reversible airway remodeling. The backbone of controller therapy is inhaled corticosteroids. Oral thrush, which can be alleviated by using a spacer device and cleaning the mouth after ICS use, and dysphonia, which can be managed by switching to a different delivery system, are two common side effects. If control is inadequate, long-acting beta agonists (LABA) or leukotriene modifiers can be added to ICS, with the former being more successful than the latter. Long-acting muscarinic antagonists, which are routinely used to treat COPD, do not appear to be superior to LABA and are normally reserved for patients with severe disease. LABA mono-therapy is unsuitable and should never be given for a patient with asthma, as it was associated with an elevated asthma-related mortalities.[52-56]

### **Rescue Medications:**

All patients should be provided with an albuterol rescue inhaler. In individuals with acute symptoms, this short-acting drug delivers rapid bronchodilator and can be used up to four times per day. Albuterol administered by metered-dose inhaler (MDI) without spacer, MDI with spacer, or nebulizer is equally effective in an adult patient with appropriate guidance. In practice, though, spacers and nebulizers are more reliable at delivering medication. Patients with episodic asthma may be able to manage their symptoms with just a short-acting beta-agonist; however, those with chronic asthma (those who use their albuterol inhaler more than twice weekly to relieve symptoms) will require the addition of an inhaled corticosteroid (ICS).[57,58]

### **Other Treatment Options in Severe Asthma:**

A typical target is the allergic inflammatory pathway. Omalizumab is an anti-IgE antibody that interacts with mast cells, eosinophils, and basophils in the allergic pathway. It's approved for people with mild to severe asthma who don't respond to ICS and have confirmed allergen sensitivity. It's given every 2 to 4 weeks. Antibodies such as mepolizumab, reslizumab, and benralizumab target the interleukin-5 pathway, which is implicated in eosinophil recruitment and activation.[59-62] Bronchial thermoplasty (BT) is an endoscopic technique that uses radiofrequency heat radiation to ablate airway smooth muscle, diminishing its ability to cause bronchoconstriction. In people with severe asthma, data shows that BT lowers exacerbations, minimizes asthma-related health-care utilization, and enhances the quality of life. [63,64]

**Conclusion:**

Asthma is one of the most known respiratory disorders that require long-term managements. Diagnosis of asthma is complicated by the common features between asthma and other respiratory diseases. Severe asthma is considered when the patient is not responding to the normal medication, where inhaled corticosteroids are the main management technique beside avoidance of the trigger that causes case exacerbation. Patient awareness about the disease and adequate methods for its management is an important step in the management of this disorder.

UNDER PEER REVIEW

## References:

1. Villarroel MA, Blackwell DL. Percentage of adults aged 18–64 years with current asthma,\* by State-National Health Interview Survey,(Dagger) 2014–2016. Atlanta (GA): Centers Disease Control 1600 Clifton Rd; 2018. 30333 USA.
2. Pavord ID, Beasley R, Agusti A, Anderson GP, Bel E, Brusselle G, et al. After asthma: redefining airways diseases. *Lancet*. (London, England). (2018) 391:350–400. doi: 10.1016/S0140-6736(17)30879-6
3. Haldar P, Pavord ID, Shaw DE, Berry MA, Thomas M, Brightling CE, et al. Cluster analysis and clinical asthma phenotypes. *Am J Respirat Critic Care Med*. (2008) 178:218–24. doi: 10.1164/rccm.200711-1754OC.
4. Winer RA, Qin X, Harrington T, et al. Asthma incidence among children and adults: findings from the behavioral risk factor surveillance system asthma call-back survey—United States, 2006–2008. *J Asthma* 2012;49(1):16–22. [PubMed: 22236442]
5. Huovinen E, Kaprio J, Koskenvuo M. Factors associated to lifestyle and risk of adult onset asthma. *Respir Med* 2003;97(3):273–80. [PubMed: 12645835]
6. Guerra S, Sherrill DL, Martinez FD, et al. Rhinitis as an independent risk factor for adult-onset asthma. *J Allergy Clin Immunol* 2002;109(3):419–25. [PubMed: 11897985]
7. Jamrozik E, Knuiiman MW, James A, et al. Risk factors for adult-onset asthma: a 14-year longitudinal study. *Respirology* 2009;14(6):814–21. [PubMed: 19703063].
8. Centers for Disease Control and Prevention. 2012 National Health Interview Survey (NHIS) data. Available at: <https://www.cdc.gov/asthma/nhis/2012/data.htm>. Accessed May 24, 2017.
9. Centers for Disease Control and Prevention. Asthma Facts—CDC’s National Asthma Control Program Grantees. Atlanta, Ga: US Department of Health and Human Services, Centers for Disease Control and Prevention; 2013.
10. Chung KF, Wenzel SE, Brozek JL, et al. International ERS/ATS guidelines on definition, evaluation and treatment of severe asthma. *Eur Respir J*. 2014;43:343-373.
11. Nunes C, Pereira AM, Morais-Almeida M. Asthma costs and social impact. *Asthma Res Pract*. 2017;3:1.
12. Bahadori K, Doyle-Waters MM, Marra C, et al. Economic burden of asthma: a systematic review. *BMC Pulm Med*. 2009;9:24.
13. Mazurek JM. Prevalence of asthma, asthma attacks, and Emergency Department visits for asthma among working adults — national health interview survey, 2011 — 2016. *MMWR Morb Mortal Wkly Rep* 2018;67 10.15585/mmwr.mm6713a1.
14. 7 Chung KF, Wenzel SE, Brozek JL, et al. International ERS/ ATS guidelines on definition, evaluation and treatment of severe asthma. *Eur Respir J* 2014; 43: 343-373.
15. 8 Bel EH, ten Brinke A. New anti-eosinophil drugs for asthma and COPD. *Chest* 2017; 152: 1276-1282.

16. 9 Grainge CL, Maltby S, Gibson PG, et al. Targeted therapeutics for severe refractory asthma: monoclonal antibodies. *Expert Rev Clin Pharmacol* 2016; 9: 927-941.
17. 10 Upham JW, Chung LP. Optimising treatment for severe asthma. *Med J Aust* 209 (2 Suppl): S22-S27.
18. 11 Bel EH, Sousa A, Fleming L, et al; Unbiased Biomarkers for the Prediction of Respiratory Disease Outcome (U-BIOPRED) Consortium. Diagnosis and definition of severe refractory asthma: an international consensus statement from the Innovative Medicine Initiative (IMI). *Thorax* 2011; 66: 910-917.
19. 12 Agustí A, Bafadhel M, Beasley R, et al. Precision medicine in airway diseases: moving to clinical practice. *Eur Respir J* 2017; 50: 1701655.
20. Price D, Bjermer L, Bergin DA, Martinez R. Asthma referrals: a key component of asthma management that needs to be addressed. *J Asthma Allergy*. 2017;10:209-223.
21. Rance KS. Helping patients attain and maintain asthma control: reviewing the role of the nurse practitioner. *J Multidiscip Healthc*. 2011;4:299-309.
22. Murphy KR, Meltzer EO, Blaiss MS, Nathan RA, Stoloff SW, Doherty DE. Asthma management and control in the United States: results of the 2009 Asthma Insight and Management survey. *Allergy Asthma Proc*. 2012;33:54-64.
23. Schierhorn C. Avoiding pitfalls in caring for patients with asthma. Available at: <https://thedo.osteopathic.org/2014/03/avoiding-pitfalls-in-caring-for-patients-with-asthma>. Accessed September 26, 2017. 2014.
24. Reddel HK, Sawyer SM, Everett PW, et al. Asthma control in Australia: a cross-sectional web-based survey in a nationally representative population. *Med J Aust* 2015; 202: 492-496. <https://www.mja.com.au/journal/2015/202/9/asthma-control-australia-cross-sectional-web-based-survey-nationally>.
25. Gibson PG, McDonald VM. Management of severe asthma: targeting the airways, comorbidities and risk factors. *Intern Med J* 2017; 47: 623-631.
26. Centre of Excellence in Severe Asthma [website]. 2017 [www.severeasthma.org.au](http://www.severeasthma.org.au) (viewed May 2018).
27. Tay TR, Lee J, Radhakrishna N, et al. A structured approach to specialist-referred difficult asthma patients improves control of comorbidities and enhances asthma outcomes. *J Allergy Clin Immunol Pract* 2017; 5: 956-964.
28. van der Meer A-N, Pasma H, Kempenaar-Okkema W, et al. A 1-day visit in a severe asthma centre: effect on asthma control, quality of life and healthcare use. *Eur Respir J* 2016; 48: 726-733.
29. Gibeon D, Heaney LG, Brightling CE, et al. Dedicated Severe asthma services improve health-care use and quality of life. *Chest* 2015; 148: 870-876.
30. Global Initiative for Asthma. Global strategy for asthma management and prevention. Available at: <http://www.ginasthma.org>. Accessed May 24, 2017.
31. National Heart, Lung and Blood Institute. National asthma education and prevention program. In: Expert Panel Report 3: Guidelines for the Diagnosis and Management of Asthma. Full Report 2007. Bethesda, Md:NHLBI Health Information Center; 2007.
32. Chung KF, Wenzel SE, Brozek JL, et al. International ERS/ATS guidelines on definition, evaluation and treatment of severe asthma. *Eur Respir J*. 2014;43:343-373.

33. Cordier T, Slabaugh SL, Havens E, et al. A health plan's investigation of Healthy Days and chronic conditions. *Am J Manag Care*. 2017;23:e323-e330.
34. O'Byrne PM, Pedersen S, Schatz M, et al. The poorly explored impact of uncontrolled asthma. *Chest*. 2013;143:511-523.
35. Shen H, Hua W, Wang P, et al. A new phenotype of asthma: chest tightness as the sole presenting manifestation. *Ann Allergy Asthma Immunol* 2013;111(3):226-7. [PubMed: 23987204]
36. Dixon AE. Rhinosinusitis and asthma: the missing link. *Curr Opin Pulm Med* 2009; 15(1):19-24. [PubMed: 19077701]
37. Vernon MK, Wiklund I, Bell JA, et al. What do we know about asthma triggers? a review of the literature. *J Asthma* 2012;49(10):991-8. [PubMed: 23574397]
38. Henneberger PK, Redlich CA, Callahan DB, et al. An official American Thoracic Society statement: work-exacerbated asthma. *Am J Respir Crit Care Med* 2011;184(3):368-78. [PubMed: 21804122]
39. Tarlo SM, Lemiere C. Occupational asthma. *N Engl J Med* 2014;370(7):640-9. [PubMed: 24521110]
40. Work-related asthma: number of cases by classification and state, 2009-2012 Available at: [https://wwwn.cdc.gov/eworld/Data/Work-related\\_asthma\\_Number\\_of\\_cases\\_by\\_classification\\_and\\_state\\_20092012/924](https://wwwn.cdc.gov/eworld/Data/Work-related_asthma_Number_of_cases_by_classification_and_state_20092012/924). Accessed September 14, 2018.
41. Pellegrino R, Viegi G, Brusasco V, et al. Interpretative strategies for lung function tests. *Eur Respir J* 2005;26(5):948-68. [PubMed: 16264058]
42. Bateman ED, Hurd SS, Barnes PJ, et al. Global strategy for asthma management and prevention: GINA executive summary (vol 31, pg 143, 2008). *Eur Respir J* 2018;51(2):143-78.
43. Busse WW, Boushey HA, Camargo CA, et al. Expert panel report 3: guidelines for the diagnosis and management of asthma. Washington (DC): US Department of Health and Human Services, National Heart Lung and Blood Institute; 2007 p. 1-417.
44. Borak J, Lefkowitz RY. Bronchial hyperresponsiveness. *Occup Med (Lond)* 2016; 66(2):95-105. [PubMed: 26590957]
45. Cockcroft DW. Direct challenge tests: airway hyperresponsiveness in asthma: its measurement and clinical significance. *Chest* 2010;138(2, Supplement): 18S-24S. [PubMed: 20668014]
46. Cockcroft DW, Murdock KY, Berscheid BA, et al. Sensitivity and specificity of histamine PC20 determination in a random selection of young college students. *J Allergy Clin Immunol* 1992;89(1, Part 1):23-30. [PubMed: 1730837]
47. Kitch BT, Paltiel AD, Kuntz KM, et al. A single measure of FEV1 is associated with risk of asthma attacks in long-term follow-up. *Chest* 2004;126(6):1875-82. [PubMed: 15596687]
48. Gautier C, Charpin D. Environmental triggers and avoidance in the management of asthma. *J Asthma Allergy* 2017;10:47-56. [PubMed: 28331347]
49. Gøtzsche PC, Johansen HK. House dust mite control measures for asthma. *Cochrane Database Syst Rev* 2008;(2). 10.1002/14651858.CD001187.pub3.

50. Platts-Mills TAE. Allergen avoidance in the treatment of asthma and rhinitis. *N Engl J Med* 2003;349(3):207–8. [PubMed: 12867602].
51. Juel CT-B, Ali Z, Nilas L, et al. Asthma and obesity: does weight loss improve asthma control? a systematic review. *J Asthma Allergy* 2012;5:21–6. [PubMed: 22791994].
52. Halm EA, Mora P, Leventhal H. No symptoms, no asthma: the acute episodic disease belief is associated with poor self-management among inner-city adults with persistent asthma. *Chest* 2006;129(3):573–80. [PubMed: 16537854]
53. Chauhan BF, Ducharme FM. Addition to inhaled corticosteroids of long-acting beta2-agonists versus anti-leukotrienes for chronic asthma. *Cochrane Database Syst Rev* 2014;(1). 10.1002/14651858.CD003137.pub5.
54. Ducharme F, Ni Chroinin M, Greenstone I, et al. The addition of long-acting beta2-agonists to inhaled steroids compared to higher doses of inhaled steroids alone as maintenance treatment for chronic asthma. *Cochrane Database Syst Rev* 2010;(4). 10.1002/14651858.CD005533.pub2.
55. Kew K, Dahri K. Long-acting muscarinic antagonists (LAMA) added to combination long-acting beta2-agonists and inhaled corticosteroids (LABA/ICS) versus LABA/ICS for adults with asthma. *Cochrane Database Syst Rev* 2016;(1). 10.1002/14651858.CD011721.pub2.
56. Sobieraj DM, Baker WL, Nguyen E, et al. Association of inhaled corticosteroids and long-acting muscarinic antagonists with asthma control in patients with uncontrolled, persistent asthma: a systematic review and meta-analysis. *JAMA* 2018;319(14):1473–84. [PubMed: 29554174]
57. Newman KB, Milne S, Hamilton C, et al. A comparison of albuterol administered by metered-dose inhaler and spacer with albuterol by nebulizer in adults presenting to an urban emergency department with acute asthma. *Chest* 2002;121(4): 1036–41. [PubMed: 11948030]
58. Newman SP. Spacer devices for metered dose inhalers. *Clin Pharmacokinet* 2004;43(6):349–60. [PubMed: 15086274]
59. Thomson NC, Chaudhuri R. Omalizumab: clinical use for the management of asthma. *Clin Med Insights Circ Respir Pulm Med* 2012;6:27–40. [PubMed: 22745565]
60. Ortega HG, Liu MC, Pavord ID, et al. Mepolizumab treatment in patients with severe eosinophilic asthma. *N Engl J Med* 2014;371(13):1198–207. [PubMed: 25199059]
61. Castro M, Zangrilli J, Wechsler ME, et al. Reslizumab for inadequately controlled asthma with elevated blood eosinophil counts: results from two multicentre, parallel, double-blind, randomised, placebo-controlled, phase 3 trials. *Lancet Respir Med* 2015;3(5):355–66. [PubMed: 25736990]
62. FitzGerald JM, Bleecker ER, Nair P, et al. Benralizumab, an anti-interleukin-5 receptor a monoclonal antibody, as add-on treatment for patients with severe, uncontrolled, eosinophilic asthma (CALIMA): a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet* 2016;388(10056):2128–41. [PubMed: 27609406]
63. Dombret M-C, Alagha K, Boulet LP, et al. Bronchial thermoplasty: a new therapeutic option for the treatment of severe, uncontrolled asthma in adults. *Eur Respir Rev* 2014;23(134):510–8. [PubMed: 25445950]

64. Chupp G, Laviolette M, Cohn L, et al. Long-term outcomes of bronchial thermoplasty in subjects with severe asthma: a comparison of 3-year follow-up results from two prospective multicentre studies. *Eur Respir J* 2017;50(2). 10.1183/13993003.00017-2017.

UNDER PEER REVIEW