

Original Research Article

In-vitro Antioxidant and Pharmacognostic Studies of *Phaseolus vulgaris* (Linn) seed

coat

ABSTRACT

Aim: The objective of developing novel anti-epileptic herbal agent with fast therapeutic action and low toxicity profile of *Phaseolus vulgaris* (Linn) seed coat was evaluated.

Study Design: The pharmacognostic profile along with *in-vitro* estimation of bioactive compounds present were studied.

Place and Duration of Study: Department of Pharmacology, I. T. S College of Pharmacy, Muradnagar, Ghaziabad, India between May 2020 and December 2020.

Methodology: Morphology revealed dark brownish red seed, kidney to oval shaped, medium size and bland taste. Seed length, width, thickness, and surface area were also determined. Microscopically, the transverse section also showed the presence of proteinaceous aleurone cells, macro-sclereids and starch granules with irregular oval shape in the cotyledon specify the energy reservoir of seeds. In physico-chemical parameters such as extractive value ash value, moisture content, swelling index were recorded. Phytochemical screening displayed the presence of alkaloids, flavonoids, phenol, amino acid, tannins, carbohydrates and saponins. HPTLC & *in-vitro* estimations were done.

Result: HPTLC study was performed to standardize the extract of *Phaseolus vulgaris* seed coats for the presence of flavonoids. The antioxidant profile revealed TFC as 13.62 mg /g QE and TPC as 32.03 ± 1.50 mg/g GAE. IC₅₀ value for vitamin C was found to be 369.03 µg/ml as compared to *Phaseolus vulgaris* seed coat 423.00 µg/ml.

Conclusion: The study can serve as a valuable source of information and it could be beneficial in the treatment of epilepsy.

Keywords: *Phaseolus vulgaris*, pharmacognosy, antioxidant, bioflavonoids, epilepsy.

1. INTRODUCTION

Epilepsy is chronic mental disorder, generally ensuing in doubtful, indefensible recurrent seizure [1]. “Epilambanein” a greek word, which means to attack or seize is now being termed as “Epilepsy” [2]. Seizures are classified according to modified version of International Classification of Epileptic seizure (ICES) that is partial (focal) seizure, generalized seizure such as atonic, clonic, tonic-clonic (grand-mal), absence (petit-mal) seizure and unclassified seizure (neonatal, infantile) [3] Anti-epileptic drugs (AEDs) like valproic acid, phenytoin, carbamazepine, phenobarbital, clonazepam etc. are mostly used to control epileptic seizure [4].

AEDs have low efficiency, limited supply, high cost and adverse effects such as dizziness, disturbance in coordination, sedation, mood alteration, sexual dysfunction etc, as compared to herbal medicines. Herbal medicines are generally well tolerated with lesser side effects. They have a broad spectrum because they are an assortment of bioactive compounds [5].

Phaseolus vulgaris (Red kidney bean) have excellent sources of proteins, energy, carbohydrates, minerals and vitamins. It contains flavonoids such as kaempferol [6], quercetin [7], naringin [8], rutin [9] etc. These flavonoids have protective effect in epilepsy. *Phaseolus vulgaris* is used as anti-oxidant, anti-inflammatory [10], anti-diabetes [11], anti-proliferative [12] and effective in neurodegenerative disease such as anti-parkinsonism [13].

2. MATERIAL AND METHODS

2.1 Procurement and Authentication: The seeds of *Phaseolus vulgaris* were identified and procured from the local market of Modinagar, Ghaziabad. The material was authenticated by **Dr. Sunita Garg**, Emeritus Scientist, CSIR-National Institute of Science Communication and Information resources (NISCAIR), Pusa Campus, New Delhi. A voucher specimen was deposited at RHMD, New Delhi.

3. PHARMACOGNOSY

3.1 Vernacular name: Vernacular names are mentioned in Table 1 [14].

Table 1: Vernacular name of *Phaseolus vulgaris*

Languages	Names
English	Kidney bean
Hindi	Rajma
Bengali	Barbati Beej, Raajma
Telugu	Chikkuduginjalu, nallachikkudu
Kannada	Capparadavare
Oriya	Baragudi Chhuin, Rajma
Malayalam	Rajma
Tamil	Sigappu Kaaramani
Urdu	Lal lobia
Portuguese	Feijao (dry), Feijao-vagem (green)
Italian	Fagiolo, Faxoe, Faisoe (Liguria), Fasoel
Spanish	Caraota, Chaucha

3.2 Taxonomical classification: Taxonomical classification has been mentioned in Table 2.

Table 2: Taxonomical classification of *Phaseolus vulgaris*

Kingdom	Plantae
Sub-kingdom	Viridiplantae
Super-division	Embryophyta
Division	Tracheophyta
Subdivision	Spermatoptina
Class	Magnoliopsida
Order	Fabales
Family	Fabaceae
Genus	Phaseolus
Species	<i>Phaseolus vulgaris</i> L
Synonym (s)	<i>Phaseolus vulgaris</i> var. <i>humilis</i> , <i>Phaseolus</i>

3.3 Nutritional value: The seeds of *Phaseolus vulgaris* are nutritionally essential and have following crucial components shown in Table 3 [15].

Table 3: Nutritional value of *Phaseolus vulgaris*

Basic components (mg/g)		Fatty acid (mg/g)	
Total lipids	10.60	Total saturated	1.54
Protein	225.30	Total monounsaturated	0.82
Carbohydrates	612.90	Total polyunsaturated	5.86
Essential minerals (mg/g)		Vitamins (mg/g)	
Macro-minerals		Ascorbic acid (C)	0.045
Calcium	0.83	Thiamine (B1)	0.00608
Magnesium	1.38	α -tocopherol (E)	0.0021
Potassium	13.59	Folate	0.00394
Phosphorus	4.06	Niacin (B3)	0.0211
Sodium	0.12	Phylloquinone (k)	0.056 μ g/g
Micro-minerals		Pyridoxine (B6)	0.00397
Zinc	0.0279	Retinol	---
Iron	0.0669	Riboflavin (B2)	0.00212
Total dietary fibre	0.1520	Caloric value	3.37 kcal/g

3.4 Phytoconstituents: For various pharmacological activities, the main phenolic compounds in common beans can be summarized as phenolic acids, flavonoids, proanthocyanidins, and coumarins. In brief it contains Quercetin 3-O-glucoside, kaempferol 3-O-glucoside, myricetin, *p*-coumaric acid derivatives, ferulic acid derivatives, ferulic acid, caffeic acid, vanillin aldehyde [16], Catechin, epicatechin, epigallocatechin, quercetin, naringenin, chlorogenic acid, cichoric acid, coumaric acid, vanillic acid [12].

3.5 Traditional Medicinal use: The seeds of *Phaseolus vulgaris* are recorded as diuretic chiefly in kidney and heart disease. They are also effective in lenient cases of diarrhoea [17].

3.6 Pharmacological activities: The extract of *Phaseolus vulgaris* seeds are used as antioxidant and anti-inflammatory [10], anti-diabetic [11], anti-Parkinson [13], anti-proliferative [12], hepato-protective [18], trypsin, α -amylase [19, 20], analgesic, anti-fertility, litholytic [21] and antidepressant [22].

4. MORPHOLOGY

4.1 Physical Qualitative Characteristics: It includes following characteristics of seed viz, colour; shape; size; odour; taste and seed coat pattern.

The physical qualitative characteristics of seeds were evaluated as follows:

- 1) Seed colour – Dark brownish red
- 2) Seed shape – Kidney to oval shaped
- 3) Seed size – Medium
- 4) Seed coat Pattern – single colour on entire seed
- 5) Taste – Bland
- 6) Odour - None

4.2 Physical Quantitative Characteristics: Quantitative seed descriptors includes the physical evaluation (Table 4) of the following seven characteristics: average of 1 and 100 seed weight; seed length (L); seed thickness (T); seed width (W); diameter; volume and surface area. 1 and 100 seed weight were measured in six repetitions using a digital weighing balance. 10-randomly selected fully developed undamaged seeds were measured in six repetitions using a Vernier calliper (least count of 0.1mm). Length, Thickness and Width were measured from the highest, lowest aligned to hilum, and from hilum to the opposite side respectively. Various diameter means viz. Arithmetic (AMD), Geometric (GMD) and sphericity (ϕ) of kidney bean was calculated using equations given by.²³ Also, parameters like volume (V) and surface area (S) which depends on axial dimension (length) was calculated for single bean [24].

Table 4: Physical Quantitative Characteristics of *Phaseolus vulgaris*

Sr. No.	Parameters	Mean \pm S.E.M
---------	------------	------------------

1.	Seed length (L in cm)	2.090 ± 0.023
2.	Seed thickness (T in cm)	0.887 ± 0.010
3.	Seed width (W in cm)	1.150 ± 0.011
4.	Arithmetic mean diameter (AMD in cm)	1.376 ± 0.012
5.	Geometric mean diameter (GMD in cm)	1.287 ± 0.011
6.	Volume (in cm ³)	1.310 ± 0.035
7.	Surface area (in cm ²)	6.103 ± 0.110
8.	Sphericity (φ)	0.616 ± 0.004
9.	Weight variation within seed (one seed/g)	0.34 ± 0.03 – 0.72 ± 0.03
10.	Weight of 100 seeds (in g)	47.06 ± 0.373

4.3 Physico-chemical Parameters: Various physico-chemical parameters were estimated (Table 5) in triplicates, viz. moisture content, extractive values, ash values and swelling index. It gives an idea about the quality and purity of crude drugs.

Table 5: Physico-chemical Parameters of *Phaseolus vulgaris*

Sr. No.	Parameters	Determined values (% w/w)
A.	Moisture Content	1.03 ± 0.23
B.	Extractive values	
1.	Alcohol soluble extractive value	10.46 ± 0.65
2.	Water soluble extractive value	19.75 ± 0.41
C.	Ash Values	
1.	Total ash	1.87 ± 0.02
2.	Acid insoluble ash	0.33 ± 0.02
3.	Water soluble ash	0.82 ± 0.03
D.	Swelling index (in cm)	1.01 ± 0.06

5. MICROSCOPY

The microscopy was done using optical microscope (Olympus vanoz-s-AH-2, Japan) with various optical magnification and images were captured employing a digital camera. The cross section in Fig. 1 of soaked whole red kidney bean, visualizes the presence of three cell layers: the **cotyledon (A)**, **endosperm (B)** and **testa /seed coat (C)** (helps from mechanical injury, predators & drying out).




Figure 1: Optical light microscopy of red kidney bean (cross-section)

The transverse section (Fig 2) also showed the presence of **proteinaceous aleurone cells (blue arrow)**, **macro-sclereids (black arrow)**, which are important for the absorption of water by the seed are observed. The presence of **starch granules (yellow arrow)** with irregular oval shape in the cotyledon indicates the energy reserves of seeds.




Figure 2: Transverse-section of red kidney bean

6. EXTRACTION & PHYTOCHEMICAL SCREENING

6.1 Soaking & Extraction procedure: The drug was collected and shade-dried at room temperature of about $25\pm 2^{\circ}\text{C}$. 250g dry mature seeds of *Phaseolus vulgaris* were soaked overnight, for 16 h, in distilled water, on the proportion of 100 g per 300 mL of water (Figure 3). After soaking, seed coats were manually separated from cotyledons. Seed coats were further dried at room temperature, for an average period of 24 hours. Dried coats (7.97gm) were extracted without previous milling, with the ethanol: water (60:40, v/v) solution, followed by sonication. At the end using rotary flash evaporator under vacuum, the extract was concentrated to a semi-solid mass with the recovery of solvent. The traces of the solvents were separated by using lyophilizer. The resultant extract was stored at 4°C for future use.

The percentage yield was determined as follows:

$$\text{Percentage yield} = \frac{\text{Final weight of dried extract}}{\text{Initial weight of powder}} \times 100$$

$$\text{Percentage yield} = 23.30\text{gm} / 39\text{gm} \times 100 = \mathbf{58.97\%}$$




Figure 3: Soaking and extraction of red kidney bean

6.2 Preliminary phytochemical screening: Screening was performed as per standard protocol and results are depicted in Table 6 [25].

Table 6: Phytochemical screening of *Phaseolus vulgaris*

BIOACTIVE COMPONENTS	RESULT
Alkaloids	Present
Carbohydrates	Present
Flavonoids	Present
Tannins	Present
Saponin	Present
Antraquinone	Absent
Phenol	Present
Steroids	Absent

7. HPTLC OF BIOACTIVE COMPONENTS:

The ethanolic extract and flavonoid fraction were analyzed for the presence of flavonoids by comparing with the R_f value and spectral comparison with co-chromatographic standard compounds, Quercetin. HPTLC study was performed to standardize the extract of *Phaseolus vulgaris* seed coats for the presence of flavonoids in Figure 4 [26].

HPTLC fingerprinting was performed by using winCATS software. Sample application was executed by CAMAG Linomat 5 and inert gas spray and methanol as solvent type was used. After application chromatogram developed Twin Trough Chamber 20x10cm with Tol :EA:FA (10:3:1) at 60° C for 5 minutes. CAMAG TLC Scanner 3 was used to detect spots (Figure 5). The result from HPTLC fingerprinting scanned at 254nm and 366nm for extract by using CAMAG Reprostar 3 illumination instrument.




Figure 4: Spectral comparison of *Phaseolus vulgaris* seed coats with co-chromatographic standard, Quercetin.




Figure 5: Detection of active phytoconstituents in *Phaseolus vulgaris* seed coat by HPTLC

8. *In-vitro* ESTIMATION OF BIOACTIVE COMPONENTS

8.1 Determination of total flavonoids (TF): The total flavonoid content was confirmed according to the procedure given [27]. 1 ml extract of 1000 µg/ml concentration 4 ml of purified water was mixed and then 0.3 ml NaNO₂ & 0.3 ml AlCl₃ was added to solution after that mixture was incubated for 5 minutes at room temperature. Sodium Hydroxide (2ml) and purified water (2.4ml) was added to the incubated solution and the absorbance was measured at 510 nm with the help of spectrophotometer. Standard curve was used to determine Total Flavonoid content. Quercetin was used as standard and TF content was indicated as Quercetin equivalents (QE) in mg/g of dry sample. The results were shown in Table 7.

8.2 Determination of total phenols: Total phenolic content was evaluated by using Folin-Ciocalteu (FC) reagent. The evaluation was carried out spectrophotometrically as stated by [28] with minimum moderation. In a test tube, 0.1 ml of extract (1mg/ml) was taken and then 1.9 ml distilled water and 1.0 ml of Folin– Ciocalteu's reagent was added in a test tube, after that 1.0 ml of 100 g/L Na₂CO₃ was added to the solution. The mixture was incubated at room temperature for 2 hours and the absorbance of the solution was measured at 765 nm using spectrophotometer. The standard

curve of gallic acid was used to estimation of Total Phenolic Content. The total phenolic compounds of the plant extracts were indicated as gallic acid equivalents (GAE) which showed the phenolic content equal to the gallic acid (mg/g) of dry material. The results were shown in Table 7.

Table 7: *In-vitro* antioxidant potential of *Phaseolus vulgaris* seed coat

<i>Phaseolus</i>	Total Flavonoids Content	Total Phenolic Content
<i>vulgaris</i>	(TFC in mg /g QE)	(TPC in mg/g GAE)
seed coat	13.62 ± 0.49	32.03 ± 1.50

8.3 DPPH Radical scavenging assay: According to [29] the free radical scavenging activity was evaluated with the help of an improved DPPH assay. 2.7 mL (0.2 mM) DPPH solution was added to 0.3 mL of the extract of different concentrations. Then, the mixture was shaken efficiently and incubated at room temperature for 1 h in dark before the absorbance was taken at 517 nm.

$$\text{Percentage inhibition} = [(A_s - A_i)/A_s] \times 100$$

Where, A_s is the absorbance of pure DPPH and A_i is the absorbance of DPPH in the presence of different extracts. Vitamin C was used as reference. *In-vitro* antioxidant potential and % inhibition was depicted in table 8 and Figure 6.

Table 8: *In-vitro* antioxidant potential of *Phaseolus vulgaris* seed coat

IC_{50} value	Concentration of Vitamin C ($\mu\text{g/ml}$)	Concentration of seed coat ($\mu\text{g/ml}$)
	354.93 ± 7.37	429 ± 4.07




Figure 6: DPPH assay of *Phaseolus vulgaris* seed coat

9. RESULT AND DISCUSSION

Medicinal plants are very useful and they have been used for more than hundred years by mankind in the prevention and treatment of numerous diseases [30]. From the above context, this study was designed to describe pharmacognostic profile and evaluated antioxidant activity of extract from *Phaseolus vulgaris*. To signify quality control profile, the initial step is macroscopy and microscopy study of any herbal or medicinal plant when used in pharmaceutical industry.² In this study we are discussed taxonomical classification, nutritional value in which plant seed contain higher amount of protein, lipids and carbohydrates and other essential components such as macro-minerals and micro-minerals, vitamins and dietary fibres [31].

This manuscript explains the morphology of red kidney bean as physical qualitative characterization, physical quantitative characterization and physico-chemical parameters. These physical characterizations of seed are significant for evaluating the product quality. The result for physical properties *i.e.* AMD, GMD, volume and surface area depends on axial dimension (length) of seed, whereas sphericity is depended on the lowest volume of the seed. Also, it helps various personals like plant breeders, machine manufacturers, food scientists, etc. Harvesting, grasping, shipment, detaching, aeration, examining, storing, filling and the other measurement prescribed machines and

equipment and it is helpful to design relevant machine and equipment. Currently, there is no exclusive standard method is registered in prompting the physical dimensions of farming outcomes [32].

Microscopical examination of seed displayed three cell layers and showed the presence of proteinaceous cells, macro-sclerides and starch granule [33]. The highest percentage yield of hydroalcoholic extract of seed coat of *Phaseolus vulgaris* was obtained 58%. Phytochemical evaluation showed the existence of many bioactive compounds like alkaloids, glycosides, carbohydrates, tannins, saponin, steroids and several phenolic compounds such as flavonoids proteins. This study suggested hydroalcoholic extract of *Phaseolus vulgaris* have antioxidant potential owing to the presence of higher amount of phenol, flavonoids, saponins [34, 35]. In HPTLC, *Phaseolus vulgaris* exerted beneficial effects as compared with Quercetin as standard [36] and its Rf value was found 0.48. In antioxidant profile, higher value of total flavonoid and total phenol contents showed the presence of polyphenolic constituents and recommended its antioxidant action [37]. Scavenging of DPPH is one of the imperative parameters to assess the antioxidant effect of crude extracts. In this study extract exhibited higher percentage of DPPH scavenging activity and the study suggested plant extract contain flavonoids and related polyphenols [38].

10. CONCLUSION

Phaseolus vulgaris serve as decisive source of protein, minerals, vitamins, dietary fibres. This study described morphology, microscopy, HPTLC and in-vitro bioactive compound estimation. On the basis of result it is suggested that *Phaseolus vulgaris* have antioxidant property. Previously, it is proposed that *Phaseolus vulgaris* contains flavonoids such as Quercetin, kaempferol, naringin, naringenin, ferulic acid, myrcetin and it is used as anti-diabetic, anti-inflammatory, anti-depressant, anti-parkinsonism, analgesic, anti-proliferative, cardio-protective, hepato-protective. As we know epilepsy may occur due to oxidative stress and these flavonoids are useful in the treatment of epilepsy so, *Phaseolus vulgaris* could be beneficial in the treatment of epilepsy. In future, the present protocol may form the basis for the selection of plant species for further investigation in potent bioactive compounds for in-vivo activities.

The study highlights the efficacy of " Herbal " which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

REFERENCES

1. Shorvan SD. The etiologic classification of epilepsy, *Epilepsia*, 2011; 52(6): 1052-1057.
2. WHO. 2010. Epilepsy: key facts. [online] Available at: www.who.int/mediacentre/factsheets/fs999/en Accessed 7 June, 2010.
3. Lowenstein DH. 2008. Seizures and epilepsy. In: Fauci AS, Kasper DL, Longo DL, ed. *Harrison's Principles of Internal Medicine*. New York: McGraw-Hill 2498-2512.
4. McNamara JO. 1999. Drugs effective in the therapy of the epilepsies. In: Hardman JG, Limbird LE, Molinoff PB, Ruddon RW, ed. *Goodman & Gilman's The Pharmacological Basis of Therapeutics*. New York: McGraw-Hill 461-486.
5. Xiao F, Yan B, Chen L, Zhou D. Review of the use of botanicals for epilepsy in complementary medical system- traditional Chinese medicine. *Epilepsy behave*, 2015; 52: 281-89.
6. Copmans D, Steurs G. Methylated flavonoids as anti-seizure agents: naringenin 4, 7-dimethyl ether attenuates epileptic seizures in zebrafish and mouse models, *Neurochem Int*, 2018; 112: 124-133.
7. Sefil F, Ibrahim Kahraman, Neslihan Pinar. Ameliorating effect of Quercetin on acute PTZ induced seizures in Rats. *International Journal of Clinical and experimental medicine*, 2014; 7(9): 2471-2477.
8. Kola PK, Akula A, NissankaraRao LS, Danduga RSR. Protective effect of naringin on pentylenetetrazole (PTZ) induced kindling, *Epilepsy Behav*, 2017; 75: 114-126.

9. Nassiri M, Reza Martazavi, Fatemeh Samiee-Rad, Esmail Abbasi. The effect of rutin on the development of PTZ kindling and memory retrieval in rats. *Epilepsy behave.* 2010.
10. Akash, Sumitra S, Shailendra KS. Antioxidant and anti-inflammatory potential of kidney bean (*Phaseolus vulgaris* L). *International research journal of pharmacy,* 2018.
11. Luka CD, Olatunda A, tijjani H and Olisa-Enewe IA. Effect of aqueous extract of *Phaseolus vulgaris* (red kidney bean) on alloxan induced diabetic wistar rats. *International Journal of Science Invention Today,* 2013; 2(9): 292-301.
12. Ombra MN, d'Acierno A, Nazzaro F, Riccardi R, Spigno P, Zaccardelli M, Fratianni F. Phenolic composition and antioxidant and antiproliferative activities of the extracts of twelve common bean (*Phaseolus vulgaris* L.) endemic ecotypes of Southern Italy before and after cooking. *Oxidative Medicine and Cellular Longevity,* 2016; 13(9): 82-98.
13. Rashmi A, Bhakti S, Yash N, Tejas G. Evaluation and comparison of Anti Parkinson activity of methanolic extract of *Phaseolus vulgaris* with L-DOPA. *International journal of Pharmacology Research,* 2015; 103-108.
14. Porcher MH. *Phaseolus vulgaris* L. In: Multilingual multiscrypt plant name database. <http://www.plantnames.unimelb.edu.au/Sorting/Phaseolus.html> 2013 [cited 2016 14th Oct].
15. USDA. 2005. National Nutrient Database for Standard Reference Release 27: Basic Report 16032, Beans, Kidney, red, mature seeds, raw. [cited 2016 Sept 23, 03:45EDT]; Available at: <http://ndb.nal.usda.gov/ndb/nutrients/index> Accessed 23 September 2016.
16. Lin LZ, Harnly JM, Pastor-Corrales MS, Luthria DL. The polyphenolic profiles of common bean (*Phaseolus vulgaris* L.). *Food Chemistry,* 2008; 107(1): 399–410.
17. CSIR 1979. Anonymous. *The Wealth of India: Raw Materials.* Ph-Re. New Delhi, India: Publications and Information Directorate.

18. Oomah BD, Corbé A and Balasubramanian P. Antioxidant and anti-inflammatory activities of bean (*Phaseolus vulgaris* L.) hulls. *Journal of Agricultural and Food Chemistry*, 2010; 58(14): 8225-8230.
19. Reynoso C, Ramos G and Loarca P. 2006. Bioactive components in common beans (*Phaseolus vulgaris* L.): Advances in Agricultural and Food Biotechnology. In: GuevaraGonzález RG, Torres-Pacheco I, ed. Research Signpost. Trivandrum, Kerala, India, 217-236.
20. Marshall JJ and Lauda CM. Purification and properties of phaseolamin, an inhibitor of alpha-amylase from the kidney bean *Phaseolus vulgaris*. *Journal of Biological Chemistry*, 1975; 250(20): 8030-8037.
21. Rana JH, Patel UR, Patel SJ, Lambole V, Shah, DP. Pharmacological activities of *Phaseolus vulgaris*: A review. *Pharma Science Monitor*, 2016; 7(2): 107-115.
22. Madhu S, Ramica S. Antidepressant activity of aqueous extract of *Phaseolus vulgaris* in rodent models of depression. *International Journal of Nutrition, Pharmacology, Neurological Diseases*, 2014; 4:118-124.
23. Mohsenin NN. 1986. Physical properties of plant animal material II. NEWYORK USA: Gordon and Breach Sci. Pub.
24. Jain RK, Bal D. Properties of pearl millet. *J Agric Engg Res*, 66 (2): 85-91.
25. Kokate CK, 1999. Practical pharmacognosy. 4th edition, New Delhi, India: Vallabh Prakashan Publication.
26. Harborne JB. 1998. Phytochemical methods, a guide to modern techniques of plant analysis. New Delhi, India: Springer Pvt.Ltd.
27. Kim D, Jeong S, Lee C. Antioxidant capacity of phenolic phytochemicals from various cultivars of plums. *Food Chemistry*, 2003; 81(6): 321-326.
28. Slinkard K, Singleton VL. Total phenol analysis: automation and comparison with manual methods. *Am J Enol Vitic*, 1977; 28: 49-55.

29. Lillian B, Telma C, Paula B, Leticia ME, Isabel F. Wild and commercial mushrooms as source of nutrients and nutraceuticals. *Food and Chemical Toxicology*, 2008; 46: 2742-2747.
30. Baldivia DDS, Leite DF, deCastro DTH, Campos JF, DosSantos UP, Paredes-Gamero EJ, Carollo CA, Silva DB, dePicoli SK, DosSantos EL. Evaluation of in vitro antioxidant and anticancer properties of aqueous extract from stem bark of *Stryphnodendron adstringens*. *Int J Mol Sci* , 2018; 19: 2432.
31. Zainish MS, Salman A, Muhammad M H. *Phaseolus vulgaris* L.: Botany, medicinal uses, phytochemistry and pharmacology. *World Journal of Pharmaceutical Research*, 2016; 5: 1611-1615.
32. Waziri AN, Mittal JP. Design –related physical properties of selected agricultural products. *Food and Agriculture Organization of the United Nations*, 2013; 14: 59-62.
33. Joseph SG, Barry GS. Microstructure of black, green and red gram. *Food structure*, 1993; 12(2): 155-162.
34. Peerzada AM, Ali HH, Naeem M, Latif M, Bukhari AH, Tanveer A. *Cyperus rotundus* L.: Traditional uses, phytochemistry, and pharmacological activities. *J Ethnopharmacol*, 2015; 174: 540-560.
35. Akhtar S, Ismail T, Fraternali D, Sestili P. Pomegranate peel and peel extracts: Chemistry and food features. *Food Chem*, 2015; 174: 417-425.
36. Mishra J, Singh RD, Jadon VS, Gusain MP, Aradhana. HPTLC profile of Quercetin content of common bean. *American Journal of Food Technology*, 2012; 7(2): 96-100.
37. Khorasani EA, Mat TR, Mohajer S, Banisalam B. Antioxidant activity of total phenolic and flavonoid content of various solvent extracts from in vivo and in vitro grown *Trifolium pratense* L. (Red Clover). *BioMed Res Int*, 2015: 643285.
38. Nambooppha B, Photichai K, Wongsawan K, Chaummitri P. quercetin manipulates the expression of genes involved in the reactive oxygen species (ROS) process in chicken heterophils. *J Vet Med Sci*, 2018; 80: 1204-1211.

UNDER PEER REVIEW