

Original Research Article

ANTIBACTERIAL AND ANTIFUNGAL ACTIVITY OF SELECTED INDIAN TRADITIONAL MEDICINAL PLANTS

ABSTRACT

Antibacterial and antifungal activities are the universal supporting activities for other therapeutic activities like anti-inflammatory, wound healing etc. The traditional medicinal plants *Pongamia pinnata* and *Macaranga peltata* selected. The selected plants different extracts (Ethanol, Chloroform, Petroleum ether, Ethyl acetate and Methanol) were prepared and preliminary phytochemical screening was performed. The antibacterial and antifungal activity was performed by Agar well method by using Ciprofloxacin and Erythromycin standards. In this study we come to conclusion that the selected medicinal plants alcohol extract is having significant antibacterial and antifungal activity.

Key words: *Pongamia*, *Macaranga*, Antibacterial, Antifungal etc.

INTRODUCTION

Medicinal plants are used for curing innumerable diseases. In terms of medicinal uses mixture of constituents found in extracts of plants are more effective than isolated compounds. Many herbs in nature possess tissue regenerating property as they possess pharmacologically active compound in minute quantity along with energy boosting molecules such as carbohydrates, lipids and proteins.

The plants possess various therapeutic activities which should be brought to the notice of the scientific field for the systematic evaluation. Hence an attempt was made to select the plants possessing antibacterial and antifungal activity.

In the present study *Macaranga peltata* leaves and *Pongamia pinnata* seeds selected since scientifically antibacterial extracts and principles are not explored and standardized for the proposed activity. Both plants are traditional medicinal plants possessing anti-inflammatory, antioxidant other relevant activities.

METHODOLOGY

Collection and Authentication of Plants:

The *Macaranga peltata* leaves were collected from Paneer, Deralakatte areas in Mangalore. The *Pongamia pinnata* seeds were purchased from Pioneer Agro industry, Coimbatore, Tamilnadu. The collected plant materials were subjected to authentication at Foundation for

Revitalization of Local Health Traditions (FRLHT), Bangalore and the voucher specimen (4717-18/28.0/.2021) was deposited

Extraction: The authenticated seeds and leaves were dried in hot air oven, powder and sieved to obtain coarse powder. The powders were subjected to maceration with Ethanol, Chloroform, Petroleum ether, Ethyl acetate and methanol respectively. Filtered and dried to obtain the extract and preserved in desiccator.

Preliminary phytochemical screening for Phytochemicals: The extracts were subjected to preliminary phytochemical screening to know the presence of Phytochemicals such as Alkaloids, Carbohydrates, Tannins, Proteins, Saponins, Steroids, Tri terpenoids, Flavonoids and fixed oils.

Test for Alkaloids:

- Dragendroff's Test: The extract sample was dissolved in chloroform and Dragendroff's reagent was added, Reddish brown colour precipitate obtained indicates presence of alkaloids(Positive result)
- Mayer's Test: The extract sample was dissolved in chloroform and Mayer's reagent was added, Cream colour precipitate.
- Hager's Test: The extract sample was dissolved in chloroform and Hager's reagent was added, yellow colour precipitate.
- Wagner's Test: The extract sample was dissolved in chloroform and Dragendroff's reagent was added, Reddish brown colour precipitate.

Carbohydrates:

- Fehling's Test: To 1 mL of Fehling's solution A added 1 mL of Fehling solution B , Added 2 mL of the plant extract solution, mixed well and boil for 45minutes. The red precipitate formed at the bottom of test tube.
- Molisch's Test: To 2 ml extract solution in a test tube, added 1 drop of Molisch's reagent (10% α -naphthol in ethanol) than added 1-2 mL of conc. H₂SO₄ down the side of the test tube, it formed a layer at the bottom of the tube.

Tannins:

- FeCl₂ Test: To the 2ml of extract sample added few drops of ferric chloride formed brackish green precipitate (Tannins Positive)

- Lead acetate Test: To the 2ml of extract sample added few drops of 10% lead acetate solution formed white precipitate

Protein's test:

- Biuret test: To the 2ml of extract sample added few drops of copper sulphate solution and sodium hydroxide solution formed violet color(Proteins Positive)
- Millon's Test: To the 2ml of extract sample added few drops of Millons reagent, Heat gently and sodium nitrite solution added leads to formation of red precipitate

Saponin Glycosides:

- Foam Test: Shaken the 2ml of extract solution allowed to stand for 15 min foam retained indicated saponins present
- Haemolysis Test: To the 2ml of extract solution added few drops of blood, haemolysis of RBC and liberation of haemoglobin shows tannins present

Steroids and Triterpenoids:

- Liebermann Burchard Test: To the 2mg of extract dissolved in acetic anhydride, boiled, cooled the 1ml of concentrated H_2SO_4 added along the sides of test tube, Pink colour indicates the presence of Triterpenoids(Green colour for steroids)
- Salkowski Test: To the 2mg of extract dissolved in 1ml of concentrated H_2SO_4 yellow color formation indicates presence of Triterpenoids(Red colour for steroids)

Flavonoid:

- Shinoda Test: To the 2ml of extract solution, added Magnesium turnings and concentrated HCL drop wise magenta colour indicates presence of Flavonoids

Fixed oil and Fat:

- **Solubility Test**
 - Non Polar solvent-** Fixed oils are freely soluble in chloroform and diethyl ether.
 - Polar Solvent -** Fixed oils are insoluble in water
- Translucent spot test: Put a spot of 2ml of extract solution on filter paper, residue remained after drying indicates fixed oils present.
- The extracts were subjected to Antibacterial (Staphylococcus aureus, Escherichia coli, Streptococcus Pneumonia and P. aeruginosa) and antifungal(Candida albicans) activity by

Agar well diffusion method. The bacteria's were grown and preserved in the nutrient broth at 4°C for future use.

Antibacterial and Antifungal activity: Agar well diffusion method

The microbial cultures were subcultured one day before performing the antimicrobial activity. The nutrient agar media was prepared by dissolving the nutrient agar powder (24grams) Himedia in 1000ml distil water. The agar solution media was cooled to 50°C and inoculated with the Bacterial samples. The agar solutions containing bacteria were poured to the Petridishes and allowed for solidification. The wells were created in solidified agar media using borer.

The Standards (Ciprofloxacin 10µg/ml, Erythromycin 10µg/ml), Blank (DMSO) and test samples of plants extracts (Macaranga peltata 100mg/ml and Pongamia pinnata100mg/ml) were prepared. 0.1ml of Sample solutions, Blank and Standards were dropped to agar wells in petriplates with 100mm diameter using 1ml graduated pipette and Incubated in BOD incubator(SIPLAB, Serve well Instruments Pvt Ltd) for 24 hours at 37°C. The zone of inhibition was measured and compared with blank and standards to know the bactericidal activity of selected medicinal plant extracts. The same procedure was used for antifungal activity and standard Fluconazole was used for the present study.

RESULTS AND DISCUSSION

The Pongamia pinnata seed extracts and Macaranga peltata leaves extracts were described as follows:

PONGAMIA PINNATA

❖ Description of Ethanol extract:

- Nature: Semisolid paste
- Colour: Yellowish brown
- Odor: Oily odor(Fixed oil smell)
- After complete drying oily nature retained
- % yield: 10.8

❖ Description of Chloroform extract:

- Nature: Oily extract/Oil
- Colour: Slightly dark yellow colour
- Odor: Acrid oily odour
- % yield: 21.72

❖ Description of Petroleum ether extract:

- Nature: Viscous Oily extract/Fixed Oil
- Colour: Light yellow colour

- Odor: Oily odor
- % yield: 21.64
- ❖ **Description of Ethyl acetate extract:**
- Nature: Jelly liquid/Viscous liquid
- Colour: Yellow cream red colour
- Odor: Characteristic odor
- % yield: 21.12%
- ❖ **Description of Methanol extract:**
- ❖ Nature: Jelly Solid
- ❖ Colour: Reddish brown colour
- ❖ Odor: Bitter Characteristic odor(Astringent odor)
- ❖ % yield: 10.12

1. MACARANGA PELTATA

- ❖ **Description of Ethanol extract:**
- Nature: Amorphous solid
- Colour: Bark blackish green colour
- Odor: Characteristic
- % yield: 24.16
- ❖ **Description of Chloroform extract:**
- Nature: Amorphous solid
- Colour: light greenish yellow color
- Odor: characteristic odor
- % yield: 10.35
- ❖ **Description of Petroleum ether extract:**
- Nature: Amorphous solid
- Colour: yellowish brown colour
- Odor: Characteristic
- % yield: 10.15
- ❖ **Description of Ethyl acetate extract:**
- Nature: Amorphous solid
- Colour: Light yellow colour
- Odor: Characteristic
- % yield: 14.64
- ❖ **Description of Methanol extract:**
- ❖ Nature: Amorphous Solid
- ❖ Colour: Reddish brown colour
- ❖ Odor: Characteristic Astringent odor
- ❖ % yield: 16.36

Table 1: PERCENTAGE YIELD OF EXTRACT

SL NO	SOLVENT USED FOR EXTRACTION	PERCENTAGE YIELD OF EXTRACT (GRAM EXTRACT/ 100GRAM POWDER)	
		PONGAMIA PINNATA SEED	MACARANGA PELTATA LEAF

01	Ethanol	10.8	24.16
02	Chloroform	21.72	10.35
03	Petroleum Ether	21.64	10.15
04	Ethyl acetate	21.12	14.64
05	Methanol	10.12	16.36

UNDER PEER REVIEW

Table 2: PRELIMINARY PHYTOCHEMICAL SCREENING OF EXTRACTS:

SL NO	CHEMICAL TEST	PONGAMIA PINNATA SEED					MACARANGA PELTATA LEAF				
		Ethanol extract	Chloroform extract	Petroleum ether extract	Ethyl acetate extract	Methanol extract	Ethanol extract	Chloroform extract	Petroleum ether extract	Ethyl acetate extract	Methanol extract
01	Alkaloids: <ul style="list-style-type: none"> • Dragendroff's Test • Mayer's Test • Hager's Test • Wagner's Test 	-	+	+	+	-	-	+	+	+	-
02	Carbohydrates: <ul style="list-style-type: none"> • Fehling's Test • Molisch's Test 	+	-	-	-	+	+	+	+	+	
03	Tannins: <ul style="list-style-type: none"> • Fecl₂ Test • Lead acetate Test 	+	-	-	-	+	+	+	-	-	+
04	Protein's test <ul style="list-style-type: none"> • Biuret test • Millon's Test 	+	+	-	-	+	+	+	-	-	+
05	Saponin Glycosides <ul style="list-style-type: none"> • Foam Test 	-	+	+	+	-	-	-	-	-	-

	• Haemolysis Test	-	+	+	+	-	-	-	-	-	-
06	Steroids and Triterpenoids										
	• Liebermann Burchard Test	-	+	+	+	-	-	+	+	+	-
	• Salkowski Test	-	+	+	+	-	-	+	+	+	-
07	Flavonoid Test										
	• Shinoda Test	+	-	-	-	+	+	+	-	-	+
	• Conc HCL	+	-	-	-	+	+	+	-	-	+
	• Alkali Test	+	-	-	-	+	+	+	-	-	+
08	Fixed oil and Fat										
	• Solubility Test										
	iii. Non Polar solvent										
	a. Chloroform	+	+	+	+	+	-	+	-	-	-
	b. Diethyl ether	+	+	+	+	+	-	+	-	-	-
	iv. Polar Solvent										
	a. Water	--	--	--	--	--	+	-	-	-	+
	• Translucent spot test	+	+	+	+	+	-	+	-	-	-

(+) = Result positive;

(--)= Result Negative

Table 3 : ANTIBACTERIAL ACTIVITY OF PONGAMIA PINNATA SEED EXTRACTS AND MECARANGA PELTATA LEAF EXTRACTS

SL NO	ZONE OF INHIBITION OF <i>PONGAMIA PINNATA</i> EXTRACTS					ZONE OF INHIBITION OF <i>MECARANGA PELTATA</i> LEAF EXTRACTS			
	Samples/Standards	S aureus	E coli	S pneumonia	P acronerous	S aureus	E coli	S pneumonia	P acronerous
01	Negative Control: (DMSO)	10 ± 0.5	12.5 ± 0.5	11.5 ± 0.5	10 ± 0.5	10 ± 0.5	10 ± 0.5	10 ± 0.5	12.0 ± 0.5
02	Positive Control 01: Ciprofloxacin 0.1 ml(100µg) of Extract(1mg/1ml)	30 ± 0.2	39 ± 0.3	25.8 ± 0.4	31 ± 0.2	30 ± 0.2	30 ± 0.4	20 ± 0.7	25.0 ± 0.2
03	Positive Control 02: Erythromycin 0.1 ml(100µg) of Extract(1mg/1ml)	34 ± 0.7	35 ± 0.5	30 ± 1	31 ± 0.5	32.5 ± 0.3	35 ± 0.5	24 ± 0.2	33.5 ± 0.3
04	Alcohol Extract 0.1 ml(10mg) of Extract(100mg/ml)	23.5 ± 0.5	19 ± 0.5	18 ± 0.5	19 ± 0.2	18 ± 0.3	19 ± 1	19.5 ± 0.4	17.5 ± 0.5
05	Chloroform Extract 0.1 ml(10mg) of Extract(100mg/ml)	19 ± 0.1	18 ± 0.5	16 ± 0.4	19 ± 0.4	20 ± 0.2	14 ± 0.5	11.5 ± 0.5	17.0 ± 0.4
06	Petroleum Ether 0.1 ml(10mg) of Extract(100mg/ml)	12 ± 0.2	20 ± 0.3	11.5 ± 0.3	12.5 ± 0.5	17 ± 0.3	12.5 ± 0.5	11 ± 0.5	11.0 ± 0.5
07	Ethyl Acetate Extract 0.1 ml(10mg) of Extract(100mg/ml)	15.5 ± 0.5	15 ± 0.4	11.5 ± 0.5	13 ± 0.3	18 ± 0.4	12 ± 0.1	13 ± 0.2	12.0 ± 0.2
08	Methanol Extract 0.1 ml(10mg) of	14.5 ± 0.4	25.5 ± 0.5	10 ± 0.2	11.0 ± 0.1	19 ± 0.5	19 ± 0.3	14 ± 1	12.0 ± 0.2

Extract(100mg/ml)									
-------------------	--	--	--	--	--	--	--	--	--


Values are mean (in mm) \pm Standard Deviation, DMSO-Dimethyl sulphoxide

UNDER PEER REVIEW

Table 4 : ANTIFUNGAL ACTIVITY OF PONGAMIA PINNATA SEED EXTRACTS AND MECARANGA PELTATA LEAF EXTRACTS(CANDIDA ALBICANS)

SL NO	<i>PONGAMIA PINNATA</i> EXTRACTS	ZONE OF INHIBITION against <i>Candida albicans</i>	<i>MECARANGA PELTATA</i> EXTRACTS	ZONE OF INHIBITION against <i>Candida albicans</i>
01	Negative Control (DMSO)	10 ±00	Blank (DMSO)	10 ±00
02	Positive Control Fluconazole 0.1 ml(100µg) of Extract(1mg/1ml)	20 ± 00	Standard 01 Fluconazole (10µg/ml)	20 ± 00
03	Alcohol Extract 0.1 ml(10mg) of Extract(100mg/ml)	16 ± 0.4	Alcohol Extract (100mg/ml)	22.5 ± 0.5
04	Chloroform Extract 0.1 ml(10mg) of Extract(100mg/ml)	17.5 ± 0.2	Chloroform Extract (100mg/ml)	21.5 ± 0.1
05	Petroleum Ether 0.1 ml(10mg) of Extract(100mg/ml)	14 ± 0.5	Petroleum Ether Extract(100mg/ml)	12 ± 0.5
06	Ethyl Acetate Extract 0.1 ml(10mg) of Extract(100mg/ml)	17.0 ± 0.4	Ethyl Acetate Extract(100mg/ml)	17.0 ± 0.2
07	Methanol Extract 0.1 ml(10mg) of Extract(100mg/ml)	19 ± 0.5	Methanol Extract (100mg/ml)	26.5 ± 0.1


Values are mean (in mm) ± Standard Deviation, DMSO-Dimethyl sulphoxide



1. Negative Control	4. Chloroform extract	7. Methanol extract
2. Positive Control 01(Ciprofloxacin)	5. Petroleum ether extract	8. Methanol extract
3. Positive Control 01(Erythromycin)	6. Ethyl acetate extract	9. Negative Control

Fig 1 : ANTIBACTERIAL ACTIVITY OF PONGAMIA PINNATA EXTRACTS ON E COLI

Fig 2 : ANTIFUNGAL ACTIVITY OF PONGAMIA PINNATA SEED AND MACARANGA PELTATA LEAF EXTRACTS



1. Negative control	4. Pongamia pinnata chloroform extract	7. Pongamia pinnata methanol extract	10. Macaranga peltata petroleum ether extract
2. Standard	5. Pongamia pinnata petroleum ether extract	8. Macaranga peltata alcohol extract	11. Macaranga peltata ethyl acetate extract
3. Pongamia pinnata ethanol extract	6. Pongamia pinnata ethyl acetate extract	9. Macaranga peltata chloroform extract	12. Macaranga peltata methanol extract

Conclusions

❖ **Antibacterial activity:**

1. *Pongamia pinnata* alcoholic extract shown significant anti-bacterial activity against *S. aureus*, whereas other extracts(Chloroform, Petroleum ether, Ethyl acetate, methanol) of *Macaranga peltata* shown significant anti-bacterial activity against *S. aureus*
2. *P. pinnata* methanol extract shown good bactericidal activity against *E. coli*
3. The ethanolic extracts of both plants shown prominent bactericidal activity compared to all other solvent extracts
4. The descending order of antibacterial activity: Ethanolic extract> Chloroform extract> Methanol extract> Petroleum ether= Ethyl acetate
5. All the extracts shown very less antibacterial activity against *Streptococcus Pneumonia* (*S. Pneumonia* resistant to selected plant extracts)
6. Selected Second Standard erythromycin shown better bactericidal activity than Ciprofloxacin and many clear zone of inhibition were noticed in erythromycin treated wells
7. Against *S. aureus* *P. pinnata* alcoholic extract and *M. peltata* Chloroform, P. ether, Ethyl acetate and methanol extracts shown significant bactericidal activity
8. *S. Pneumonia* is found to be slightly resistant to selected standard antibiotics (Ciprofloxacin and Erythromycin)
9. All the extracts of both plants shown very less activity against *P. acro nerous*
10. The *S. aureus* and *E. coli* infections can be well treated with the selected medicinal plants extracts

❖ **Antifungal activity:**

1. The standard Fluconazole and sample extracts have shown antifungal activity compared to blank.
2. *Macaranga peltata* leaf methanol extract, Ethanolic extract and chloroform extract have shown better antifungal activity compared to selected dose of standard Fluconazole(10µg/ml)
3. All the extracts of *Pongamia pinnata* shown less antifungal activity than standard.

4.

5. **COMPETING INTERESTS DISCLAIMER:**

6.

7. Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge.

Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

8.
9.

10.

References:

1. Singh RK, Pandey BL. Anti-inflammatory activity of seed extracts of *Pongamia pinnata* in rat. Indian J. of Physio. Pharmacol. 1996 Oct 1; 40: 355-8.
2. Kumar P, Semalty A, Semalty M. Antibacterial and Antifungal activity of *Pongamia pinnata*. Indian drugs.2010; 47(3): 48- 50
3. Rani MS, Dayanand CD, Shetty J, Vegi PK, Kutty AM. Evaluation of antibacterial activity of *Pongamia pinnata* linn on pathogens of clinical isolates. Am. J. Phytomed. Clin. Thera.. 2013; 1(8):645-51.
4. Kage DN, Tabassum N, Malashetty VB, Deshpande R, Seetharam YN. Isolation and pharmacological studies of karanjachromene from the seeds of *Pongamia pinnata* (L. Pierre). Int. J Cur Res Rev. 2016; 8(17):35.
5. Dwivedi D, Dwivedi M, Malviya S, Singh V. Evaluation of wound healing, anti-microbial and antioxidant potential of *Pongamia pinnata* in wistar rats. J. Tr. Compl. Medi. 2017; 7(1):79-85.
6. Nehete M, Tatke P. Antioxidant, antimicrobial and wound healing potential of *macaranga peltata* bark extracts. Eur. J. Pharm. Med. Res. 2016; 3(10): 313-320
7. Badarudheen R, Lubna N, Dufaida KM, Mashhoor K. Antibacterial Activity of *Macaranga peltata*. Int. J. Adv. Res. Emerging Discipli. 2015; 3(2): 77-83

8. Verma M. NK, Thakar M, Subrahmanyam VM, Rao V, Dhanaraj SA. Investigation of antibacterial and antifungal potentials of *macaranga peltata*. Int. J. Curr. Res. Rev. 2013; 5(7): 17-24
9. Bijesh K, Sebastian D. Isolation and characterization of antibacterial compounds from *Macaranga peltata* against clinical isolates of *Staphylococcus aureus*. Int. J. Biol. Pharm. Res. 2013; 4: 1196-203.
10. Subrahmanyam VM, Rao V, Dhanaraj SA, Vasanth Raj P. Investigation of antibacterial and antifungal potentials of *Macaranga peltata*. Int J Cur Res Rev. 2013 Apr; 5(07).

UNDER PEER REVIEW