

Analysis of antimicrobial activity of aqua alcoholic extract of *Boerhaviadiffusa* against oral pathogens -An Invitro study

ABSTRACT

INTRODUCTION

Boerhaviadiffusa(*Mukkirattai*) of the *Nyctaginaceae* family is a widely used folk medicinal plant that grows as a common weed with its leaves, seeds and roots are useful parts with pharmacological activities with a cure for twenty-three ailments. Phytochemical constituents of *B.diffusa* showing antimicrobial activity includes phenols, flavonoids, tannins, saponins, alkaloids, glycosides.

AIM

The present study aimed to evaluate the qualitative analysis of the antimicrobial activity of aqua alcoholic extracts of *Boerhaviadiffusa* L. (Family: *Nyctaginaceae*) leaves.

MATERIALS AND METHODS

Boerhaviadiffusa (*Mukkirattai*) was freshly procured as a powdered form and the antimicrobial activity of different solvent extracts of *B. diffusa* were tested against the Gram-positive bacteria and fungal strains by observing the zone of inhibition. The Gram-positive bacteria used in the test were *Staphylococcus aureus*, *Enterococcus faecalis*, *Streptococcus mutans* and fungal strains *Candida albicans* were used. The obtained data were analysed statistically by Non parametric Spearman correlation analysis.

RESULTS

Antimicrobial activity was observed in the aqueous-alcoholic extracts against Gram-positive bacilli , cocci and fungal strains. The aqua alcoholic extract of *B. diffusa* showed positive correlation with p value less than 0.05 with antimicrobial effect against fungal strains(e.g. *C. albicans*, the zone of inhibition diameter of 26mm) and against Gram-positive bacteria(e.g. *S. aureus*, the zone of inhibition diameter of 20 mm) and then Gram-positive cocci (e.g. *S.Mutans*, the zone of inhibition diameter of 18 mm) when compared to *E.faecalis*.

CONCLUSION

The Aqua Alcoholic extract of *Boerhaviadiffusa* has a very strong Antimicrobial activity against *Candida Albicans* and *Staphylococcus Aureus* while it showed minimal antimicrobial activity against *Enterococcus faecalis* and *Streptococcus mutans* and showed excellent potential as an antimicrobial agent and also as a biofriendly, inexpensive.

Keywords*Boerhaviadiffusa*, Antimicrobial, Aqua alcoholic extract, Green synthesis, Zone of inhibition, *Nyctaginaceae*

INTRODUCTION

The *Boerhavia* genus distributed in the tropics of Asia, Africa, and Australia has around forty species. *Boerhaviadiffusa* of the *Nyctaginaceae* family is a widely used folk medicinal plant⁽¹⁾. *B.diffusa* grows as a common weed with its leaves, seeds and roots are the useful parts with pharmacological activities with a cure for twenty-three ailments includes cardioprotective effect, treatment of prostatic hyperplasia, anti-inflammatory action, anxiolytic activity, protective effect on gastrointestinal problems, anticancer activity, antimicrobial activity, protection against harmful radiations, hepatoprotective activity, anti-arthritic activity and antidiabetic activity.^(2,3) Other Indian names include varshabhu, Tambadivasu, Snathikari. It also helps in resolving abdominal pain, jaundice, diabetes, elephantiasis, ulcers, etc.,⁽⁴⁾ Phenols, flavonoids, tannins, saponins, alkaloids, glycosides are the basic aqua-alcoholic phytochemical constituents of *B.diffusa* showing antimicrobial activity and in previous studies showed the presence of flavonoids, alkaloids, steroids, triterpenoids, lipids, lignins, carbohydrates, proteins and glycoproteins in *B.diffusa* extracts⁽⁵⁻⁷⁾. Alcoholic and alkaline phytochemical flavonoids extract of *B.diffusa* is more compared to phenols and alkaloids⁽⁸⁾

The seeds of *Boerhaviadiffusa* have anti-bacterial, anti-fungal and anti-pathogenic activity against *E.faecalis*, *S.aureus*, *S.mutans*, *C.albicans*.^(9,10) Antibiotics result in multidrug resistance in treating infectious disease also associated with anaphylactic reactions.⁽¹¹⁾ This forced the physicians to create new antimicrobial substances from medicinal plants with fewer hypersensitivity reactions. *B. diffusa* is also used in Ayurvedic medicine in India and as an Unani medicine in Arab countries for the treatment of various factors like inflammation, jaundice, enlargement of spleen, congestive heart failure, diabetes, stress, dyspepsia, abdominal pain.⁽¹²⁻¹⁵⁾ It has also been reported that *Boerhaaviadiffusa* is useful in the treatment of corneal ulcers, nephritic syndrome, elephantiasis, night blindness.⁽¹⁶⁻¹⁸⁾

Several screening studies have been carried out in different parts of the world. Plant-based antimicrobials are the source of medicines with multiple therapeutic potentials.⁽¹⁹⁾ Both stem and leaf extract was tested against seven fungi species and six bacterial species, two types of yeast which showed antimicrobial activity in a dose-dependent manner (300-1800µg) and our team has extensive knowledge and research experience that has translate into high quality publications^{(20),(21-34),(35-39)} In this present investigation, we have analysed the antimicrobial activity of aqua alcoholic extract of *Boerhaaviadiffusa*.

MATERIALS AND METHODS

Preparation of plant extract

Boerhaviadiffusa (Mukkirattai) was freshly procured from Nature and Nurture health care pvt limited, New delhi as a powdered form which is the main advantage in our study. *Boerhaviadiffusa* is a commonly available leaf in southern India and is well known for its health benefits. It is the best green medicine for diabetes. Aqueous-alcoholic extract was prepared in the nanotechnology lab of Saveetha Dental College and Hospital. This original

study protocol was reviewed and approved by the research ethical committee of Saveetha Dental College and Hospitals, Chennai, Tamilnadu, India.

For preparing an aqueous-alcoholic extract, 50 ml of ethanol is measured using a measuring cylinder. 5g of powdered *B. diffusaw* was added to the 50 ml of ethanol and mixed well. Now the extract is transferred to the glass beaker and 50 ml of distilled water is added to the alcoholic extract. The beaker with aqueous-alcoholic extract of *Boerhaviadiffusa* is covered with aluminum foil paper and then kept in an orbital shaker for general mixing of the extract at 79.20 rpm. After 24 hours the extract was transferred to the measuring cylinder and the extract was boiled at 10% for 20 minutes and the extract was cooled at room temperature.

Antimicrobial activity

Antibacterial Activity

Antibacterial activity of respective nanoparticles against the strain *staphylococcus aureus*, *Enterococcus faecalis*, and *staphylococcus mutans*. MHA agar was utilized for this activity to determine the zone of inhibition. Muller Hinton agar was prepared and sterilized for 45 minutes at 120lbs. Media poured into the sterilized plates and let them stabilize for solidification. The wells were cut using the well cutter and the test organisms were swabbed. The plant extract at 25 μ L, 50 μ L, 100 μ L, 150 μ L concentrations were loaded and plates were incubated for 24 hours at 37 ° C. After the incubation time the zone of inhibition was measured. Manual labelling of the organisms should be given to avoid manual error.

Antifungal activity

Candida albicans are used as test pathogens by agar well diffusion assay. Sabouraud's dextrose Agar is used to prepare the medium. The prepared and sterilized medium was swabbed with test organisms and nanoparticles with different concentrations were added to the wells. The plates were incubated at 28° C for 48-72hours. After the incubation time, the zone of inhibition was measured and tabulated. The data obtained were tabulated and analysed by non parametric spearman correlation analysis using SPSS version 23.

RESULTS AND DISCUSSION

The anti-microbial activity of aqua-alcoholic extract of *Boerhaaviadiffusa* against *Streptococcus mutans*, *Staphylococcus aureus*, *Enterococcus faecalis*, and *Candida albicans* summarized in Figure 3. The result revealed that *Boerhaaviadiffusa* showed antimicrobial activity of different magnitudes. The zone of inhibition at different concentrations (25 μ L, 50 μ L, 100 μ L, 150 μ L) was done showing a reduction of the diameter of colonies around 26mm for *C.albicans*, 20mm for *S.aureus*, 18mm for *S.mutans* and 17mm for *S.faecalis*. Microorganisms are sensitive to different components of extracts of *Boerhaaviadiffusa*. Spearman correlation analysis showed positive correlation ($r=1$) of decrease in zone of inhibition (mm) with increase in concentration and significant p value of less than 0.023. The sensitivity of bacterial and fungal species is observed in the following decreasing order *C.albicans*>*S.aureus*>*S.mutans*>*E.faecalis*.




Figure 1 Schematic representation of Preparation of *Boerhaviadiffusa* extract. A-powdered extract of *Boerhaviadiffusa*. B-Aqua Alcoholic extract of *Boerhaviadiffusa*.




Figure 2 Anti-microbial activity of *B.diffusa* extract on A - *Streptococcus mutans*, B - *Staphylococcus aureus*, C - *Enterococcus faecalis*, and D - *C.albicans*




Figure 3: Bar graph depicts the concentration of antimicrobial activity (x axis) of Aqua -alcoholic extract of *Boerhaaviadiffusa* and the zone of inhibition (y axis). Positive correlation ($r=1$) with p value <0.05 is observed with increasing concentration of Aqua -alcoholic extract of *Boerhaaviadiffusa*. Blue colour denotes staphylococcus faecalis, green represents staphylococcus aureus, grey represents streptococcus mutans, purple represents candida albicans. From this it is inferred that albicans showed maximum zone of inhibition with the least seen in faecalis.




Figure 4 Line Graph depicts the zone of inhibition of *Candida Albicans* was higher with increase in concentration.

TABLE-1 - Depicts the zone of inhibition for *Streptococcus mutans*, *Staphylococcus aureus*, *Enterococcus faecalis*, *Candida albicans* at increasing concentrations.

Concentration	25 μL	50 μL	100μL	150 μL
<i>E.faecalis</i>	9mm	11mm	15mm	17mm
<i>S. aureus</i>	9mm	12mm	18mm	20mm
<i>S. mutans</i>	10mm	13mm	15mm	18mm
<i>C.albicans</i>	14mm	18mm	25mm	26mm

In the present study, the zone of inhibition for *Enterococcus faecalis* at 25μL was 9mm, at 50μL it was 11mm and for 100μL it was 15mm and for 150μL it is 17mm. The zone of inhibition for *staphylococcus aureus* at 25μL is 9mm and at 50μL it is 12mm and at 100μL it is 18mm and at 150μL it is 20mm. The zone of inhibition for *streptococcus mutans* at 25μL is 10mm and at 50μL it is 13mm and for 100μL it is 15mm and at 150μL it is 18mm. The zone of inhibition for *Candida albicans* at 25μL is 14mm and at 50μL it is 18mm and at 100μL it is 25mm and at 150μL it is 26mm.

Rotenoids are prototype compound named rotenone and isoflavonoids derivative a mitochondrial inhibitor causes ion inhibition electron transport chain in mitochondria at complex I, “toxophore” prenyl-derived ring of rotenoid structure and dimethoxy substitute to rotenone. Rotenoids isolated from *Boerhaaviadiffusa* are noncytotoxic due to the lack of isoprenoid residue.⁽⁴⁰⁾

Staphylococcus and *Streptococcus*, Gram-positive bacteria invades skin, tissues, and bloodstreams. Coagulases, proteins activate the hemostatic factor prothrombin of the host, Surface of bacteria display agglutinins, proteins and fibrin are the virulence factors of *S. aureus* infections leading to the destruction of immune cells, resulting in purulent exudate.⁽⁴¹⁾ *S. aureus* has superantigen (SSL5&SSL10), prevents rolling and adherence of neutrophil along endothelium. *S. aureus* diminishes opsonization by targeting complement activation systems.⁽⁴²⁾

Candida albicans adhesion(adhesins) and invasion into host cells with the secretion of hydrolases, yeast hyphae transition, thigmotropism, phenotypic switching, altered pH biofilm formation (Hwp1 and Als3) Secreted aspartic proteases (Saps) lead to endocytosis of *C. Albicans*. The utilization of lipases and amino acids results in hyphae formation. Hog1-, Mkc1-, Cek1-MAP kinase pathway responsible for maintaining the integrity of candida to the host surface.⁽⁴³⁾ *Candida albicans* adhere to the host surface with adhesin expression yeast-hyphae transition, growth by thigmotropism, Invasins mediate endocytosis of fungus into the host cell breaking down the barriers. Heat shock proteins, amino acids, lipases, ammonia excretion, and different trace compounds like zinc, carbon, manganese are responsible for hyphae formation.

Extract of *Boerhaaviadiffusa* induced systemic resistance active component BDP-30 a glycoprotein, pI greater than 9.0 with amino acid sequence KLYDIPPLR is responsible for antimicrobial activity by inhibiting bacterial transduction between the host and the recipient cells also inhibits candida albicans biofilm formation by preventing dimorphism and switching of candidal hyphae formation⁽⁴⁴⁾. The limitations of the study are constrained with four microorganisms at different concentrations. In future similar study in large scale productions for targeted drug delivery to treat and prevent a wide array of oral microbial infections.

CONCLUSION

The Aqua Alcoholic extract of *Boerhaviadiffusa* has a very strong Antimicrobial activity against *Candida Albicans* and *Staphylococcus Aureus* while it showed minimal antimicrobial activity against *Enterococcus faecalis* and *Streptococcus mutans* and showed excellent potential as an antimicrobial agent and also as a biofriendly, inexpensive.

• **COMPETING INTERESTS DISCLAIMER:**

-
- Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.
-

-
-
-

REFERENCE

1. Nugraha AS. Bioactive compounds from *Boerhavia erecta* L.: an African medicinal plant. 2009; Available from: <http://ro.uow.edu.au/cgi/viewcontent.cgi?article=4676&context=theses>
2. Patel SS, Raghuwanshi R, Masood M, Acharya A, Jain SK. Medicinal plants with acetylcholinesterase inhibitory activity. *Rev Neurosci*. 2018 Jul 26;29(5):491–529.
3. Preethikaa S, Brundha MP. Awareness of diabetes mellitus among general population. *Research Journal of Pharmacy and Technology*. 2018;11(5):1825–9.
4. Subramoniam A. *Anti-Diabetes Mellitus Plants: Active Principles, Mechanisms of Action and Sustainable Utilization*. CRC Press; 2016. 390 p.
5. Agarwal RR, Dutt SS. Chemical examination of punarnava or *Boerhaaviadiffusa* Linn. II. Isolation of an alkaloid punarnavine. *Chem Abst*. 1936;30:3585.
6. Basu NK, Lal SB. Investigations on Indian medicinal plants. *Q J Pharm Pharmacol*. 1947 Jan;20(1):38; passim.
7. Surange SR, Pendse GS. Pharmacognostic study of roots of *Boerhaaviadiffusa* Willd.(punarnava). *J Res Indian Med*. 1972;7(1).
8. Jayachitra J, Janani B, Bharathi V, Manikandan R. Phytochemical analysis and mineral composition of Methanolic extract of *Boerhavia diffusa* L [Internet]. Vol. 13, *Research Journal of Pharmacy and Technology*. 2020. p. 4856. Available from: <http://dx.doi.org/10.5958/0974-360x.2020.00854.9>
9. Prasad R, Jha AK, Prasad K. *Exploring the Realms of Nature for Nanosynthesis*. Springer; 2018. 414 p.
10. Brundha MP, Pathmashri VP, Sundari S. Quantitative Changes of Red Blood cells in Cancer Patients under Palliative Radiotherapy-A Retrospective Study. *Research Journal of Pharmacy and Technology*. 2019;12(2):687–92.
11. Lin DM, Koskella B, Lin HC. Phage therapy: An alternative to antibiotics in the age of multi-drug resistance. *World journal of gastrointestinal* [Internet]. 2017; Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/pmc5547374/>
12. Kīrtikara KR, Basu BD. *Indian medicinal plants* / [Internet]. 1918. Available from: <http://dx.doi.org/10.5962/bhl.title.137025>
13. Dagogo-Jack S. *Diabetes Mellitus in Developing Countries and Underserved Communities*. Springer; 2016. 294 p.
14. Chopra RN, Nayar SL, Chopra IC, Others. *Glossary of Indian medicinal plants*. Vol. 1. Council of Scientific & Industrial Research New Delhi; 1956.

15. Mishra JP, Others. Studies on the effect of indigenous drug Boerhaaviadiffusa Rom. on kidney regeneration. *Indian J Pharm.* 1980;12(59):1487–98.
16. Singh RH, Udupa KN. Studies on the Indian indigenous drug Punarnava (*Boerhaaviadiffusa L.*) Part I, Identification and pharmacological studies. *J Res Indian Med.* 1972;7:1–12.
17. Bharali R, Azad MRH, Tabassum J. Chemopreventive action of Boerhaaviadiffusa on DMBA-induced skin carcinogenesis in mice. *Indian J PhysiolPharmacol.* 2003 Oct;47(4):459–64.
18. Timothy CN, Samyuktha PS, Brundha MP. Dental pulp Stem Cells in Regenerative Medicine--A Literature Review. *Research Journal of Pharmacy and Technology.* 2019;12(8):4052–6.
19. van Vuuren S, Viljoen A. Plant-based antimicrobial studies--methods and approaches to study the interaction between natural products. *Planta Med.* 2011 Jul;77(11):1168–82.
20. Anita R, Paramasivam A, Priyadharsini JV, Chitra S. The m6A readers YTHDF1 and YTHDF3 aberrations associated with metastasis and predict poor prognosis in breast cancer patients. *Am J Cancer Res.* 2020 Aug 1;10(8):2546–54.
21. Jayaseelan VP, Paramasivam A. Emerging role of NET inhibitors in cardiovascular diseases. *Hypertens Res.* 2020 Dec;43(12):1459–61.
22. Sivakumar S, SmilineGirija AS, VijayashreePriyadharsini J. Evaluation of the inhibitory effect of caffeic acid and gallic acid on tetR and tetM efflux pumps mediating tetracycline resistance in *Streptococcus sp.*, using computational approach. *Journal of King Saud University - Science.* 2020 Jan 1;32(1):904–9.
23. SmilineGirija AS. Delineating the Immuno-Dominant Antigenic Vaccine Peptides Against gacS-Sensor Kinase in *Acinetobacter baumannii*: An in silico Investigational Approach. *Front Microbiol.* 2020 Sep 8;11:2078.
24. IswaryaJaisankar A, SmilineGirija AS, Gunasekaran S, VijayashreePriyadharsini J. Molecular characterisation of csgA gene among ESBL strains of *A. baumannii* and targeting with essential oil compounds from *Azadirachta indica*. *Journal of King Saud University - Science.* 2020 Dec 1;32(8):3380–7.
25. Girija ASS. Fox3+ CD25+ CD4+ T-regulatory cells may transform the nCoV's final destiny to CNS! *J Med Virol* [Internet]. 2020 Sep 3; Available from: <http://dx.doi.org/10.1002/jmv.26482>
26. Jayaseelan VP, Ramesh A, Arumugam P. Breast cancer and DDT: putative interactions, associated gene alterations, and molecular pathways. *Environ Sci Pollut Res Int.* 2021 Jun;28(21):27162–73.
27. Arumugam P, George R, Jayaseelan VP. Aberrations of m6A regulators are associated with tumorigenesis and metastasis in head and neck squamous cell carcinoma. *Arch Oral Biol.* 2021 Feb;122:105030.
28. Kumar SP, Girija ASS, Priyadharsini JV. Targeting NM23-H1-mediated inhibition of tumour metastasis in viral hepatitis with bioactive compounds from *Ganoderma lucidum*: A

computational study. pharmaceutical-sciences [Internet]. 2020;82(2). Available from: <https://www.ijpsonline.com/articles/targeting-nm23h1mediated-inhibition-of-tumour-metastasis-in-viral-hepatitis-with-bioactive-compounds-from-ganoderma-lucidum-a-comp-3883.html>

29. Girija SA, Priyadharsini JV, Paramasivam A. Prevalence of carbapenem-hydrolyzing OXA-type β -lactamases among *Acinetobacter baumannii* in patients with severe urinary tract infection. *Acta Microbiol Immunol Hung*. 2019 Dec 9;67(1):49–55.
30. Priyadharsini JV, Paramasivam A. RNA editors: key regulators of viral response in cancer patients. *Epigenomics*. 2021 Feb;13(3):165–7.
31. Mathivadani V, Smiline AS, Priyadharsini JV. Targeting Epstein-Barr virus nuclear antigen 1 (EBNA-1) with Murrayakoengii bio-compounds: An in-silico approach. *Acta Virol*. 2020;64(1):93–9.
32. Girija As S, Priyadharsini J V, A P. Prevalence of Acb and non-Acb complex in elderly population with urinary tract infection (UTI). *Acta Clin Belg*. 2021 Apr;76(2):106–12.
33. Anchana SR, Girija SAS, Gunasekaran S, Priyadharsini VJ. Detection of csgA gene in carbapenem-resistant *Acinetobacter baumannii* strains and targeting with *Ocimum sanctum* biocompounds. *Iran J Basic Med Sci*. 2021 May;24(5):690–8.
34. Girija ASS, Shoba G, Priyadharsini JV. Accessing the T-Cell and B-Cell Immuno-Dominant Peptides from *A.baumannii* Biofilm Associated Protein (bap) as Vaccine Candidates: A Computational Approach. *Int J Pept Res Ther*. 2021 Mar 1;27(1):37–45.
35. Arvind P TR, Jain RK. Skeletally anchored forsus fatigue resistant device for correction of Class II malocclusions-A systematic review and meta-analysis. *OrthodCraniofac Res*. 2021 Feb;24(1):52–61.
36. Venugopal A, Vaid N, Bowman SJ. Outstanding, yet redundant? After all, you may be another *Choluteca* Bridge! *Semin Orthod*. 2021 Mar 1;27(1):53–6.
37. Ramadurai N, Gurunathan D, Samuel AV, Subramanian E, Rodrigues SJL. Effectiveness of 2% Articaine as an anesthetic agent in children: randomized controlled trial. *Clin Oral Investig*. 2019 Sep;23(9):3543–50.
38. Varghese SS, Ramesh A, Veeraiyan DN. Blended Module-Based Teaching in Biostatistics and Research Methodology: A Retrospective Study with Postgraduate Dental Students. *J Dent Educ*. 2019 Apr;83(4):445–50.
39. Mathew MG, Samuel SR, Soni AJ, Roopa KB. Evaluation of adhesion of *Streptococcus mutans*, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary molars: randomized controlled trial [Internet]. Vol. 24, *Clinical Oral Investigations*. 2020. p. 3275–80. Available from: <http://dx.doi.org/10.1007/s00784-020-03204-9>
40. Crombie L, Josephs JL, Cayley J, Larkin J, Weston JB. The rotenoid core structure: Modifications to define the requirements of the toxophore. *Bioorg Med Chem Lett*. 1992 Jan 1;2(1):13–6.

41. Thomer L, Schneewind O, Missiakas D. Pathogenesis of *Staphylococcus aureus* Bloodstream Infections. *Annu Rev Pathol: Mech Dis*. 2016 May 23;11(1):343–64.
42. Stapels DAC, Ramyar KX, Ricklin D, Milder FJJ, Bischoff M, Herrmann M, et al. Extracellular adherence protein (Eap) of *Staphylococcus aureus* evades innate immunity by inhibiting complement activation and neutrophil elastase [Internet]. Vol. 217, *Immunobiology*. 2012. p. 1170. Available from: <http://dx.doi.org/10.1016/j.imbio.2012.08.118>
43. Prasad R. *Candida Albicans: Cellular and Molecular Biology*. Springer Science & Business Media; 2012. 267 p.
44. Srivastava S, Verma HN, Srivastava A, Prasad V. BDP-30, a systemic resistance inducer from *Boerhaaviadiffusa* L., suppresses TMV infection, and displays homology with ribosome-inactivating proteins. *J Biosci*. 2015 Mar;40(1):125–35.

UNDER PEER REVIEW