

**THE PREVALENCE OF ANTENATAL DEPRESSION AMONG WOMEN IN MEDINA, SAUDI ARABIA**

**Abstract**

**Background:** Antenatal depression is a recognized global burden. However, it has remained an ignored part of public health. Most of the maternal depression interventions have been directed towards post-natal depression.

**Purpose of the study:** The purpose of this study was to understand the third-trimester antenatal risk factors and prevalence among Saudi Arabian women.

**Method:** This study employed cross-sectional study design from April 2020 to June 2020. The collection of the data was through face-to-face interviews with pregnant women meeting eligibility criteria who were visiting the clinic of antenatal care at the Maternity and Children's Hospital. Data were collected using a structured and validated Patient Health Questionnaire-9 (PHQ-9). All the collected data were analyzed using SPSS software.

**Results:** Antenatal depression among participants stands at 35.8% for normal/ minimal depression, 31.3% for mild depression, 3.9% for moderate depression, 5.5% for moderately severe, and 3.5% for severe depression. Most of the risk factors for antenatal depression affect women mostly in the third trimester.

**Conclusion:** The antenatal depression prevalence among pregnant women in Medina is high. Screening for mental health problems and provision of supportive mental health services as a component of antenatal care services is recommended

**Keyword:** Depression, Women, Antenatal depression, Medina, Third trimester, Saudi Arabia.

## 1.0. Introduction

Depression is a global burden that is highly prevalent in pregnancy (1). Mothers transitioning through pregnancy to parenthood usually face psychological and social changes that involve anxiety (2). Up to 25% of women suffer from depression during pregnancy (2, 3). The early, mid (3<sup>rd</sup> trimester), and post pregnancy prevalence have been estimated at 15.5%, 11.1%, and 8.7% respectively (2). Pregnancy poses a high vulnerability for depression in women. Surprisingly, a lot of effort has been directed to understanding postpartum depression, ignoring pregnancy associated depression (1). Yet, by virtue of prevalence, pregnancy associated depression is more wanting to be addressed.

The depressive episodes that begin during pregnancy provide a basis for defining antenatal depression and usually predict possible postnatal depression (4, 5). Evidence shows that the incidence rate of depression in females with a known depression history is 42.5% and that in females without a known depression history is 21.7% (6, 7).

Depression affects the health, career, social, and family life of the affected pregnant women (8).

The manifestation of depression is linked to alterations in women's sex hormones and life stressors during or before pregnancy (8, 9). The antenatal depression risk factors are highly identical with those of any other form of depression at any one time. Research shows that the most common depression risk factors during pregnancy are perceived stress, life satisfaction, and social support (10). A review of 50 studies additionally showed that previous history of depression, life stressors, unplanned pregnancy, lower education, single marital status, Medicaid insurance, domestic violence, lack of social support, lower-income, and smoking significantly contribute to antenatal depression (11). Other studies have identified other risk factors for depression during pregnancy such as fear of childbirth, pregnancy issues, history of pregnancy loss, poor antenatal care, and poor nutrition (2, 12). In Saudi Arabia, there is little literature on depression during pregnancy and associated risk factors.

Antenatal depression is usually associated with significant psychological and medical issues that can affect both the baby and mother before and affect delivery (13). Research has mentioned that antenatal depression is highly associated with operative deliveries such as instrumental vaginal deliveries or C-sections, pre-eclampsia, use of epidural analgesics during delivery, postnatal depression, suicidal thoughts, and spontaneous preterm births (14). For the baby, the risk of low birth weight is highly associated with antenatal depression, decreased fetal movement, and even sudden death (15). Subsequently, the babies of depressed mothers stand a high likelihood of developing psychological and physical issues after birth (16, 17). Hence, carrying the effects forward to their pregnant life when they grow up. As such, increased early detection of antenatal depression and awareness with suitable psychological interventions particularly for women in the third trimester could decrease the high risk of adverse effects for the pregnant woman's life, family, and newborn child. The purpose of this study, therefore, was to understand the third-trimester antenatal depression risk factors and prevalence among Saudi Arabian women.

### **1.1. Objectives**

- To determine the antenatal depression prevalence for pregnant women attending antenatal care clinic in the Maternity and Children's Hospital in Medina, Saudi Arabia.
- To establish the risk factors for antenatal depression among pregnant women attending antenatal care clinic in the Maternity and Children's Hospital in Medina, Saudi Arabia.

### **2.0 Methods**

This study employed a descriptive cross-sectional study design. The study was conducted at Medina in the Maternity and Children's Hospital (MCH) from April 2020 to June 2020. The study participants were pregnant women who visited the outpatient clinic of antenatal care in MCH. Only women who were in their third trimester pregnancy were included in this study. Pregnant women who had a known familial history or already been diagnosed with depression before pregnancy were excluded from this study. The sample size was calculated using the formula:  $n = \frac{P(1-P) * Z^2}{d^2}$  where n is calculated sample size, p is expected proportion in population based on previous studies. Z is the z-

value for the selected level of confidence (95%), and  $d$  is absolute error or precision (0.05) (18). These participants were selected randomly. Every 5 to 10 participants per day who reported to the outpatient clinic for antenatal care at MCH and consented to be part of the study were enrolled in the study. This procedure was repeated daily until the sample size was reached.

An already structured Patient Health Questionnaire-9 (PHQ-9) (19) that was researcher self-administered was used to collect data from the participants. This questionnaire consisted of three parts; the socio-demographic data part, the obstetric history part, and the screening for depression among pregnant females part. This tool has previously been used in several other studies; hence, its validity and reliability are sound with high internal consistency. However, to understand this tool's reliability in the current study, it was piloted among 10 participants from a hospital different from the current study setting. Given the limitations of the Covid-19 pandemic, all the hospital Covid-19 initiated standard operating procedures were observed for safety of the researchers and the participants.

### **2.1. Data analysis**

Data were analyzed using Statistical Package for Social Sciences (SPSS) program version 22. Descriptive statistics were calculated for all the major variables and presented as frequencies. The independent student t-test was used to compare means of depression scores among participants in the first, second, and third trimesters. The statistical significance level was set at a p-value of  $<0.05$ .

### **3.0. Results**

As shown in table 1, most of the participants (56.7%) were in the third trimester, followed by 32.3% who were in the second trimester, and 10.9% in the first trimester. Seventy percent of the participants reported no disease associated with their pregnancies. Among those who reported at least a disease associated with their pregnancies, 5.5%, 5.5%, 4.0%, 1.5%, 1.5%, 1.0%, and 0.5% suffer from hypertension, hypothyroidism, Iron deficiency anemia, hypotension, chronic inflammations, asthma, and hyperthyroidism respectively. 68.2% of the participants reported having less than 5 pregnancies and 31.8% reported having 5 and more pregnancies. Regarding the number of children, a participant had, 81.1 % reported having less than 5 children while 18.9% have 5 and more children.

35.8% of the participants reported having no or minimal depression, 31.3% had mild depression, 23.9% had moderate depression, 5.5% had moderately severe depression, and 3.5% had severe depression. Concerning how the problems were affecting the participants, most of the participants (43.8%) reported no difficulty at all, 23.9% had somewhat difficulty, 19.4% had reported the problem being very difficult, and 12.9% reported the problem being extremely difficult to them.

**Table 1: Frequencies of the social-demographic variables**

<b>Variable</b>	<b>Frequency (n=201)</b>	<b>Percentage (%)</b>
<b>Trimester</b>		
1st trimester	22	10.9
2nd trimester	65	32.3
3rd trimester	114	56.7
<b>No diseases associated with pregnancy</b>	142	70.6
<b>Diseases associated with pregnancy</b>		
Diabetes	20	10
Hypertension	11	5.5
Hypothyroidism	11	5.5
Iron deficiency anemia	8	4.0
Hypotension	3	1.5
Chronic inflammations	3	1.5
Asthma	2	1.0
Hyperthyroidism	1	0.5
<b>Number of pregnancies</b>		
Less than 5 pregnancies	137	68.2
5 and more pregnancies	64	31.8
<b>Number of children per a woman</b>		

less than 5 children	163	81.1
5 and more children	38	18.9
<b>Levels of depression</b>		
None/minimal depression	72	35.8
Mild depression	63	31.3
Moderate depression	48	23.9
Moderately severe depression	11	5.5
Severe depression	7	3.5
<b>Impact of problem (depression)</b>		
not difficult at all	88	43.8
somewhat difficult	48	23.9
very difficult	39	19.4
extremely difficult	26	12.9

Table 2 shows that the participant's mean age was  $30.13 \pm 5.97$  years, the mean number of pregnancies was  $33.71 \pm 2.26$  pregnancy, and the mean total score of depression was  $7.33 \pm 5.36$ .

**Table2: Mean age and number of pregnancies and mean total score of depression of the participants**

Variable	Mean	Std. Deviation
Age of participants	30.13	5.975
Number of pregnancies	3.71	2.262
Total depression score	7.3383	5.36050

Table 3 shows that the mean score among pregnant women in the 1<sup>st</sup>, 2<sup>nd</sup>, and 3<sup>rd</sup> trimesters is  $5.86 \pm 4.47528$ ,  $6.18 \pm 4.39422$ , and  $8.28 \pm 5.84082$  respectively and this difference is statistically significant ( $p = .016$ ) on ANOVA. A post hoc analysis test was used to identify which exact pairs were different. Only the mean depression score of women in the 2<sup>nd</sup> and 3<sup>rd</sup> trimester pair was statically significant ( $p = 0.011$ ). Comparing the means of this pair, the participants in the 3<sup>rd</sup> trimester were more susceptible to depression compared to those in the 2<sup>nd</sup> trimester. Surprisingly, when participants with chronic illnesses in pregnancy, their number of pregnancies, and children were compared with

mean depression scores on t-tests, the findings were not statistically significant.

Table 3: Comparison of the mean depression scores among participants in the first, second, and third trimester of pregnancy

<i>Trimester</i>	<i>N</i>	<i>Mean</i>	<i>Std. Deviation</i>	<i>P-value</i>
1 <sup>st</sup>	22	5.8636	4.47528	.016
2 <sup>nd</sup>	65	6.1846	4.39422	
3 <sup>rd</sup>	114	8.2807	5.84082	
Total	201	7.3383	5.36050	

#### 4.0. Discussion

Determining the prevalence of and risk factors for antenatal depression among Saudi Arabian pregnant women were the specific objectives of this study. This determination is fundamental to the prevention and management of antenatal depression (1). Data from 201 participants were analyzed and the mean age of the participants was  $30.13 \pm 5.97$  years with the majority (56.7%) of the participants in the third trimester of pregnancy. This finding indicates that the majority of the pregnant women in Saudi Arabia that go for antenatal care are young women between 25 and 35 years. This observation resonates with the findings in studies elsewhere (20, 21) where the majority of the pregnant women seeking antenatal care are younger in age compared to the older ones.

In this study, the prevalence of antenatal depression stands at 31.3% in mild depression, 23.9% in moderate depression, 5.5% in moderately severe depression, and 3.5% in severe depression. In a study by Beyene et al. (20) where depression was not categorized, the prevalence of antenatal depression was found to be 24.45%. This observation is comparable to that of this current study. Additionally, Bunevicius et al. (22) found out that in their study 7.4% of the women had a diagnosis of a depressive disorder at least once during pregnancy. Depressive disorders have become a common disorder among pregnant women however with increasing prevalence in the third trimester. In the current study, most of the depressive disorders were significantly registered among participants in

the third trimester. In the study by Bunevicius et al. (22) this observation is linked to unplanned and unwanted pregnancy, high neuroticism, and occurrence of psychosocial stressors among women.

In the current study, the observed antenatal depression disorders have been associated with pregnancy related diseases and an increasing number of pregnancies. The majority (43.8%) of the participants reported that depression did not create any difficulty for them. This observation might be explained by the findings in the study by Beyene et al. (22) that indicate that antenatal depression is largely a neglected public health problem among pregnant women. Subsequently, Dadi et al. (23) in their study indicated that antenatal depression does not cause fatal outcomes has contributed largely to the low intervention priority among the potential victims. However, it's important to note that when not well attended to, antenatal depression increases the risk of adverse pregnancy outcomes, particularly preterm birth risk (23). This association can be explained in three ways: First, depression may cause the production of Hypothalamic-Pituitary-Adrenocortical Axis that stimulates the production of stress hormones such as cortisol. Cortisol interrupts adequate oxygen and nutrient flow to the fetus (22). Secondly, depression can disrupt the immune system functioning causing infections that affect fetal growth (20). Lastly, depressed mothers have a poor appetite and can even smoke and take alcohol that nutritionally affects the fetus (21).

Although not included in this study, evidence shows that other factors are associated with antenatal depression. These factors include; low level of education, low economic income, unplanned pregnancy, a history of psychological disorders like depression, anxiety, stress, and low social support during pregnancy (1, 21). Additionally, pregnant women from poor marital relationships and experiencing domestic violence are more susceptible to antenatal depression than the rest of the pregnant women (23).

#### **4.1. Conclusion**

The objectives of this study were met. The prevalence of antenatal depression among the participants in this study varies from 31.3% for mild depression, 23.9% for moderate depression, 5.5% for moderately severe depression, and 3.5% for severe depression. Although several factors can be associated with antenatal depression among women, in this study the third trimester of pregnancy significantly demonstrated being riskier to the development of antenatal depressive disorders compared to the first and second trimesters. Screening for mental health problems and provision of supportive mental health services as a component of antenatal care services is recommended.

#### **4.2. Limitations**

The study requires a larger sample size to ensure a representative distribution of the population.

#### **4.3. Recommendations**

Given the above findings and discussion, there is a need to screen all pregnant women for depression as part of routine antenatal care so that appropriate care is given to them. The health care system needs to consider antenatal depression a priority in obstetrics for better maternal and child health.

#### **7.0. Ethical consideration:**

The ethical approval was obtained from the Review Ethical Committee of the faculty of medicine, Taibah University. Additional approval was obtained from the administration of the Maternity and Children's Hospital. Informed consent was sought from the participants before interviewing them.

#### **COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

#### **9.0. References**

1. Míguez, M. C., & Vázquez, M. B. (2021). Risk factors for antenatal depression: A review. *World Journal of Psychiatry, 11*(7), 325–336. <https://doi.org/10.5498/wjp.v11.i7.325>

2. Ajinkya, S., Jadhav, P., & Srivastava, N. (2013). Depression during pregnancy: Prevalence and obstetric risk factors among pregnant women attending a tertiary care hospital in Navi Mumbai. *Industrial Psychiatry Journal*, 22(1), 37. <https://doi.org/10.4103/0972-6748.123615>
3. Rwakarema M, Premji S, Nyanza E, Riziki P, Palacios-Derflingher L. (2015). Antenatal depression is associated with pregnancy-related anxiety, partner relations, and wealth in women in Northern Tanzania: A cross-sectional study. *BMC Women's Health*. 15(1).
4. Orr, S., Blazer, D., James, S., & Reiter, J. (2007). Depressive Symptoms and Indicators of Maternal Health Status during Pregnancy. *Journal of Women's Health*. 16(4):535-542.
5. Klier, C. (2006). Mother–infant bonding disorders in patients with postnatal depression: The Postpartum Bonding Questionnaire in clinical practice. *Archives of Women's Mental Health*. 9(5):289-291.
6. Srivastava S, Bhatia M, Bohra N. (2015). Depression in women in Indian context. *Indian Journal of Psychiatry*. 57(6):239.
7. Al-Azri M, Al-Lawati I, Al-Kamyani R, Al-Kiyumi M, Al-Rawahi A, Davidson R et al. (2016). Prevalence and Risk Factors of Antenatal Depression among Omani Women in a Primary Care Setting: Cross-sectional study. *Sultan Qaboos University Medical Journal*. 16(1):e35-41.
8. Ashley, J., Harper, B., Arms-Chavez, C. and LoBello, S. (2015). Estimated prevalence of antenatal depression in the US population. *Archives of Women's Mental Health*, 19(2), pp.395-400.
9. Zeng, Y., Cui, Y., & Li, J. (2015). Prevalence and predictors of antenatal depressive symptoms among Chinese women in their third trimester: a cross-sectional survey. *BMC Psychiatry*. 15(1).
10. Bindt, C., Appiah-Poku, J., Te Bonle, M., Schoppen, S., Feldt, T., Barkmann, C., Koffi, M., Baum, J., Nguah, S., Tagbor, H., Guo, N., N'Goran, E. and Ehrhardt, S. (2012). Antepartum Depression and Anxiety Associated with Disability in African Women: Cross-Sectional Results from the CDS Study in Ghana and Côte d'Ivoire. *PLoS ONE*, 7(10), p.e48396.
11. Fisher J, Cabral de Mello M, Patel V, Rahman A, Tran T, & Holton S et al. (2011). Prevalence and determinants

- of common perinatal mental disorders in women in low- and lower-middle-income countries: a systematic review. *Bulletin of the World Health Organization*. 90(2):139-149H.
12. Shidhaye P, Shidhaye R, & Phalke V. (2017). Association of gender disadvantage factors and gender preference with antenatal depression in women: a cross-sectional study from rural Maharashtra. *Social Psychiatry and Psychiatric Epidemiology*. 52(6):737-748.
  13. Abdelhai R, & Mosleh H. (2015). Screening for antepartum anxiety and depression and their association with domestic violence among Egyptian pregnant women. *Journal of the Egyptian Public Health Association*. 90(3):101-108.
  14. O'Connor E, Rossom R, Henninger M, Groom H, & Burda B. (2016). Primary Care Screening for and Treatment of Depression in Pregnant and Postpartum Women. *JAMA*. 315(4):388.
  15. Akçalı Aslan P, Aydın N, Yazıcı E, Aksoy A, Kirkan T, & Daloglu G. (2014). Prevalence of depressive disorders and related factors in women in the first trimester of their pregnancies in Erzurum, Turkey. *International Journal of Social Psychiatry*. 2014;60(8):809-817.
  16. Friesen K, Peterson W, Squires J, & Fortier C. (2017). Validation of the Edinburgh Postnatal Depression Scale for Use with Young Childbearing Women. *Journal of Nursing Measurement*. 25(1):1-16.
  17. Fisher J, Cabral de Mello M, Patel V, Rahman A, Tran T, Holton S et al. (2011). Prevalence and determinants of common perinatal mental disorders in women in low- and lower-middle-income countries: a systematic review. *Bulletin of the World Health Organization*. 90(2):139-149H.
  18. Charan, J., & Biswas, T. (2013). How to calculate sample size for different study designs in medical research? *Indian Journal of Psychological Medicine*, 35(2), 121–126. <https://doi.org/10.4103/0253-7176.116232>
  19. Spitzer, L., Williams, W., & Kroenke, K. (n.d.). *Patient Health Questionnaire-9 (PHQ-9)*. Retrieved from <http://www.riss.kr/link?id=T13240980>
  20. Beyene, G. M., Azale, T., Gelaye, K. A., & Ayele, T. A. (2021). Depression remains a neglected public health problem among pregnant women in Northwest Ethiopia. *Archives of Public Health*, 79(1), 1–12.

<https://doi.org/10.1186/s13690-021-00649-6>

21. Mirieri, H. K., Mweu, M. M., & Olenja, J. M. (2020). Determinants of prenatal depression among women attending the antenatal clinic at a referral facility in Mombasa County, Kenya: a case control study. *F1000Research*, 9, 36. <https://doi.org/10.12688/f1000research.22017.2>
22. Bunevicius, R., Kusminskas, L., Bunevicius, A., Nadisauskiene, R. J., Jureniene, K., & Pop, V. J. M. (2009). Psychosocial risk factors for depression during pregnancy. *Acta Obstetrica et Gynecologica Scandinavica*, 88(5), 599–605. <https://doi.org/10.1080/00016340902846049>
23. Dadi, A. F., Miller, E. R., & Mwanri, L. (2020). Antenatal depression and its association with adverse birth outcomes in low and middleincome countries: A systematic review and meta-analysis. *PLoS ONE*, 15(1), 1–23. <https://doi.org/10.1371/journal.pone.0227323>