

# EVALUATION OF ANTIDIABETIC POTENTIAL OF ETHANOLIC EXTRACTS OF *CISSUS QUADRANGULARIS* AND *CINNAMOMUM TAMALA* - AN *IN VITRO* STUDY

## ABSTRACT

**BACKGROUND:** Diabetes is a rapidly growing metabolic disorder of the present generation. Uncontrolled diabetes leads to various complications which in turn affects the vascular system, neuropathy, and more. Medicinal plants are always a good source of drug equivalent currently present. Antidiabetic drugs lead to a large amount of side effects. A comparative study was made between *Cissus quadrangularis* and *Cinnamomum tamala* for its antidiabetic potential.

**AIM:** To evaluate the antioxidant and antidiabetic potential of ethanolic extract of *Cissus quadrangularis* and *Cinnamomum tamala* and to compare its efficacy.

**MATERIALS AND METHODS:** Ethanolic extracts of *Cissus quadrangularis* and *Cinnamomum tamala* were tested for its phytoconstituents, antioxidant and antidiabetic potential. The data were analyzed statistically by one-way analysis of variance (ANOVA) followed by Duncan's multiple range test and it was used to see the statistical significance among the groups. The results with the  $p < 0.05$  level were considered to be statistically significant.

**RESULT:** Phytochemical screening showed a strong presence of flavonoids and terpenoids in both the plant extract. Results showed that plant extract had antidiabetic and antioxidant activity. Among them *Cissus quadrangularis* exhibited significantly more antidiabetic and antioxidant activity.

**CONCLUSION:** The ethanolic extract of *Cissus* and *Cinnamomum tamala* exhibits potent antidiabetic and antioxidant properties. With further *in vivo* and *in vitro* studies the plant extract can be formulated into a potent antidiabetic drug.

**KEYWORDS:** Phytochemicals, antidiabetic potential, antioxidants, innovative technology, novel method.

**Running title:** Antidiabetic potential of *Cissus quadrangularis* and *Cinnamomum tamala*.

## INTRODUCTION

Diabetes is a growing problem worldwide causing enormous financial burden and maintaining care policy issues (Keter and Mutiso, 2012). According to the international diabetes federation (IDF) the number of individuals with diabetes in 2011 crossed 3.66 million with an estimation of 4.6 million death each year (Keter and Mutiso, 2012; Dong *et al.*, 2019). Indian subcontinent emerged as the capital of the diabetes epidemic. Indians show a significantly higher age related prevalence of diabetes when compared to other folks. Diabetes is characterized by metabolic dysregulation primarily of carbohydrate manifested by hyperglycemia leading to defects in insulin secretion, impaired insulin action or in many cases both the above mentioned stages. Untreated or uncontrolled diabetes leads to a plethora of complications which in turn affect the vascular disease, neuropathy, nephropathy. Diabetes is referred to as a chronic metabolic disorder that affects around 40.9 million population in India. Antidiabetic drugs such as biguanides and sulfonylureas can be used but they have prolonged side effects.

Medicinal plants have always been a good source of drugs and currently there are many drugs which are directly or indirectly extracted from them (Liu and Yaniv, 2005). The ethnobotanical information says that about 800 plant species possess antidiabetic potential among which *Momordica charantia*, *Pterocarpus marsupium*, *Cissus quadrangularis*, *Trigonella foenum graecum* are reported beneficial for treatment of diabetees.

Several herbs have shown antidiabetic activity when evaluated using different experimental techniques ('Antidiabetic Activity of Lansau Herb Extract in Glucose Induced Diabetic Male Rats', 2021). Wide arrays of plant derived active principles representing different kinds of biological activity has been explored. Among which alkaloids, glycosides, galactomannan, peptidoglycan, amino acids, terpenoids, glycoprotein, carbohydrates and other inorganic ions are demonstrated activity including treatment for diabetes.

Several classes of oral hypoglycemic drugs exert antidiabetic effect through different mechanisms like sulfonylureas, biguanides,  $\alpha$ -glucosidase inhibitors, thiazolidinediones and non-sulfonylureas secretagogues. Although synthetic oral hypoglycemic drugs alongside insulin are the main cause for controlling diabetes they fail to reverse the course of its complication completely and further worsen it by producing side effects (Ravindran, Nirmal-Babu and Shylaja, 2003; Singh *et al.*, 2015; Saboo, 2016; Scheen and Paquot, 2020). This forms the main alternative source discoveries for antidiabetic agents. Despite the significant progress made in treatment of diabetes using oral antidiabetic agents in the past 3 decades, the results of treatment of diabetic patients are still far from perfect. There are several disadvantages reported related to the use of oral hypoglycemic agents, including drug resistance which reduces the efficiency, adverse effect and even the toxicity (Ravindran, Nirmal-Babu and Shylaja, 2003; Singh *et al.*, 2015; Scheen and Paquot, 2020). Due to several limitations the search for newer antidiabetic drugs from natural sources continues.

One of the popular plants which helps in antidiabetic treatment is *Cissus quadrangularis* which belongs to the family Vitaceae and are distributed in hotter parts of India and Sri Lanka. The stem of the plant has been reputed in Ayurveda as an alternative, anthelmintic, dyspeptic, digestive, analgesic in eye and ear disease, irregular menstruation, asthmatic and mainly as antidiabetic. It also proves to reduce swelling and proves to promote the relevance of pain and healing of simple fractures as well as in curing the allied disorders associated with fracture (Muthusami *et al.*, 2011). Some phyto-constituents like flavonoids, phytosterol, tannins and triterpene present in the plant extract are responsible for antidiabetic activity. *Cinnamomum tamala*, Indian bay leaves which is also known as tejpat, is a tree which belongs to the Lauraceae family native to India, Nepal, Bhutan and China. They are commercially cultivated in certain parts of the country for leaf production and essential oils (Ravindran, Nirmal-Babu and Shylaja, 2003). Previous studies suggest that about 3g/d for 30 days decreases the risk factor of diabetes, cardiovascular disease and is beneficial for people with type-1 diabetes (Yang *et al.*, 2021). *Cinnamomum tamala* also possesses antidiabetic, gastroprotective, hypolipidemic, immunomodulatory, cytoprotective and antigenotoxic effects. The antidiabetic activity of *Cinnamomum tamala* is due to the presence of steroids, terpenoids, tannins and polysaccharides (Kumar, Vasudeva and Sharma, 2012). (Wu *et al.*, 2019), (Chen *et al.*, 2019), (Li *et al.*, 2020), (Babu and Jayaraman, 2020), (Malaikolundhan *et al.*, 2020), (Han *et al.*, 2019), (Gothai *et al.*, 2018), (Veeraraghavan, Hussain, *et al.*, 2021), (Sathya

*et al.*, 2020),(Yang *et al.*, 2020),(Rajendran *et al.*, 2020),(Barma *et al.*, 2021),(Samuel, 2021),(Samuel *et al.*, 2021),(Tang *et al.*, 2021),(Yin *et al.*, 2021),(Veeraraghavan, Periadurai, *et al.*, 2021),(Mickymaray *et al.*, 2021),(Teja and Ramesh, 2020),(Kadanakuppe and Hiremath, 2016). This research is carried out to evaluate the antioxidant and antidiabetic potential of ethanolic extract of *Cissus quadrangularis* and *Cinnamomum tamala* and to compare its efficacy(Ravindran, Nirmal-Babu and Shylaja, 2003; Singh *et al.*, 2015).

## **MATERIALS AND METHODS**

### **1. Phytochemical Screening test**

#### **Test for phlobatannin**

1ml of the extract was treated with 1ml of 1% HCl and boiled for 10 mins. The formation of red color precipitate indicates the presence of phlobatannin.

#### **Test for Carbohydrates**

Three to five drops of Molisch reagent was added with 1 mL of the extract and then 1 mL of concentrated sulphuric acid was added carefully through the side of the test tube. The mixture was then allowed to stand for two minutes and diluted with 5 mL of distilled water. The Development of a red or dull violet ring at the junction of the liquids showed the presence of carbohydrates.

#### **Test for Flavonoids**

Few drops of 1% liquid ammonia were taken in a test tube and along with it 1ml of the extract was added resulting in the formation of yellow color thereby indicating the presence of flavonoids.

#### **Test for Alkaloids**

2ml of sample was mixed with 2ml of HCl. Then 6 drops of HCN was added and further 2 drops of picric acid was added that resulted in a creamish pale yellow ppt indicating the presence of alkaloids.

#### **Test for Terpenoids**

2 ml of sample along with 2ml of chloroform and 3ml of con.  $H_2SO_4$  was added. Red color ppt obtained indicates the presence of terpenoids.

## **Test for proteins**

One milliliter of ninhydrin was dissolved in 1 mL of acetone and then a small amount of extract was added with ninhydrin. The formation of purple colour revealed the presence of protein.

## **Detection of saponins**

Foam test: A fraction of the extract was vigorously shaken with water and observed for persistent foam.

## **Test for steroids**

One milliliter of chloroform was mixed with 1 mL of extract and then ten drops of acetic anhydride and five drops of concentrated sulphuric acid were added and mixed. The formation of dark red colour or dark pink colour indicates the presence of steroids.

## **2. DPPH free radical scavenging activity of *Cissus quadrangularis* and *Cinnamomum Tamala***

Scavenging of 2, 2-Diphenyl-1-picrylhydrazyl (DPPH) radicals was assessed by the method of Hatano et al, (1989). DPPH solution (1.0 ml) was added to 1.0 ml of extract at different concentrations (0.1 to 0.5mg/ml). The mixture was kept at room temperature for 50 minutes and the activity was measured at 517 nm. Ascorbic acid at the same concentrations was used as standard. The capability to scavenge the DPPH radical was calculated and expressed in percentage (%) using following formula:

$$\text{DPPH radical scavenging (\%)} = \frac{\text{Control OD} - \text{Sample OD}}{\text{Control OD}} \times 100$$

## **3. *In vitro* antidiabetic activity**

### **3.1. *In vitro* alpha amylase inhibitory activity of *Cissus quadrangularis* and *Cinnamomum tamala***

$\alpha$ -amylase inhibitory activity of the extract was carried out according to the standard method of Ademiluyi *et al* [14]. In a test tube, reaction mixture containing 500 microliters of phosphate buffer (100 Mm, pH = 6.8), 100  $\mu$ l of  $\alpha$ -amylase (2 U/ml) and varying concentrations of *T. email* (0.1 to 0.5 mg/ml) was pre incubated at 3 degree Celsius for 20 min. Then, 200  $\mu$ l of 1% soluble starch (100mM phosphate buffer of pH = 6.8) was added as a substrate and incubated further at 37 degree Celsius for 30min; 1000 microliters of 3,5- dinitrosalicylic acid (DNS) color reagent

was then added and boiled for 10 min. The absorbance of the resulting mixture was measured at 540 nm using multiple readers. Acarbose at various concentrations (0.1-0.5 mg/ml) was used as standard. The results were expressed as percentage inhibition, which was then calculated using the formula,

$$\text{Inhibitory activity (\%)} = (1 - A_s/A_c) \times 100 \text{ Where,}$$

$A_s$  – absorbance in presence of test substance,  $A_c$  – absorbance of control

### 3.2. $\alpha$ -glucosidase inhibitory activity of *Cissus quadrangularis* and *Cinnamomum Tamala*

$\alpha$ -glucosidase inhibitory activity of extract was carried out according to the standard method of Ademiluyi et al [14]. In a test tube, reaction mixture containing 500  $\mu$ l of phosphate buffer (100 Mm, pH = 6.8), 100 microliters of  $\alpha$ -glucosidase (1 U/ml) and varying concentrations of *T. ammi* oil (0.1 to 0.5 mg/ml) was pre incubated at 37<sup>o</sup> C for 20 min. The reaction was stopped by adding 50  $\mu$ l of Na<sub>2</sub>CO<sub>3</sub> (0.1 M). The absorbance of the released p- nitrophenol was measured at 405 nm using multiple readers. Acarbose at various concentrations (0.1-0.5mg/ml) was used as a standard. The result was expressed as percentage inhibition which was calculated using the formula,

$$\text{Inhibitory activity (\%)} = (1 - A_s/A_c) \times 100 \text{ Where,}$$

$A_s$  – absorbance in presence of test substance,  $A_c$  – absorbance of control

### 4. Statistical Analysis

The data were subjected to statistical analysis using one – way analysis of variance (ANOVA) and Duncan’s multiple range test to assess the significance of individual variations between the groups. In Duncan’s test, significance was considered at the level of  $p < 0.05$ .

## RESULTS AND DISCUSSION:

Table1: Phytochemical analysis of Ethanolic extract of *Cissus quadrangularis* and *Cinnamomum tamala*

Phytochemical	<i>Cissus quadrangularis</i>	<i>Cinnamomum tamala</i>
Protein	+	+
Amino Acids	-	-

<b>Flavonoids</b>	+	-
<b>Alkaloids</b>	+	++
<b>Terpenoids</b>	+	+
<b>Steroids</b>	-	+
<b>Saponins</b>	+	+

**Figure 1:** *In vitro* antioxidant activity of ethanolic extracts of *Cissus quadrangularis* and *Cinnamomum tamala*




Figure 1 : Bar graph depicts the *In vitro* antioxidant activity of ethanolic extract of *Cissus quadrangularis* and *Cinnamomum tamala*. The X axis represents the different concentrations of *Cissus quadrangularis* and *Cinnamomum tamala* and the Y axis represents the percentage of inhibition. The blue colour denotes Vitamin C, the

orange colour denotes *Cissus quadrangularis* ethanolic extracts and purple denotes ethanolic extracts of *Cinnamomum tamala*. The difference was statistically significant. Each line represents Mean  $\pm$  SEM of 3 independent observations. Significance at  $p \leq 0.05$ .

Figure 2: *In vitro* alpha amylase inhibitory activity of ethanolic extracts of *Cissus quadrangularis* and *Cinnamomum tamala*




Figure 2 : Bar graph depicts the In vitro alpha amylase inhibitory activity of ethanolic extracts of *Cissus quadrangularis* and *Cinnamomum tamala*. The X axis represents the different concentrations of ethanolic extracts of *Cissus quadrangularis* and *Cinnamomum tamala* and the Y axis represents the percentage of inhibition. Blue colour denotes the concentration of the standard drug Metformin, red colour denotes the ethanolic extracts of *Cissus quadrangularis* and purple denotes the ethanolic extracts of *Cinnamomum tamala*. The difference was statistically significant. Each line represents Mean  $\pm$  SEM of 3 independent observations. Significance at  $p \leq 0.05$ .

Figure 3: *In vitro* alpha glucosidase inhibitory activity of ethanolic extracts of *Cissus quadrangularis* and *Cinnamomum tamala*




Figure 3 : Bar graph depicts the In vitro alpha glucosidase inhibitory activity of ethanolic extracts of *Cissus quadrangularis* and *Cinnamomum tamala*. The X axis represents the different concentrations of ethanolic extracts of *Cissus quadrangularis* and *Cinnamomum tamala* and the Y axis represents the percentage of inhibition. Blue colour denotes the concentration of the standard drug Metformin, orange colour denotes the ethanolic extracts of *Cissus quadrangularis* and green denotes the ethanolic extracts of *Cinnamomum tamala*. The difference was statistically significant. Each line represents Mean  $\pm$  SEM of 3 independent observations. Significance at  $p \leq 0.05$ .

The qualitative phytochemical analysis of *Cissus quadrangularis* ethanolic extract showed the presence of proteins, flavonoids, alkaloids, terpenoids and saponins whereas ethanolic extracts of *Cinnamomum tamala* showed the presence of proteins, alkaloids, terpenoids, steroids and saponins ( Table 1). Phytochemical screening is generally done for identification of medically active substances. The ethanolic extracts of *Cissus quadrangularis* and *Cinnamomum tamala* showed antioxidant activity ( figure 1), alpha amylase inhibitory activity ( figure 2) and alpha glucosidase inhibitory activity ( figure 3).

Low levels of insulin achieves adequate response and insulin resistance in target tissues. Some of the patients with type 2 diabetes are asymptomatic during their early phase of diabetes, but symptoms include polyuria, polydipsia, weight loss, blurred vision and polyphagia(Craig, Hattersley and Donaghue, 2009). Where uncontrolled diabetes leads to complications like stupor, coma, if not treated may even lead to death, due to the ketoacidosis or from nonketotic hyperosmolar syndrome.(Craig, Hattersley and Donaghue, 2009; Galtier, Brunet and Bringer, 2010). Diabetes is generally associated with patients with endocrine disease such as acromegaly,

cushing syndrome, glucagonoma and pheochromocytoma. It can also be seen in patients with genetic syndrome like down syndrome, klinefelter syndrome, turners syndrome and wolfram syndrome. As in discussion with prediabetes it correlates with increased cardiovascular mortality and cancer(Huang *et al.*, 2014).

The phytochemical analysis of the plant extract *Cissus quadrangularis* and *Cinnamomum tamal* were studied. Where the extracts exhibited a strong presence of saponins, terpenoids, flavonoids and alkaloids. Phytochemical screening refers to the identification, extraction and screening of medicinally active substances which are found in plants. They are widely used in order to enhance the values of particular plants for particular disease and disorders. Previous studies on the plant *Ephedra intermedia* for the presence of phytochemicals reveals the presence of tannins, terpenoids, glycosides and saponins (Gul *et al.*, 2017).

Oxidants are free radical compounds which cause harm if their levels are high in our body. They are linked to multiple illnesses like diabetes, heart disease and cancer. Antioxidants protect the cells from free radicals and play an important role in heart disease, cancer and other diseases. Free radicals are produced when the human body encounters breakdown of food and when they are exposed to toxins(Chakraborty and Mueen Ahmed, 2011). The most commonly known antioxidants are Vit C and Vit B. Ethanolic extract of the plants showed in vitro antioxidant activity in a concentration dependent manner where vit c is used as a standard drug . Antioxidant activity with  $IC_{50} = 320\mu\text{g/ml}$  and  $380\mu\text{g/ml}$  was exhibited by *Cissus quadrangularis* and *Cinnamomum tamal* respectively ( figure 1).

One of the antidiabetic therapeutic strategies is the inhibition of carbohydrate digesting enzymes like alpha amylase and alpha glucosidase. Where glucosidase enzymes lead to catalysis of starch to simple sugar and in humans they increase the blood glucose levels. In vitro antidiabetic activity was analysed for both the plant extracts and the result prove that ethanolic extract of *Cissus quadrangularis* showed a significantly increased activity compared to *Cinnamomum tamala* ( figure 2 and figure 3).

## **CONCLUSION:**

From the study it can be concluded that ethanolic extracts of *Cissus quadrangularis* and *Cinnamomum tamala* showed a potent antioxidant and antidiabetic activity which was evident

from DPPH scavenging activity, alpha amylase inhibitory activity and alpha glucosidase inhibitory activity respectively. A dose-dependent antidiabetic activity was observed from extracts and the standard drug Metformin. In the present study, the standard drug Metformin showed greater activity compared to extracts in all concentrations though the natural drugs tested showed potential, significantly so. Validation needs to be done on natural herbal extracts to make it into a potential alternative for synthetic drugs which possess a lot of side effects.

#### **COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

#### **REFERENCES**

‘Antidiabetic Activity of Lansau Herb Extract in Glucose Induced Diabetic Male Rats’ (2021) *International Journal of Pharmaceutical Research*. doi: 10.31838/ijpr/2021.13.02.094.

Babu, S. and Jayaraman, S. (2020) ‘An update on  $\beta$ -sitosterol: A potential herbal nutraceutical for diabetic management’, *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie*, 131, p. 110702.

Barma, M. D. *et al.* (2021) ‘Inhibition of *Streptococcus mutans*, antioxidant property and cytotoxicity of novel nano-zinc oxide varnish’, *Archives of oral biology*, 126, p. 105132.

Chakraborty, S. and Mueen Ahmed, K. K. (2011) ‘Introducing a New Journal: Free Radicals Antioxidants’, *Free Radicals and Antioxidants*, pp. 1–3. doi: 10.5530/ax.2011.1.1.

Chen, F. *et al.* (2019) ‘6-shogaol, a active constituents of ginger prevents UVB radiation mediated inflammation and oxidative stress through modulating Nrf2 signaling in human epidermal keratinocytes (HaCaT cells)’, *Journal of photochemistry and photobiology. B, Biology*, 197, p. 111518.

Craig, M. E., Hattersley, A. and Donaghue, K. C. (2009) ‘Definition, epidemiology and classification of diabetes in children and adolescents’, *Pediatric diabetes*, 10 Suppl 12, pp. 3–12.

Dong, H. *et al.* (2019) ‘Safety and feasibility of simultaneous endovascular therapy for supra-arch multivessel stenosis in 256 Chinese patients’, *Catheterization and cardiovascular interventions: official journal of the Society for Cardiac Angiography & Interventions*, 93(S1), pp. 846–850.

Galtier, F., Brunet, C. and Bringer, J. (2010) 'Diabète et grossesse', *Diabétologie*, pp. 305–316. doi: 10.1016/b978-2-294-70868-8.50017-7.

Gothai, S. *et al.* (2018) 'Pharmacological insights into antioxidants against colorectal cancer: A detailed review of the possible mechanisms', *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie*, 107, pp. 1514–1522.

Gul, R. *et al.* (2017) 'Preliminary Phytochemical Screening, Quantitative Analysis of Alkaloids, and Antioxidant Activity of Crude Plant Extracts from Ephedra intermedia Indigenous to Balochistan', *The Scientific World Journal*, pp. 1–7. doi: 10.1155/2017/5873648.

Han, X. *et al.* (2019) 'Anticarcinogenic potential of gold nanoparticles synthesized from *Trichosanthes kirilowii* in colon cancer cells through the induction of apoptotic pathway', *Artificial cells, nanomedicine, and biotechnology*, 47(1), pp. 3577–3584.

Huang, Y. *et al.* (2014) 'Prediabetes and the risk of cancer: a meta-analysis', *Diabetologia*, 57(11), pp. 2261–2269.

Kadanakuppe, S. and Hiremath, S. (2016) 'Social and Behavioural Factors Associated with Dental Caries Experience among Adolescent School Children in Bengaluru City, India', *British Journal of Medicine and Medical Research*, pp. 1–10. doi: 10.9734/bjmmr/2016/24021.

Keter, L. K. and Mutiso, P. C. (2012) 'Ethnobotanical studies of medicinal plants used by Traditional Health Practitioners in the management of diabetes in Lower Eastern Province, Kenya', *Journal of ethnopharmacology*, 139(1), pp. 74–80.

Kumar, S., Vasudeva, N. and Sharma, S. (2012) 'GC-MS analysis and screening of antidiabetic, antioxidant and hypolipidemic potential of *Cinnamomum tamala* oil in streptozotocin induced diabetes mellitus in rats', *Cardiovascular diabetology*, 11, p. 95.

Liu, C.-X. and Yaniv, Z. (2005) 'Research and Development of New Drugs Originating from Chinese Medicinal Plants', *Handbook of Medicinal Plants*, pp. 61–96. doi: 10.1201/9781482278026-4.

Li, Z. *et al.* (2020) 'Apoptotic induction and anti-metastatic activity of eugenol encapsulated chitosan nanopolymer on rat glioma C6 cells via alleviating the MMP signaling pathway', *Journal of Photochemistry and Photobiology B: Biology*, p. 111773. doi: 10.1016/j.jphotobiol.2019.111773.

Malaikolundhan, H. *et al.* (2020) 'Anticarcinogenic effect of gold nanoparticles synthesized from *Albizia lebbek* on HCT-116 colon cancer cell lines', *Artificial cells, nanomedicine, and biotechnology*, 48(1), pp. 1206–1213.

Mickymaray, S. *et al.* (2021) 'Rhaponticin suppresses osteosarcoma through the inhibition of PI3K-Akt-mTOR pathway', *Saudi journal of biological sciences*, 28(7), pp. 3641–3649.

Muthusami, S. *et al.* (2011) 'Cissus quadrangularis augments IGF system components in human osteoblast like SaOS-2 cells', *Growth Hormone & IGF Research*, pp. 343–348. doi:

10.1016/j.ghir.2011.09.002.

Rajendran, P. *et al.* (2020) 'Consumption of reused vegetable oil intensifies BRCA1 mutations', *Critical reviews in food science and nutrition*, pp. 1–8.

Ravindran, P. N., Nirmal-Babu, K. and Shylaja, M. (2003) *Cinnamon and Cassia: The Genus Cinnamomum*. CRC Press.

Saboo, B. (2016) 'Oral Hypoglycemic Agents: Drug Interaction and Toxicity', *Fundamentals of Diabetes*, pp. 239–239. doi: 10.5005/jp/books/12733\_20.

Samuel, S. R. (2021) 'Can 5-year-olds sensibly self-report the impact of developmental enamel defects on their quality of life?', *International journal of paediatric dentistry / the British Paedodontic Society [and] the International Association of Dentistry for Children*, 31(2), pp. 285–286.

Samuel, S. R. *et al.* (2021) 'Dental pain, parental SARS-CoV-2 fear and distress on quality of life of 2 to 6 year-old children during COVID-19', *International journal of paediatric dentistry / the British Paedodontic Society [and] the International Association of Dentistry for Children*, 31(3), pp. 436–441.

Sathya, S. *et al.* (2020) 'An in vitro study on hexavalent chromium [Cr(VI)] remediation using iron oxide nanoparticles based beads', *Environmental Nanotechnology, Monitoring & Management*, 14, p. 100333.

Scheen, A. J. and Paquot, N. (2020) '[Which injectable therapy after failure of oral antidiabetic agents in type 2 diabetes ?]', *Revue medicale de Liege*, 75(1), pp. 60–66.

Singh, V. *et al.* (2015) 'Combined potentiating action of phytochemical(s) from Cinnamomum tamala and Aloe vera for their anti-diabetic and insulinomimetic effect using in vivo rat and in vitro NIH/3T3 cell culture system', *Applied biochemistry and biotechnology*, 175(5), pp. 2542–2563.

Tang, Y. *et al.* (2021) 'Osteogenic differentiation and mineralization potential of zinc oxide nanoparticles from *Scutellaria baicalensis* on human osteoblast-like MG-63 cells', *Materials Science and Engineering: C*, p. 111656. doi: 10.1016/j.msec.2020.111656.

Teja, K. V. and Ramesh, S. (2020) 'Is a filled lateral canal – A sign of superiority?', *Journal of Dental Sciences*, pp. 562–563. doi: 10.1016/j.jds.2020.02.009.

Veeraraghavan, V. P., Hussain, S., *et al.* (2021) 'A Comprehensive and Critical Review on Ethnopharmacological Importance of Desert Truffles: *Terfezia clavaryi*, *Terfezia boudieri*, and *Tirmania nivea*', *Food Reviews International*, pp. 1–20. doi: 10.1080/87559129.2021.1889581.

Veeraraghavan, V. P., Periadurai, N. D., *et al.* (2021) 'Green synthesis of silver nanoparticles from aqueous extract of *Scutellaria barbata* and coating on the cotton fabric for antimicrobial applications and wound healing activity in fibroblast cells (L929)', *Saudi journal of biological sciences*, 28(7), pp. 3633–3640.

Wu, F. *et al.* (2019) 'Biologically synthesized green gold nanoparticles from Siberian ginseng induce growth-inhibitory effect on melanoma cells (B16)', *Artificial cells, nanomedicine, and biotechnology*, 47(1), pp. 3297–3305.

Yang H.-X. *et al.* (2021) '[Long-term follow-up of efficacy of insulin pump in the treatment of children with type 1 diabetes mellitus]', *Zhongguo dang dai er ke za zhi = Chinese journal of contemporary pediatrics*, 23(2), pp. 116–120.

Yang, Z. *et al.* (2020) 'Piperine loaded zinc oxide nanocomposite inhibits the PI3K/AKT/mTOR signaling pathway via attenuating the development of gastric carcinoma: In vitro and in vivo studies', *Arabian Journal of Chemistry*, 13(5), pp. 5501–5516.

Yin, Z. *et al.* (2021) 'Potential chemotherapeutic effect of betalain against human non-small cell lung cancer through PI3K/Akt/mTOR signaling pathway', *Environmental toxicology*, 36(6), pp. 1011–1020.

UNDER PEER REVIEW