

# Unintended pregnancy and associated factors of among pregnant women in Rural Ethiopia

## Abstract

**Background:** Unintended pregnancy is a worldwide problem that affects the health of women, families and relatives. This study was aimed to identify the major factors associated with unintended pregnancy among women in rural Ethiopia.

**Methods:** We used 2016 Ethiopian Demographic and Health Survey (EDHS) data on 853 rural pregnant women aged 15–49. Chi-square test was used to filter candidate independent variables to be included in a multivariable logistic regression. And the multivariable logistic regression model was employed to estimate the relative association of factors associated with unintended pregnancy

**Results:** The prevalence of unintended pregnancy in study area was 20.9% (95% CI: 18.2, 23.8%). The unintended pregnancy was associated with region Afar (AOR: 0.308, 95% CI: 0.101, 0.943) and Somalia (AOR: 0.064, 95% CI: 0.012, 0.331), women aged 35 or older (AOR: 2.016, 95% CI: 1.014, 4.009), women who had primary education (AOR: 0.284, 95% CI: 0.190, 0.425) and secondary/higher education (AOR: 0.201, 95% CI: 0.123, 0.329), Women who knew about contraceptives (AOR: 0.481, 95% CI: 0.379, 0.610), and parity 1 or 2 (AOR: 0.782, 95% CI: 0.750, 0.815).

**Conclusion:** About one-fifth of the pregnancies were unintended. Region, age, educational level, knowledge about contraceptive and parity were significantly associated factors of unintended pregnancy. Providing health education to women needs to be encouraged.

**Key Words:** Unintended pregnancy, pregnant women, rural Ethiopia

## 1. INTRODUCTION

Unintended pregnancy is defined as pregnancy that is either mistimed (pregnancy that occurred earlier than the desired time) or unwanted totally (pregnancy that occurred when no more children were desired) [1]. Sexual activity without using contraception through choice or coercion is the predominant cause of unplanned pregnancy [2].

Unintended pregnancy is a worldwide problem that affects the health of women, families and society [3]. Around 26.5 million unplanned pregnancies happened annually due to the incorrect use or failure of contraceptives globally [4]. In developing countries, unintended pregnancy accounted for 65% of all pregnancies and 59% of these pregnancies end with abortion [5].

Unintended pregnancy results in maternal mortality and morbidity because of the complications of unplanned births, unsafe abortion, and miscarriage which has a negative impact on health, economical and psychosocial costs to individual and society as well as significant emotional distress to women, families, and society [6]. The unintended pregnancy's impact is higher during the adolescent period, which levy to dropping out of

school, unstable and lack of proper management of family relationships [7]. In developing countries around 22000 women lost their lives because of the complication of abortion annually [8].

According to 2016 Ethiopian Demographic and Health Survey (EDHS) report, in Ethiopia, 25% of all births in the past five years and current pregnancies were unintended [9]. In Ethiopia, 24.7% and 42% of unintended pregnancy caused by contraceptive failure and not using contraceptive methods respectively [10].

Unintended pregnancy is affected by various factors such as educational status, knowledge on contraceptives, use of contraceptive methods, marital status, residence, religion, number of living children, age, parity, working status, economic status, family size, No of births in past 5 years [11, 12,13, 14, 15, 16,17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31].

Now days, one of the intervention areas for minimizing maternal mortality is preventing unintended pregnancy [32]. Identifying factors associated with unintended pregnancy is essential to inform the policy makers for the provision of safe and reliable service to reproductive-aged women. Hence, this study was aimed to identify factors associated with unintended pregnancy among pregnant women in rural Ethiopia based on 2016 Ethiopian Demographic and Health Survey data.

## **2. METHODS**

### **2.1. Data source**

This study used the 2016 Ethiopian demographic and health survey (EDHS) data, which was collected from January 18 to June 25, 2016. The 2016 EDHS was designed to provide estimates for the health and demographic variables of interest at national and regional level.

### **2.2.Variables in the study**

#### **2.2.1. Dependent variable**

Dependent variable is unintended pregnancy. The pregnancy is intended if it is planned at a time of survey and unintended if it is not planned at a time of survey or it is not planned at all.

#### **2.2.2. Independent variables**

Region, age, religion, marital status, educational status, husband/partner education, household wealth index, family size, working status, knowledge of contraceptive, sex of household head, parity, number of births in past five years, and ever terminated pregnancy

### **2.3. Statistical Data Analysis**

The data were analysed using statistical packages SPSS version 25. Frequencies and percentages were used to summarize the background characteristics of the women. And the Pearson's chi-square test of association was used to filter the significant independent variables that are associated to the dependent variable unintended pregnancy. A binary logistic regression model was fitted to fit the predictor variables associated with the unintended pregnancy. The model goodness of fit was checked using the Hosmer and Lemeshow test.

### 3. RESULTS

#### 3.1. Prevalence of unintended pregnancy

The overall prevalence of unintended pregnancy among pregnant women in rural Ethiopia was 20.9 % (95% CI: 18.2, 23.8%).

#### 3.2. Background characteristics of pregnant women

Of all 853 women included in the study, 138(16.2%) of them were living in Oromia region. Nearly half (48.3%) of women were in the age range of 25-34 years. More than three fifth (65.4%) of the women did not attend the formal education. Majority (97.2%) of them were in married/ living together at a time of survey. Majority (63.9%) of the women were from the households with the poor wealth index. Nearly three fifth (58.4%) of the women were Muslim religion followers. 254(29.8%) women had five or more ever born children. Majority (620(72.7%)) of women had 3 to 4 births in the last five years. More than three fourth (719(84.3%)) women were from household headed by males. More than half (490(57.4%)) of women were from family of size 5 or less. Only 86(10.1%) women ever had a terminated pregnancy. Majority (713(83.6%)) of women know about the contraceptive (Table 1).

Table1: Background characteristics of pregnant women in Ethiopia (n=853)

<b>Variables</b>	<b>Frequency (%)</b>
<b>Region</b>	
Tigray	66(7.7)
Afar	114 (13.4)
Amhara	82(9.6)
Oromia	138(16.2)
Somalia	135(15.8)
Benshangul Gummuz	73(8.6)
SNNPR	128(15.0)
Gambela	41(4.8)
Harari	50(5.9)
Dire Dawa	26(3.0)
<b>Age</b>	
15-24	308(36.1)
25-34	412(48.3)
25+	133(15.6)
<b>Educational level</b>	
No education	558(65.4)
Primary	249(29.2)
Secondary/Higher	46(5.4)
<b>Marital status</b>	
Never in union	7(0.8)
Married/living together	829(97.2)
Widowed/divorced/separated	17(2.0)

<b>Wealth index</b>	
Poor	545(63.9)
Middle	138(16.2)
Rich	170(19.9)
<b>Religion</b>	
Orthodox	188(22.0)
Protestant	142(16.6)
Muslim	498(58.4)
Others	25(2.9)
<b>Parity</b>	
0	125(14.7)
1-2	246(28.8)
3-4	228(26.7)
5+	254(29.8)
<b>Number of births in last five years</b>	
No births	196(23.0)
1-2	620(72.7)
3-4	37(4.3)
<b>Sex of household</b>	
Male	719(84.3)
Female	134(15.7)
<b>Family size</b>	
5 or less	490(57.4)
More than 5	363(42.6)
<b>Ever had a terminated pregnancy</b>	
No	767(89.9)
Yes	86(10.1)
<b>Knowledge about contraceptive</b>	
No	140(16.4)
Yes	713(83.6)

### 3.3. Unintended pregnancy by characteristics of women

Among 178 women who had unintended pregnancy, 48 (27.0%) of them were living in Oromia region at a time of survey. 51.1% of the unintended pregnancy occurred among women who were in the age groups of 25 to 34 years. 111(62.4%) unintended pregnancy occurred among women who did not attend formal education. 104(58.4%) unintended pregnancy occurred among women with poor wealth index. 78(43.8%) unintended pregnancy occurred among Muslim religion follower women. 69(38.8%) unintended pregnancy occurred among women who had ever born 5 or more children. 150(84.3%) unintended pregnancy occurred among women who had 1 or 2 births in last five years. 90(50.6%) unintended pregnancy occurred among women from families of size 5 or less. 160(89.9%) unintended pregnancy occurred among women who had no ever terminated pregnancy. 62(34.8%) unintended pregnancy occurred among women who did not know contraceptive methods (Table 2).

Table 2: Unintended pregnancies by characteristics of women in Ethiopia (n = 853)

Variables	Unintended pregnancy		Chi-square value(significance)
	Yes	No	
<b>Region</b>			83.250(0.000)
Tigray	14(7.9%)	52(7.7%)	
Afar	8(4.5%)	106(15.7%)	
Amhara	21(11.8%)	61(9.0%)	
Oromia	48 (27.0%)	90(13.3%)	
Somalia	2(1.1%)	133(19.7%)	
Benshangul Gummuz	10(5.6%)	63(6.3%)	
SNNPR	45 (25.3%)	83(12.3%)	
Gambela	11(6.2%)	30(4.4%)	
Harari	10(5.6%)	10(5.9%)	
Dire Dawa	9(5.1%)	17(2.5%)	
<b>Age</b>			19.035(0.000)
15-24	44(24.7%)	264(39.1%)	
35-34	91(51.1%)	321(47.6%)	
25+	43(24.2%)	90(13.3%)	
<b>Educational level</b>			9.869(0.020)
No education	111(62.4%)	447(66.2%)	
Primary	59(33.1%)	190(28.1%)	
Secondary/ Higher	8(4.5%)	38(5.7%)	
<b>Marital status</b>			0.257(0.880)
Never in union	1(0.6%)	6(0.9%)	
Married/living together	173(97.2%)	656(97.2%)	
Widowed/divorced/separated	4(2.2%)	13(1.9%)	
<b>Wealth index</b>			4.285(0.117)
Poor	104(58.4%)	441(65.3%)	
Middle	29(16.3%)	109(16.1%)	
Rich	45(25.3%)	125(18.5%)	
<b>Religion</b>			20.306(0.000)
Orthodox	50(28.1%)	138(20.4%)	
Protestant	42(23.6%)	100(14.8%)	
Muslim	78(43.8%)	420(62.2%)	
Others	8(4.5%)	17(2.5%)	
<b>Parity</b>			22.692(0.000)
0	9(5.1%)	116(17.2%)	
1-2	45(25.3%)	201(29.8%)	
3-4	55(30.9%)	173(25.6%)	
5+	69(38.8%)	185(27.4%)	
<b>Number of births in last five years</b>			15.675(0.000)
No births	22(12.4%)	174(25.8%)	
1-2	150(84.3%)	470(69.6%)	
3-4	6(3.4%)	31(4.6%)	
<b>Sex of household</b>			1.906(0.167)
Male	156(87.6%)	563(83.4%)	
Female	22(12.4%)	112(16.6%)	
<b>Family size</b>			4.359(0.037)
5 or less	90(50.6%)	400(59.3%)	
More than 5	88(49.4%)	275(40.7%)	
<b>Ever had a terminated</b>			0.000(0.988)

<b>pregnancy</b>			
No	160(89.9%)	607(89.9%)	
Yes	18(10.1%)	68(10.1%)	
<b>Knowledge about contraceptive</b>			14.380(0.000)
No	62(34.8%)	78(11.6%)	
Yes	116(65.2%)	597(88.4%)	

### 3.4. Factor associated with unintended pregnancy

The multivariable logistic regression analysis showed that region, age, education, parity and Knowledge about contraceptive were significantly associated factors with unintended pregnancy.

The odds of unintended pregnancy among women who were living in Afar region and Somalia region were 0.308 (AOR: 0.308, 95% CI: 0.101, 0.943) and 0.064 (AOR: 0.064, 95% CI: 0.012, 0.331) respectively times lower than women who were living in Tigray region. Women aged 35 or older (AOR: 2.016, 95% CI: 1.014, 4.009) were more likely to have unintended pregnancy than women aged 15-24 years. Women who had primary education (AOR: 0.284, 95% CI: 0.190, 0.425) and secondary/higher education (AOR: 0.201, 95% CI: 0.123, 0.329) were less likely to have unintended pregnancy than women who did not attend formal education. Women who had 1 or 2 ever born children (AOR: 0.782, 95% CI: 0.750, 0.815) were less likely to have unintended pregnancy than women who had no children. Women who knew about contraceptives (AOR: 0.481, 95% CI: 0.379, 0.610) were less likely to have unintended pregnancy than women who did not know about contraceptives (Table 3).

Table 3: Factors associated with unintended pregnancy in rural Ethiopia from multivariable logistic regression analysis

Variables	AOR	(95% CI for AOR)		p-value
<b>Region</b>				
Tigray	1			
Afar	0.308	0.101	0.943	0.029*
Amhara	1.234	0.548	2.781	0.611
Oromia	2.038	0.866	4.798	0.103
Somalia	0.064	0.012	0.331	0.001*
Benshangul Gummuz	0.535	0.196	1.462	0.223
SNNPR	2.134	0.917	4.967	0.179
Gambela	1.315	0.447	3.869	0.618
Harari	0.894	0.298	2.684	0.842
Dire Dawa	2.103	0.640	6.912	0.221
<b>Age</b>				
15-24	1			
35-34	1.241	0.574	2.686	0.583
35+	2.016	1.014	4.009	0.025*
<b>Educational level</b>				
No education	1			
Primary	0.284	0.190	0.425	0.022*

Secondary/ Higher	0.201	0.123	0.329	0.007*
<b>Religion</b>				
Orthodox	1			
Protestant	0.704	0.361	1.372	0.302
Muslim	0.743	0.401	1.377	0.345
Others	1.183	0.415	3.370	0.753
<b>Parity</b>				
0	1			
1-2	0.782	0.750	0.815	0.010*
3-4	0.512	0.185	1.419	0.129
5+	0.293	0.073	1.169	0.081
<b>Number of births in last five years</b>				
No births	1			
1-2	2.619	0.574	11.947	0.214
3-4	1.813	0.520	6.317	0.350
<b>Family size</b>				
5 or less	1			
More than 5	0.981	0.606	1.588	0.938
<b>Knowledge about contraceptive</b>				
No	1			
Yes	0.481	0.379	0.610	0.036*

\*= significant at 5% level of significance

#### 4. DISCUSSION

The prevalence of unintended pregnancy among pregnant women in the study area was 20.9%, (95% CI: 18.2, 23.8%). This finding was in line with studies conducted in Gondar (20.6%) [27]. However, it is higher than the study conducted in Felege Hiwot referral hospital, Bahir Dar (13.7%) [33], whereas it is lower than the study conducted in Angola (38.3%) [34], Uganda (37%) [35], Welkaite Woreda (26%) [36], Kersa (27.9%) [37], Debre Birhan (23%) [10], Nepal (54.5%) [38], Arsi Negele Woreda (41.5%) [25]. The variation in the prevalence might be due to socio-demographic and cultural differences of participants and health service coverage.

In this study, the factors that were significantly associated with unintended pregnancy were region, age, education, parity and knowledge about contraceptive.

Women living in Afar and Somalia were less likely to have unintended pregnancy compared to women living in Tigray region. This finding is consistent with a study conducted in Ethiopia [18] and Ghana [39]. This might be due to fertility desire difference as Afar and Somalia are underdeveloped regions where the pregnancies may be appreciated and accepted compared to Tigray region in the context of Ethiopia.

Women aged 35 and above were more likely to have unintended pregnancy than women aged 15-24 years. This is consistent with the studies in Jimma [28], sub-saharan Africa [31], Arsi Negele Woreda [25]. This might be due to that when women get older, they may not want bear children. But this is inconsistent with study in Kenya [14] and Brazil [40] which found that young women were more likely to experience unintended pregnancy than older women.

Women who had primary education and secondary/higher education were less likely to have unintended pregnancy than women who did not attend formal education. This is consistent with the studies in Ethiopia [41], Ghana [39] and sub-Saharan Africa [31]. This might be due to the fact that educated women may know about the methods of preventing the pregnancies compared to uneducated women. Similarly, Women who knew about contraceptives were less likely to have unintended pregnancy than women who did not know about contraceptives. This is consistent with the studies in Jimma [28] and sub-Saharan Africa [31].

Women with parity of 1 or 2 were less likely to have unintended pregnancy than women who had no children. This is consistent with the studies in Arsi Negele Woreda [25], Debre Birhan [10], Ethiopia [12, 18, 26, 41] and six south Asian countries [15]. This might be because of that women who have had children may care about the next pregnancy.

## **5. CONCLUSIONS**

The study was aimed to identify factors that are associated with unintended pregnancy among pregnant women in Ethiopia based on EDHS 2016 data. Region, age, educational level, knowledge about contraceptive and parity were significant factors associated with unintended pregnancy among pregnant women in Ethiopia. Providing health education to women is important to prevent unintended pregnancy.

## **ETHICS APPROVAL AND CONSENT TO PARTICIPATE**

Ethical approval was not applicable

## **CONSENT FOR PUBLICATION**

Not applicable.

## **ABERRATIONS**

AOR: Adjusted odds ratio

CSA: Central Statistical Agency

DHS: Demographic and Health Survey

EDHS: Ethiopian Demographic and Health Survey

SNNPR: Southern Nations, Nationalities and People's Region

WHO: World Health Organization

## **COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## **REFERENCES**

1. WHO, *Unintended pregnancy; improving maternal and child health*, World Health Organization, 2015.

2. Moges, Y., Worku, S.A., Niguse, A. and Kelkay, B., 2020. Factors associated with the unplanned pregnancy at Suhul General Hospital, Northern Ethiopia, 2018. *Journal of Pregnancy, 2020*.
3. Klima, C.S., 1998. Unintended pregnancy: consequences and solutions for a worldwide problem. *Journal of Nurse-Midwifery, 43*(6), pp.483-491.
4. World Health Organization, 2015. *Trends in maternal mortality: 1990-2015: estimates from WHO, UNICEF, UNFPA, World Bank Group and the United Nations Population Division*. World Health Organization.
5. Bearak, J., Popinchalk, A., Alkema, L. and Sedgh, G., 2018. Global, regional, and subregional trends in unintended pregnancy and its outcomes from 1990 to 2014: estimates from a Bayesian hierarchical model. *The Lancet Global Health, 6*(4), pp.e380-e389.
6. Kabir, S.M., 1989. Causes and consequences of unwanted pregnancy from Asian women's perspectives. *International Journal of Gynecology & Obstetrics, 30*, pp.9-14.
7. World Health Organization (WHO). Adolescent pregnancy. Geneva: World Health Organization; 2013.
8. Sedgh, G., Bearak, J., Singh, S., Bankole, A., Popinchalk, A., Ganatra, B., Rossier, C., Gerdtz, C., Tunçalp, Ö., Johnson Jr, B.R. and Johnston, H.B., 2016. Abortion incidence between 1990 and 2014: global, regional, and subregional levels and trends. *The Lancet, 388*(10041), pp.258-267.
9. Central Statistical Agency (CSA) [Ethiopia] and ICF. Ethiopia Demographic and Health Survey 2016. Addis Ababa: CSA and ICF; 2016
10. Melese, K.G., Gebrie, M.H., Badi, M.B. and Mersha, W.F., 2016. Unintended pregnancy in Ethiopia: community based cross-sectional study. *Obstetrics and gynecology international, 2016*.
11. Obare, F., van der Kwaak, A. and Birungi, H., 2012. Factors associated with unintended pregnancy, poor birth outcomes and post-partum contraceptive use among HIV-positive female adolescents in Kenya. *BMC women's health, 12*(1), pp.1-8.
12. Habte, D., Teklu, S., Melese, T. and Magafu, M.G., 2013. Correlates of unintended pregnancy in Ethiopia: results from a national survey. *PloS one, 8*(12), p.e82987.
13. Merga, J., Wirtu, D., Bekuma, T.T. and Regasa, M.T., 2021. Unintended pregnancy and the factors among currently pregnant married youths in Western Oromia, Ethiopia: A mixed method. *PloS one, 16*(11), p.e0259262.
14. Ikamari, L., Izugbara, C. and Ochako, R., 2013. Prevalence and determinants of unintended pregnancy among women in Nairobi, Kenya. *BMC pregnancy and childbirth, 13*(1), pp.1-9.
15. Sarder, A., Islam, S.M.S., Talukder, A. and Ahammed, B., 2021. Prevalence of unintended pregnancy and its associated factors: Evidence from six south Asian countries. *PLoS One, 16*(2), p.e0245923.
16. Mohamed, E.A.E.B., Hamed, A.F., Yousef, F.M. and Ahmed, E.A., 2019. Prevalence, determinants, and outcomes of unintended pregnancy in Sohag district, Egypt. *Journal of the Egyptian Public Health Association, 94*(1), pp.1-9.

17. Hamdela, B., G/mariam, A. and Tilahun, T., 2012. Unwanted pregnancy and associated factors among pregnant married women in Hosanna Town, Southern Ethiopia. *PloS one*, 7(6), p.e39074.
18. Zeleke, L.B., Alemu, A.A., Kassahun, E.A., Aynalem, B.Y., Hassen, H.Y. and Kassa, G.M., 2021. Individual and community level factors associated with unintended pregnancy among pregnant women in Ethiopia. *Scientific Reports*, 11(1), pp.1-9.
19. Lamina, M.A., 2015. Prevalence and determinants of unintended pregnancy among women in South-Western Nigeria. *Ghana medical journal*, 49(3), pp.187-194.
20. Ahinkorah, B.O., 2020. Individual and contextual factors associated with mistimed and unwanted pregnancies among adolescent girls and young women in selected high fertility countries in sub-Saharan Africa: A multilevel mixed effects analysis. *PloS one*, 15(10), p.e0241050.
21. Haffejee, F., Govender, N., Reddy, P., Sibiyi, M.N., Ghuman, S., Ngxongo, T., Borg, D. and O'Connor, L., 2018. Factors associated with unintended pregnancy among women attending a public health facility in KwaZulu-Natal, South Africa. *South African Family Practice*, 60(3), pp.79-83.
22. Alene, M., Yismaw, L., Berelie, Y., Kassie, B., Yeshambel, R. and Assemie, M.A., 2020. Prevalence and determinants of unintended pregnancy in Ethiopia: A systematic review and meta-analysis of observational studies. *PloS one*, 15(4), p.e0231012.
23. Moore, A.M., Gebrehiwot, Y., Fetters, T., Wado, Y.D., Bankole, A., Singh, S., Gebreselassie, H. and Getachew, Y., 2016. The estimated incidence of induced abortion in Ethiopia, 2014: changes in the provision of services since 2008. *International perspectives on sexual and reproductive health*, 42(3), p.111.
24. Nigussie, K., Degu, G., Chanie, H. and Edemealem, H., 2021. Magnitude of Unintended Pregnancy and Associated Factors Among Pregnant Women in Debre Markos Town, East Gojjam Zone, Northwest Ethiopia: A Cross-Sectional Study. *International Journal of Women's Health*, 13, p.129.
25. Fite, R.O., Mohammedamin, A. and Abebe, T.W., 2018. Unintended pregnancy and associated factors among pregnant women in Arsi Negele Woreda, West Arsi Zone, Ethiopia. *BMC research notes*, 11(1), pp.1-7.
26. Teshale, A.B. and Tesema, G.A., 2020. Magnitude and associated factors of unintended pregnancy in Ethiopia: a multilevel analysis using 2016 EDHS data. *BMC Pregnancy and Childbirth*, 20, pp.1-8
27. Yenealem, F. and Niberet, G., 2019. Prevalence and associated factors of unintended pregnancy among pregnant woman in Gondar town, North west Ethiopia, 2014. *BMC research notes*, 12(1), pp.1-5.
28. Beyene, G.A., 2019. Prevalence of unintended pregnancy and associated factors among pregnant mothers in Jimma town, southwest Ethiopia: a cross sectional study. *Contraception and reproductive medicine*, 4(1), pp.1-8.
29. Nigussie, K., Degu, G., Chanie, H. and Edemealem, H., 2021. Magnitude of Unintended Pregnancy and Associated Factors Among Pregnant Women in Debre Markos Town, East Gojjam Zone, Northwest Ethiopia: A Cross-Sectional Study. *International Journal of Women's Health*, 13, p.129.

30. Gite, A., Liulseged, N., Seyife, H., Abrha, Y., Workineh, Y. and Shegaze, M., 2016. Unintended pregnancy: magnitude and associated factors among pregnant women in Arba Minch Town, Gamo Gofa Zone, Ethiopia, 2015. *Reprod Syst Sex Disord*, 5(193), pp.1-6.
31. Ameyaw, E.K., Budu, E., Sambah, F., Baatiema, L., Appiah, F., Seidu, A.A. and Ahinkorah, B.O., 2019. Prevalence and determinants of unintended pregnancy in sub-Saharan Africa: A multi-country analysis of demographic and health surveys. *PloS one*, 14(8), p.e0220970.
32. Howden-Chapman, P., Siri, J., Chisholm, E., Chapman, R., Doll, C.N. and Capon, A., 2017. SDG 3: Ensure healthy lives and promote wellbeing for all at all ages. *A guide to SDG interactions: from science to implementation*. Paris, France: International Council for Science, pp.81-126.
33. Gebreamlak, W., Aragaw, A., Lemma, S. and Demilew, W., 2014. Magnitude and factors influencing unintended pregnancy among pregnant women attending antenatal care at Felege Hiwot referral hospital, Northwest Ethiopia: a cross-sectional study. *Science Journal of public health*, 2(4), pp.261-269.
34. Yaya, S. and Ghose, B., 2018. Prevalence of unmet need for contraception and its association with unwanted pregnancy among married women in Angola. *PloS one*, 13(12), p.e0209801.
35. Wasswa, R., Kabagenyi, A. and Atuhaire, L., 2020. Determinants of unintended pregnancies among currently married women in Uganda. *Journal of Health, Population and Nutrition*, 39(1), pp.1-17.
36. Abayu, H., Birhanu, Z., Nega, A. and Kidanemariam, A., 2015. Prevalence and associated factors of unintended pregnancy in Welkaite Woreda, Tigray and North Ethiopia cross sectional study by 2012. *J Preg Child Health*, 2(2), pp.1-5.
37. Kassa, N., Berhane, Y. and Worku, A., 2012. Predictors of unintended pregnancy in Kersa, Eastern Ethiopia, 2010. *Reproductive health*, 9(1), pp.1-7.
38. Bastola, K., Neupane, S., Hadkhale, K. and Kinnunen, T.I., 2015. Unintended pregnancy among married pregnant women in Nepal. *J Womens Health, Issues Care* 4, 4, p.2.
39. Nyarko, S.H., 2019. Unintended pregnancy among pregnant women in Ghana: prevalence and predictors. *Journal of pregnancy*, 2019.
40. Theme-Filha, M.M., Baldisserotto, M.L., Fraga, A.C.S.A., Ayers, S., da Gama, S.G.N. and do Carmo Leal, M., 2016. Factors associated with unintended pregnancy in Brazil: cross-sectional results from the Birth in Brazil National Survey, 2011/2012. *Reproductive health*, 13(3), pp.235-243.
41. Tebekaw, Y., Aemro, B. and Teller, C., 2014. Prevalence and determinants of unintended childbirth in Ethiopia. *BMC pregnancy and childbirth*, 14(1), pp.1-9.