

## **PROBIOTICS IN DENTAL CARIES PREVENTION**

### **ABSTRACT:**

Background: Changing lifestyle has resulted in deterioration of oral health in people of all age groups. The increasing global problems with the traditional disease management strategies have prompted the investigators to be in constant search of alternatives for managing these health issues.

Highlight: Probiotics is one such promising alternative to traditional disease management that has gained a lot of importance in improving oral health. The effect of probiotics on dental caries and its related risk factors is gaining momentum in the current era.

Conclusion: This narrative review summarised the available literature on probiotics in caries prevention.

Keywords: Bifidobacterium, Dental Caries, Oral health, Probiotics, Lactobacillus, Streptococcus

### **INTRODUCTION:**

Dental caries still remains one of the most common diseases worldwide, although a decline of the prevalence has been recorded in western countries (1). The disease is triggered by the interaction over time among cariogenic microorganisms (mainly mutans streptococci and lactobacilli), a diet rich in fermentable carbohydrates and host factors, like as saliva secretion rate and buffering capacity (2). When sugared food/drinks are supplied frequently, acidogenic and aciduric strains increase selectively in the oral environment. These changes, over time, shift the demineralization/remineralization balance toward net mineral loss, leading to the caries lesion development (3). The microbial community of caries is diverse

and contains many facultatively and obligately-anaerobic bacteria (4). Mutans streptococci (MS) have been considered for a long time the major pathogens involved in caries development (5). The main characteristics of virulence of the Streptococcus mutans are their acidogenicity, ability to survival in acidic environments, the ability of biofilm formation and adherence to the tooth. In recent years, it was described that the microflora on the tooth surface changes with caries lesion development, from a predominance of non-mutans streptococci and Actinomyces spp. to dominance of MS and other non-mutans bacteria, including lactobacilli and Bifidobacterium spp (6).

### **CARIES PREVENTION:**

Prevention of dental caries is based on the reduction in the number of cariogenic bacteria which leads to the environment that facilitates remineralization. There are several conventional methods agents to prevent occurrence of dental caries. Many agents have been shown to have an anticariogenic effect or to have the potential for such an effect. They can be categorised by their modes of action into three groups: those which affect plaque and plaque bacteria, those which affect tooth enamel chemistry and those which buffers oral pH (7). According to literature the most commonly used agents are chemoprophylactic agents, antimicrobial peptides, sugar substitutes like xylitol, fluoride therapy and casein phosphopeptides (8-13)

Preventive strategies are needed and recommended to control caries risk factors mainly based on dietary changes i.e., sweeteners intake reduction and enhancing host resistance (14). Sometimes, antibacterial agents are administered in order to reduce cariogenic micro-flora, however, a complete eradication of caries-associated microorganisms has proved to be difficult and almost impossible to obtain (15). Rather it might eradicate the entire oral flora that leads to undesirable effects. In order to overcome this drawback of

conventional methods, selective inhibition methods of pathogens responsible of dental caries are being studied at recent times (16). One such approach is use of probiotics against dental caries.

### **DEFINING THE TERM PROBIOTICS:**

According to WHO the term probiotics is defined as, "Live microorganisms which, when administered in adequate amounts, confer a health benefit on the host" (17). These microorganisms belong to the natural human flora in order to survive in the acid environment during transit to the intestines. One of the main probiotic health claims is the positive change in microbiota composition traditionally investigated in regard to intestinal health (18). Probiotics are recognized to perform several actions in the digestive system as to prevent cellular adhesion and invasion of pathogenic bacteria, modify the intestinal environment and modulating the local and systemic inflammatory immune response (19). In the last decade, the use of probiotics has generated interest within the dental community with the development of studies focused on reducing caries incidence (20).

The effect of probiotics on dental caries and its related risk factors has been evaluated in several experimental studies, using different strains; *Lactobacillus rhamnosus* GG, *L. casei*, *L. reuteri*, *L. plantarum*, *L. brevis* CD2, *Bifidobacterium* spp, etc. were proposed and used to obtain caries incidence reduction, mutans streptococci and lactobacilli count change, plaque pH control and root caries lesions reversal (21-24). Several appropriate vehicles of administration of probiotic strains have been proposed. However, specifically formulated devices with slow release of the microbial strain might be needed in order to oral diseases prevention and control.

### **MECHANISM OF ACTION OF PROBIOTICS:**

Probiotic bacteria include a broad spectrum of harmless and physiologically distinct strains, primarily belonging to the genera *Lactobacillus* and *Bifidobacterium*. Research into the health-promoting effects of probiotics has mainly focused on alleviating medical conditions, such as infections of the gastrointestinal tract (25). Several mechanisms have been proposed for the potential beneficial effect of probiotics. These can be classified into four broad lines:

1. The production of antimicrobials (bacteriocins) or acids that can inhibit the proliferation of pathogens.
2. Competition for cell adhesion sites (competitive inhibition or replacement therapy) with pathogens and/or co-aggregation to biofilm.
3. Modulation of local and systemic immune functions.
4. Degradation of toxins (26).

All available data show that the effects of probiotics are species- and strain-specific (27-29). There are studies which shown the effects of probiotics on reducing the incidence of cavities, in which the reduction took *Streptococcus mutans* in the oral cavity mainly with the probiotic *L. reuteri* ATCC 55730. The strains of *L. salivarius* BGHO1, *L. fermentum* BGHO36 and BGHO 64, *L. gasseri* and *L. delbrueckii* BGHO89 subsplactis BGHO99 who had antagonistic action on *Ataphylococcus aureus*, *Enterococcus faecalis*, *Micrococcus flavus*, *Salmonella enteritidis*, *Streptococcus pneumoniae* and *Streptococcus mutans*, except on the growth of *Candida albicans* are identified from samples in the tooth surface and healthy oral mucosa. They also found that strains of *L. gasseri* and *L. salivarius* BGHO89 BGHO1 were tolerant to low pH and high concentration of bile salts (30-33).

*Streptococcus salivarius* strain JH is a potential probiotic candidate which produces multiple proteinaceous antimicrobials (bacteriocins), the inhibitory spectrum that includes *Streptococcus mutans*, one of the principle causative agents for the development of dental caries. The genome of strain JH has previously studied and it was found to show to contain

the biosynthetic loci for the bacteriocins salivaricin A3, streptin and streptococcin SA-FF22. Thus, strain JH also produces salivaricin E, a 32 aa lantibiotic with a mass of 3565.9 Da, which is responsible for the inhibition of *S. mutans* growth (34).

According to literature, there are different mechanism of action for different strains of probiotic in the oral cavity. So, with these actions of probiotic strains, the metabolic activity of biofilm in the oral cavity and composition can be altered which leads to the prevention of dental caries. However, there are between 75 and 100 bacterial species in the oral cavity of each patient, with different combinations of species. So, if that's the case, we might require a mixture of beneficial strains rather than a single strain as oral bacteriotherapy (35).

### **PROBIOTIC STRAINS:**

Based on the literature search, there are wide range of strains which include *Streptococcus salivarius*, *L. paracasei*, *L. caseii*, *L. rhamnosus*, *L. acidophilus*, *L. reuteri*, *L. plantarum*, *L. brevis* and *Bifidobacterium lactis*. In vitro studies on monocultures of different probiotic lactobacilli strains have shown a varying acid production, while clinical trials have not revealed an impact on plaque acidogenicity (46,47).

A randomised open label clinical trial has been done to evaluate the effects of the lactic acid bacterium *Lactobacillus salivarius* on caries risk factors and it was concluded that oral consumption of *L. salivarius* WB21 and TI 2711 provided better resistance to caries risk factors compared to the xylitol tablet with no side effects related to advance of caries in the oral cavity (36). A double-blinded, randomized, placebo-controlled study evaluated the efficacy of the probiotic *L. paracasei*GMNL-33 for reducing caries-associated salivary microbial counts in healthy adults. And concluded that 2-weekperiod of medication was needed for *L. paracasei*GMNL-33 via oral administration route to become effective in the probiotic action. And short-term probiotic intervention seemed to enhance its inhibitory effect

against *S. mutans* after the intervention (37). And another double-blinded, randomized, placebo-controlled trial has done to provide a preliminary evaluation of *S. salivarius*M18 for its probiotic application to the prevention, or a reduction in the risk, of dental caries and found that *S. salivarius*M18 has oral health benefits when it is administered regularly (38). *L. salivarius* can secrete intermediates that is capable of inhibiting the formation of cariogenic *S. mutans* and *C. albicans* biofilm; and also inhibits fungal morphological transformation thus reducing the pathogenicity of *C. albicans*; ultimately weakens its pathogenic potential (39). A short-term daily ingestion of lactobacilli-derived probiotics which was a prepared prepared straws or lozenges has been delivered and it reduced the levels of salivary mutans streptococci in young adults (40). yoghurt containing LGG has an inhibitory effect on oral pathogenic bacteria and is beneficial in caries prevention on its daily consumption (41). An invitro study investigated the effect of *L. lactis* HY 449 on cariogenic biofilm and concluded that it possessed inhibitory mechanisms and useful probiotic for the prevention of dental caries (43). A randomized double blinded clinical trial demonstrated that *S. salivarius*K12 over a 28-day period does not adversely affect the human host on daily ingestion and supports the safety of its oral delivery in a food-based carrier (44). It has been proved that levels of microorganisms in saliva which is responsible of development of dental caries has been reduced with probiotic Bifidobacteria in yogurt (42). *L. rhamnosus* has been concluded as a candidate for prevention of dental caries since it inhibits oral biofilm formation by decreasing glucan production of *S. mutans* and has antibacterial activity and did not integrate into oral biofilm (45).

#### **AVAILABLE FORMS OF PROBIOTICS:**

There are various forms of probiotics available that can be used on daily basis. Common probiotic bacteria used in clinical studies targeting oral health are *Lactobacillus* spp. and *Bifidobacterium* spp., in vehicles such as milk, yogurt, cheese, drops, gum, ice

cream, lozenges and tablets (48). The number of *S. mutans* have been reduced by using the milk form Lactobacillus rhamnosus GG strain, Cheese form of Lactobacilli mix and Yoghurt form of Lactobacillus reuteri ATCC 55730, Bifidobacterium animalis DN-173010 (49-52). In addition to these results, Chewing gum, tablet and Lozenge forms of Lactobacillus reuteri ATCC 55730 and ice cream form of Bifidobacterium lactics Bb-12 have shown to decrease the count of *S. mutans* on wide range of age groups (53-55).

Milk, Cereals, Drops forms of Lactobacillus rhamnosus LB21, Lactobacillus paracasei F19, Lactobacillus reuteri DSM 55730 are present and they found to decrease the outcome of dental caries (56-58). Streptococcus uberis KJ, Streptococcus oralis KJ, Streptococcus rattus JH14 are available as Tablet form and has inhibitory effect of dental caries (59). Ideally, the delivery should be suitable for all ages especially young children, since, it has been recommended that early exposure in life would bring in a permanent colonization of health promoting strains (60-62). However, still research goes on the identification of ideal administration vehicle (63).

#### **FUTURE FOR PROBIOTICS IN CARIES PREVENTION:**

The main goal of probiotics use in prevention of caries is to destroy or reduce the bacteria which is cariogenic, mainly mutans streptococci and replace it with noncariogenic bacteria (64). In order to prevent the occurrence of dental caries, probiotic bacteria should adhere to the oral mucosa and dental tissues as part of the biofilm and compete with the growth of dental pathogens (65). Probiotics had showed quite a few promising results in caries prevention, but there are only few clear clinical outcomes pertaining to prevention of caries. So, the proper scientific evidence are still in search. A regular continuous daily intake

of probiotic is probably required through vehicles like toothpastes or mouthrinses. For all anticaries product to be effective in its action, almost daily usage is needed, therefore probiotics should be incorporated in our daily preventive products like toothpaste. In spite of this, probiotics can also be administered in daily dietary supplements, and it is considered effective to some extent in prevention of caries (66).

#### **FURTHER RESEARCH AND EMERGING PROBIOTIC STRAIN:**

Caries management strategies have travelled through various paths and are now witnessing a paradigm shift toward preventive approach. Bacteriotherapy is one of the most potential concepts to prevent dental caries (67). There are previous studies on reduction of mutans streptococci count with the help of available probiotic strains such as *Lactobacillus rhamnosus* GG, *Lactobacillus reuteri*, and *Bifidobacterium*. However, these strains possess certain limitations in their mechanism of action against biofilm and their evidence is still insufficient. In order to overcome those drawbacks of previously available strains, new probiotic strains are under search. One such emerging strain is *Streptococcus thermophilus* and it was found to be an effective probiotic for oral health (68,69) *Streptococcus salivarius*, a Gram-positive bacterium, is now currently being used as an oral probiotic (70). The most commonly used probiotic *S. salivarius* strains are K12 and M18. In addition to the above strains, *S. salivarius* strain JH has also been found to possess a potent anti-MS activity (71). And future research on probiotics for caries prevention should be based on new strains, vehicle of administration and prescription methods of probiotics for patients (66).

#### **REFERENCES**

1. Campus, G.; Solinas, G.; Cagetti, M.G.; Senna, A.; Minelli, L.; Majori, S.; Montagna, M.T.; Reali, D.; Castiglia, P.; Strohmenger, L. National pathfinder survey of 12-year-old children's oral health in Italy. *Caries Res.* 2007, 41, 512–517.
2. Selwitz, R.H.; Ismail, A.I.; Pitts, N.B. Dental caries. *Lancet* 2007, 369, 51–59.
3. Eliasson, L.; Carlén, A.; Almståhl, A.; Wikström, M.; Lingström, P. Dental plaque pH and micro-organisms during hyposalivation. *J. Dent. Res.* 2006, 85, 334–338.
4. Yadav, Khushbu & Prakash, Satyam. (2016). Dental Caries: A Review. 06. 01-07.
5. Cagetti MG, Mastroberardino S, Milia E, Cocco F, Lingström P, Campus G. The use of probiotic strains in caries prevention: a systematic review. *Nutrients.* 2013;5(7):2530–50.
6. Takahashi, N.; Nyvad, B. The role of bacteria in the caries process: Ecological perspectives. *J. Dent. Res.* 2011, 90, 294–303.
7. Duckworth, R.M. The science behind caries prevention. *Int. Dent. J.*, 1993, 43, 529–539.
8. Zander, H.A. and Bibby, B.G. Penicillin and caries activity. *J. Dent. Res.*, 1947, 26, 365–368
9. Anderson, M.H. A review of the efficacy of chlorhexidine on dental caries and the caries infection. *J. Calif. Dent. Assoc.*, 2003, 31, 211–214.
10. Walker, C. Effects of sanguinarine and Sanguinaria extract on the microbiota associated with the oral cavity. *J. Can. Dent. Assoc.*, 1990, 56, 13–30.
11. Edwardsson, S., Birkhed, D. and Mejare, B. Acid production from Lycasin, maltitol, sorbitol and xylitol by oral streptococci and lactobacilli. *Acta. Odontol. Scand.*, 1977, 35, 257–263.
12. Karulf, R.E., Collier, J.A., Bartolo, D.C., et al. Anorectal physiology testing. A survey of availability and use. *Dis. Colon. Rectum.*, 1991, 34, 464–468

13. Reynolds, E.C. Anticariogenic complexes of amorphous calcium phosphate stabilized by casein phosphopeptides: A review. *Spec. Care Dentist.* 1998, 18, 8–16.
14. Petersen, P.E.; Lennon, M.A. Effective use of fluorides for the prevention of dental caries in the 21st century: The WHO approach. *Community Dent. Oral Epidemiol.* 2004, 32, 319–321
15. Zero, D.T. Dentifrices, mouthwashes and remineralization/caries arrestment strategies. *BMC Oral Health* 2006, 6(Suppl. 1), 9.
16. Allaker, R.P. and Douglas, C.W. Novel anti-microbial therapies for dental plaque-related diseases. *Int. J. Antimicrob. Agents.*, 2009, 33, 8–13.
17. World Health Organization. Available online: [http://www.who.int/foodsafety/publications/fs\\_management/en/probiotics.pdf](http://www.who.int/foodsafety/publications/fs_management/en/probiotics.pdf) (accessed on 17 April 2013).
18. Coqueiro AY, Bonvini A, Tirapegui J, Rogero MM. Probiotics supplementation as an alternative method for celiac disease treatment. *International Journal of Probiotics & Prebiotics [Internet]*. 2017;12:23–32.
19. Floch, M.H.; Walker, W.A.; Madsen, K.; Sanders, M.E.; Macfarlane, G.T.; Flint, H.J.; Dieleman, L.A.; Ringel, Y.; Guandalini, S.; Kelly, C.P.; et al. Recommendations for probiotic use—2011 update. *J. Clin. Gastroenterol.* 2011, 4, S168–S171.
20. Singh VP, Sharma J, Babu S, Rizwanulla & Singla A. Role of probiotics in health and disease: a review. *J Pak Med Assoc.* 2013;63(2):253-7
21. Taipale, T.; Pienihäkkinen, K.; Alanen, P.; Jokela, J.; Söderling, E. Administration of *Bifidobacterium animalis* subsp. *lactis* BB-12 in early childhood: A post-trial effect on caries occurrence at four years of age. *Caries Res.* 2013, 47, 364–372.
22. Marttinen, A.; Haukioja, A.; Karjalainen, S.; Nylund, L.; Satokari, R.; Öhman, C.; Holgerson, P.; Twetman, S.; Söderling, E. Short-Term consumption of probiotic

- lactobacilli has no effect on acid production of supragingival plaque. *Clin. Oral Investig.* 2012, 16, 797–8
23. Keller, M.K.; Hasslöf, P.; Dahlén, G.; Stecksén-Blicks, C.; Twetman, S. Probiotic supplements (*Lactobacillus reuteri* DSM 17938 and ATCC PTA 5289) do not affect regrowth of mutans streptococci after full-mouth disinfection with chlorhexidine: A randomized controlled multicenter trial. *Caries Res.* 2012, 46, 140–146
  24. Lexner, M.O.; Blomqvist, S.; Dahlén, G.; Twetman, S. Microbiological profiles in saliva and supragingival plaque from caries-active adolescents before and after a short-term daily intake of milk supplemented with probiotic bacteria—A pilot study. *Oral Health Prev. Dent.* 2010, 8, 383–388.
  25. Ravn, I., Dige, I., Meyer, R.L. and Nyvad, B. Colonization of the oral cavity by probiotic bacteria. *Caries Res.*, 2012, 46, 107-112
  26. Babaji, Prashant. (2012). Role of probiotics in oral health: A review of the literature. *Journal of Education and Ethics in Dentistry*. 2. 10.4103/0974-7761.121256.
  27. Hedayati-Hajikand T, Lundberg U, Eldh C & Twetman S. Effect of probiotic chewing tablets on early childhood caries – a randomized controlled trial. *BMC Oral Health*. 2015 Sep 24;15(1):112
  28. Ashwin D, Ke V, Taranath M, Ramagoni NK, Nara A & Sarpangala M. Effect of Probiotic Containing Ice-cream on Salivary Mutans Streptococci (SMS) Levels in Children of 6-12 Years of Age: A Randomized Controlled Double Blind Study with Six-months Follow Up. *J Clin Diagn Res* 2015;9(2):ZC06-9 doi: 10.7860/JCDR/2015/10942.5532. Epub 2015 Feb 1.
  29. Rebolledo M, Rojas E, Salgado F. Efecto de dos probióticos que contienen cepas de *Lactobacillus casei* variedad *rhamnosus* y *Lactobacillus johnsonii* sobre el crecimiento *in vitro* de *Streptococcus mutans*. *Int. J. Odontostomat.* 2013;7(3):415-419

30. Nozari A, Motamedifar M, Seifi N, Hatamizargaran Z, Ranjbar MA (2015) The Effect of Iranian Customary Used Probiotic Yogurt on the Children's Salivary Cariogenic Microflora. *J Dent (Shiraz)* 16: 81-86.
31. Nishihara T, Suzuki N, Yoneda M, Hirofuji T (2014) Effects of *Lactobacillus salivarius*-containing tablets on caries risk factors: a randomized open-label clinical trial. *BMC Oral Health* 14: 110.
32. Cortés-Dorantes N, Ruiz-Rodríguez MS, Karakowsky-Kleiman L, Garrocho-Rangel JA, Sánchez-Vargas LO, et al. (2015) Probiotics and their effect on oral bacteria count in children: a pilot study. *Eur J Paediatr Dent* 16: 56-60.
33. Astekar M, Sidhu GK, Kathuria NS (2014) Impact of diet alteration on oral microflora by addition of probiotics. *Indian J Med Microbiol* 32: 466-467.
34. Glenn V. Walker, Nicholas C.K. Heng, Alan Carne, John R. Tagg1 and Philip A. Wescombe Salivaricin E and abundant dextranase activity may contribute to the anti-cariogenic potential of the probiotic candidate *Streptococcus salivarius*. *JH. Microbiol.*, 2016, 162, 476–486.
35. Twetman. Are we ready for caries prevention through bacteriotherapy? *Braz Oral Res.* 2012;26(SpecIss 1):64-70
36. Tetsuyo Nishihara, Nao Suzuki, Masahiro Yoneda and Takao Hirofuji. Effects of *Lactobacillus salivarius*-containing tablets on caries risk factors: A randomized open-label clinical trial. *BMC Oral Health.*, 2014, 14, 110
37. Li-Chuan Chuan, Chiung-Shing Huang, Li-Wei Ou-Yang, Shiao-Yu Lin Probiotic *Lactobacillus paracasei* effect on cariogenic bacterial flora. *Clin. Oral. Investig.*, 2011, 15, 471-476.
38. Jeremy P. Burton, Bernadette K. Drummond, Chris N. Chilcott, John R. Tagg, W. Murray Thomson et al. Influence of the probiotic *Streptococcus salivarius* strain M18

- on indices of dental health in children: a randomized double-blind, placebo-controlled trial. *J. Med. Microbiol.*, 2013, 62, 875-884.
39. Wirginia Krzyściak, Dorota Kościelniak, Monika Papież, Palina Vyhouskaya, et al. Effect of a *Lactobacillus salivarius* Probiotic on a Double-Species *Streptococcus mutans* and *Candida albicans* caries Biofilm. *Nutr.*, 2017, 9, 1242.
40. Esber Caglar, Sule Kavaloglu, Semra Ergeneli, Nuket Sandalli et al. Salivary mutans streptococci and lactobacilli levels after ingestion of the probiotic bacterium *Lactobacillus reuteri* ATCC 55730 by straws or tablets. *Acta Odontol. Scand.*, 2006, 64, 314-318
41. Domagoj Glavina, Kristina Gorseta, Ilija Skrinjari, Dubravka Negovetic Vranic et al. Effect of LGG Yoghurt on *Streptococcus Mutans* and *Lactobacillus Spp.* Salivary Counts in Children. *Coll. Antropol.*, 2012, 36, 129-132
42. Esber Caglar, Nuket Sandalli, Svante Twetman, Sule Kavaloglu et al. Effect of yogurt with *Bifidobacterium DN-173 010* on salivary mutans streptococci and lactobacilli in young adults. *Acta Odontol. Scand.*, 2005, 63, 317-320.
43. Young-Jae Kim, Sung-Hoon Lee. Inhibitory Effect of *Lactococcus lactis* HY 449 on Cariogenic Biofilm. 2016.
44. Burton, J.P., Cowley, S., Simon, R.R., McKinney, J. et al. Evaluation of safety and human tolerance of the oral probiotic *Streptococcus salivarius* K12: A randomized, placebo-controlled, double-blind study. *Fd. Chem. Toxicol.*, 1, 2011, 49, 2356-23
45. Sung Hoon Lee, Young Jae Kim. A comparative study of the effect of probiotics on cariogenic biofilm model for preventing dental caries. 2014, 196, 8.
46. Hedberg M, Hasslof P, Sjöström I, Twetman S, Stecksén-Blicks C: Sugar fermentation in probiotic bacteria—an in vitro study. *Oral Microbiol Immunol* 2008;23: 482–485.

47. Haukioja A, Soderling E, Tenovuo J: Acid production from sugars and sugar alcohols by probiotic lactobacilli and bifidobacteria in vitro. *Caries Res* 2008; 42:449–453
48. Hasslöf, Pamela &Stecksén-Blicks, Christina. (2020). Chapter 10: Probiotic Bacteria and Dental Caries. 10.1159/000455377.
49. Nase L, Hatakka K, Savilahti E, Saxelin M, Pönkä A, Poussa T, Korpela R, Meurman JH: Effect of long-term consumption of a probiotic bacterium, *Lactobacillus rhamnosus* GG, in milk on dental caries and caries risk in children. *Caries Res* 2001;35:412–420.
50. Ahola AJ, Yli-Knuuttila H, Suomalainen T, Poussa T, Ahlström A, Meurman JH, Korpela R: Short-term consumption of probiotic-containing cheese and its effect on dental caries risk factors. *Arch Oral Biol* 2002;47:799–804.
51. Nikawa H, Makihira S, Fukushima H, Nishimura H, Ozaki Y, Ishida K, Darmawan S, Hamada T, Hara K, Matsumoto A, Takemoto T, Aimi R: *Lactobacillus reuteri* in bovine milk fermented decreases the oral carriage of mutans streptococci. *Int J Food Microbiol* 2004; 95:219–223.
52. Caglar E, Sandalli N, Twetman S, Kavaloglu S, Ergeneli S, Selvi S: Effect of yogurt with *Bifidobacterium* DN-173 010 on salivary mutans streptococci and lactobacilli in young adults. *Acta Odontol Scand* 2005;63:317–320.
53. Caglar E, Kuscu OO, Cildir SK, Kuvvetli SS, Sandalli N: A probiotic lozenge administered medical device and its effect on salivary mutans streptococci and lactobacilli. *Int J Paediatr Dent* 2008;18: 35–39.
54. Caglar E, Kavaloglu SC, Kuscu OO, Sandalli N, Holgerson PL, Twetman S: Effect of chewing gums containing xylitol or probiotic bacteria on salivary mutans streptococci and lactobacilli. *Clin Oral Investig* 2007;11:425–429.

55. Caglar E, Kuscu OO, Selvi Kuvvetli S, KavalogluCildir S, Sandalli N, Twetman S: Short-term effect of ice-cream containing *Bifidobacterium lactis* Bb-12 on the number of salivary mutans streptococci and lactobacilli. *ActaOdontolScand* 2008;66:154–158.
56. Petersson LG, Magnusson K, Hakestam U, Baigi A, Twetman S: Reversal of primary root caries lesions after daily intake of milk supplemented with fluoride and probiotic lactobacilli in older adults. *ActaOdontScand* 2011;69:321–327.
57. Hasslof P, West CE, Videhult FK, Brandelius C, Stecksén-Blicks C: Early intervention with probiotic *Lactobacillus paracasei* F19 has no long-term effect on caries experience. *Caries Res* 2013;47: 559–565
58. Stensson M, Koch G, Coric S, Abrahamsson TR, Jenmalm MC, Birkhed D, Wendt LK: Oral administration of *Lactobacillus reuteri* during the first year of life reduces caries prevalence in the primary dentition at 9 years of age. *Caries Res* 2014;48:111–117
59. Hedayati-Hajikand T, Lundberg U, Eldh C, Twetman S: Effect of probiotic chewing tablets on early childhood caries – a randomized controlled trial. *BMC Oral Health* 2015;15:112
60. Meurman, J.H. Probiotics: Do they have a role in oral medicine and dentistry? *Eur. J. Oral. Sci.*, 2005, 113, 188-196.
61. Yli-Knuutila, H., Snäll, J., Kari, K. and Meurman, J.H. Colonization of *Lactobacillus rhamnosus* G in the oral cavity. *Oral. Microbiol. Immunol.*, 2006, 21, 129-131.
62. Twetman, S. and Stecksén-Blicks, C. Probiotics and oral health effects in children. *Int. J. Paediat. Dent.*, 2008, 18, 3-10.
63. Meurman, J.H. and Stamatova, I. Probiotics: Contributions to oral health. *Oral. Dis.* 2007, 13, 443.

64. Twetman, S.; Keller, M.K. Probiotics for caries prevention and control. *Adv. Dent. Res.* 2012, 24, 98–102.
65. Van Loo J, Jonkers N. Evaluation in human volunteers of the potential anticarcinogenic activities of novel nutritional concepts: Prebiotics, probiotics and synbiotics (the SYNCAN project QLK 1-1999-00346). *NutrMetabCardiovasc Dis* 2001;11:87-93.
66. Maria Grazia Cagetti, Stefano Mastroberardino, Egle Milia, Fabio Cocco, Peter Lingström and Guglielmo Campus. The use of probiotic strains in caries prevention: A systematic review. *Nutr.*, 2013, 5, 2530-2550
67. Poorni S, Srinivasan MR, Nivedhitha MS. Probiotic *Streptococcus* strains in caries prevention: A systematic review. *J Conserv Dent* 2019;22:123-8
68. Cotter PD, Hill C. Surviving the acid test: Responses of gram-positive bacteria to low pH. *Microbiol Mol Biol Rev* 2003;67:429-53
69. Arioli S, Ragg E, Scaglioni L, Fessas D, Signorelli M, Karp M, *et al.* Alkalinizing reactions streamline cellular metabolism in acidogenic microorganisms. *PLoS One* 2010;5:e15520
70. Di Pierro F, Zanvit A, Nobili P, Risso P, Fornaini C. Cariogram outcome after 90 days of oral treatment with *Streptococcus salivarius* M18 in children at high risk for dental caries: Results of a randomized, controlled study. *ClinCosmetInvestig Dent* 2015;7:107-13
71. Walker GV, Heng NC, Carne A, Tagg JR, Wescombe PA. Salivaricin E and abundant dextranase activity may contribute to the anti-cariogenic potential of the probiotic candidate *Streptococcus salivarius* JH. *Microbiology* 2016;162:476-86