

Original Research Article

BASAL CELL CARCINOMA: IN TERTIARY CARE CENTER, DEMOGRAPHIC DETAILS AND HISTOLOGICAL SUBTYPES AT PUMHS-W NAWABSHAH, PAKISTAN.

Abstract:

Background: Skin malignancies are more common in fair skinned persons, the Caucasians, but its increasing incidence is alarming in Asian population.

Objective: To evaluate the Basal Cell Carcinoma (BCC) in a tertiary care center, in terms of frequency, age, sex, sites, and histological subtypes.

Methods: A retrospective work conducted in the Dermatology and Pathology departments, Peoples University of Medical & Health Sciences, Nawabshah during the month of November 2020 to November 2021, on 100 cases collected by nonprobability consecutive technique. All the clinical demographic data was collected in the Dermatology department and the biopsy sample was processed in Pathology department. The observations were recorded on a structured proforma, statistically analyzed on SPSS 20 and the results were tabulated.

Results: A total of 100 cases were assessed, the age of patients ranged from 22-86 years with a female preponderance. Most of the cases were detected in 61-80 years age group. Cheek was the most common site. On histological examination the nodular/solid variety was diagnosed in majority (54%) of cases.

Conclusion: All the histological subtypes in relation with both genders the density of BCC was relatively higher over the face, followed by the trunk. The nodular/solid variety is more common in post-menopause and elderly women.

Key Words: Caucasians, BCC, Dermatology, nodular/solid variety, cheeks, post –menopause.

Introduction:

Skin malignancies are classified as keratinocyte and melanocytic neoplasms. Basal cell carcinoma is categorized as keratinocytic or non-melanocytic skin cancer¹. Basal cell carcinoma (BCC) the most common skin malignancy worldwide with constantly increasing incidence due to an aging population and a predilection towards sun exposed areas especially head and neck areas^{2,3}. It is a slow growing tumor with propensity for local invasion with morbid destruction, however metastasis is seldom seen.⁴ Accounting to approximately 70% of all malignant diseases of the skin in the western countries⁵. BCCs are more common among the elderly male population with a predilection for head and neck region⁶. The higher incidence in men is probably due to increased recreational and occupational exposure to the sun, and also as the women are more concerned about their health and aesthetics, and go to the dermatologist earlier when they have lesions especially on face and neck areas.⁷ Extensive local tissue destruction is common and if not adequately managed, however metastasis is exceptionally rare⁸. Clinically the BCC is divided into several subtypes, but the main subtypes are nodular, superficial, and morpheaform BCC⁹. The histological pattern of BCC is the predictors of clinical behavior,⁶ for that purpose a skin punch or excisional biopsy is necessary for the confirmation of various histologic types of BCC¹⁰, including solid (nodular), micronodular, infiltrating, superficial,

fibroepithelial, basosquamous (metatypical), keratotic, basal cell carcinoma with adnexal differentiation, and other variants.^{5,11}

There is negligible mortality, but may be associated with significant morbidity if untreated for a long period of time.¹² The differential diagnosis of BCC includes adnexal tumors with follicular, sweat gland, or sebaceous differentiation and certain types of Squamous Cell Carcinoma (SCC). Nodular type of BCC may be confused with trichoblastoma or trichoepithelioma. Superficial BCC may mimic some inflammatory dermatoses such as psoriasis and eczema. Morphea-like BCC may be confused with a plaque of Morphea or a scar. In these cases, histopathological helps is essential to establish the diagnosis of BCC.¹³

Keeping the above mentioned facts in view, we conduct this study to see the morphological spectrum and frequency of BCC cases in a tertiary care hospital of rural area in our setup.

Methodology:

This retrospective work was conducted in the Dermatology and Pathology departments, Peoples University of Medical & Health Sciences, Nawabshah during the month of November 2020- November 2021. We evaluate the 100 cases of BCCs. All the cases were assessed in Dermatology department and the biopsy sample was processed in Pathology department. The cases were collected by nonprobability consecutive technique. A structured proforma was designed for the study and all the available data was recorded on that. The histology was evaluated by retrieving the slides and blocks of the study population, the histological diagnosis was confirmed and the tumors were classified according to the World Health Organization (WHO) classification.¹¹ The findings were analyzed statistically and the results were tabulated.

Results:

We analyzed 100 cases of BCC, the age of patients varies from 22 to 86 years with a mean age of 66.7 years. The maximum number of cases were observed in 61-80 years age group. The female (56%) patients were slightly more in number than male (44%) cases, with a male to female ratio of 1:1.3 (table-1 & Fig-1).

Table-1. Age Groups with Gender Distribution and Number of Cases (n=100)

S. No	Age Group	Number of Cases		Total
		Male	Female	
1	21-40	01	03	04
2	41-60	17	16	33
3	61-80	21	32	53
4	81-100	05	05	10
	Total	44	56	100




Fig-1. Age Groups in Study Population (n=100)

The most common site we found was cheek.

Regarding different sites of the tumor, we observed 52 (52%) cases on cheek which was the most common site, followed by 27 (27%) lesions on nose, the sternum was found the least common site having only one lesion (table-2).

Table-2. Frequency of Site of the Lesion (n=100)

S.No	Site	No of Cases
01	Cheek	52
02	Nose	27
03	Forehead	07
04	Trunk	05
05	Upper eye lid	04
06	Shoulders	02
07	Scalp	02
08	Sternum	01

On histological examination the nodular/solid tumors were detected in majority (54%) of cases, followed by infiltrating (12%), basosquamous/metatypical (7%) (table-3).

Table-3. Histological Subtypes with Number of Cases.

S. No	Histological Subtypes	No of Cases
01	Nodular/solid	54
02	Infiltrating	12
03	Basosquamous/metatypical	07
04	Superficial	04
05	Keratotic	03
06	Micronodular	03
07	BCC with adnexal differentiation	01
08	Other variants (Pigmented, adenoid, infundibulocystic)	15
09	Mixed (solid and adenoid)	01
	Total	100

Discussion:

Cancer or malignancy is commonly occurring at old age, Age with sex both are major and non-modifiable risk factors for cancer, more frequently involving male than female.^{14,15,16,17}

Skin cancers are more common tumors around the globe with more frequently involving the Caucasians than Asians.¹⁸

Our present study results showed the age of patients varies from 22 to 86 years with slight earlier occurrence of BCC in young age group, same results were recorded at Faisalabad by Janjua¹⁹, slight difference were noticed in study in which 44yrs – 89 yrs age group was noticed done at Turkey by YO¹⁸, but the results are inconsistent with the Rana¹ in Turkey where the BCC is more common in 6th and 7th decade of life.

The mean age in our study was 66.7 years, 66.9years Oh Singapore²⁰, 65yrs Y Scrivener²¹, 64.5 yrs Turkey YO¹⁸, 64.5 yrs Metib,²² 61 yrs by Rishu,²³ 59 yrs Kikuchi A²⁴ Japan, 54.5 yrs Dowerah²⁵ India.

The maximum numbers of cases were observed in 61-80 years age group. Same results were seen at Singapore by Oh²⁰ where the risk of skin cancer increased after 70yrs, Bassukas⁷ Turkey. Asians developed their first BCC at an older age than the age of first BCC in Caucasian.²⁶

The female preponderance were noticed than affected male, same results were seen

In study by Rishu²³ 82%, Kikuchi A²⁴, Japan, those were not matched with our results were noticed in Moore²⁶ found equal incidence in both sexes in Asian countries, male predominance were seen in Jordan by Al-Qarqaz²⁷, Turkey by YO¹⁸, Pakistan 1.4:1¹⁹, and 2:1 by Dowerah²⁵,

Male to female ratio in our study was 1:1.3. 0.97 kikuchi.A²⁴ Japan, 0.92 Scrivener²¹, 9;10 Turkey YO¹⁸, 1.4:1 Janjua¹⁹ Pakistan, 2:1 Dowerah²⁵ India, 1.76:1 Jordan, Al-Qarqaz.²⁷

Regarding different sites of the tumor, we observed cheek was the most common site involved, followed by nose, our results are consistent with the results of Rana¹, Turkey, but vice versa results were seen by Janjua¹⁹ at Faisalabad, Pakistan, and Jordan by Al-Qarqaz²⁷, both these anatomical sites are included in high risk areas. The sternum was found the least common site we found, limbs are the least common site involved by BCC at Jordan by Al-Qarqaz²⁷.

On histological examination the Nodular/Solid tumors were detected in majority of cases, same results were seen at Japan Kikuchi²⁴, Turkey, YO¹⁸, Pakistan, Janjua,¹⁹ India Dowerah²⁵, Scrivener²¹. The mixed type was least frequent type observed in our study opposite results were observed at Pakistan Janjua¹⁹ where pigmented variety was on second number.

Conclusion: A rising incidence of malignancy was noticed around the globe. Basal cell carcinoma was more prevalent in fair skinned population, but present study suggests that it is not a disease restricted to a geographic/racial group of peoples but instead it can involve any one. The tumor is more common in 61 – 80 years age group, the cheek was the most common site and the nodular solid variety was the most frequent lesion in our study.

Recommendations: Author recommends the establishment of central cancer statistics in country.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is

absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

References:

1. Kapukaya R, Çiloğlu O. Evaluation of the Demographic and Clinical Features of Skin Cancers: A Single-Center Experience. *Eur J Ther.* 2020 Sep 1;26(3):214-7.
2. Sreekantaswamy S, Endo J, Chen A., Butler D., Morrison L., Linos E. Aging and the treatment of basal cell carcinoma. *Clin. Dermatol.* 2019;37:373–378. doi: 10.1016/j.clindermatol.2019.06.004.
3. Rana S, Walvir NM, Khetrpal S, Jetley S, Jairajpuri ZS. Study of Basal Cell Carcinoma and Its Morphological Spectrum. *Sch J App Med Sci*2019; 10.36347/SJAMS.2019.v07i11.048
4. Fagan J, Brooks J, Ramsey ML. Basal Cell Cancer. In: *StatPearls* [Internet] 2021 Mar 1. StatPearls Publishing
5. Saldanha P, Shanthala PR, Upadhaya K. Cutaneous basal cell carcinoma: A morphological spectrum. *Archives of Medicine and Health Sciences.* 2015 Jan 1;3(1):24-28.
6. Sreeram S, Lobo FD, Naik R, Khadilkar UN, Kini H, Kini UA. Morphological spectrum of basal cell carcinoma in Southern Karnataka. *J clinic diagnostic research: JCDR.* 2016 Jun;10(6):EC04-EC07.
7. Bassukas ID, Tatsioni A. Male Sex is an Inherent Risk Factor for Basal Cell Carcinoma. *J. Skin Cancer.* 2019;2019:8304271. doi: 10.1155/2019/8304271.
8. Crowson AN. Basal cell carcinoma: biology, morphology and clinical implications. *Modern pathology.* 2006 Jan 17;19(S2):S127-147.
9. Kim DP, Kus KJB, Ruiz E. Basal Cell Carcinoma Review. *Hematol./Oncol. Clin. N. Am.* 2019;33:13–24. doi: 10.1016/j.hoc.2018.09.004.
10. Weber P, Tschandl P, Sinz C, Kittler H. Dermatoscopy of Neoplastic Skin Lesions: Recent Advances, Updates, and Revisions. *Curr Treat Options Oncol.* 2018 Sep 20;19(11):56.
11. LeBoit PE, Burg G, Weedon D, Sarasain A, editors. *Pathology and Genetics of Skin Tumors.* Lyon: IARC Press; 2006. World Health Organization Classification of Tumors; pp. 13–19.
12. Miller DL, Weinstock MA. Nonmelanoma skin cancer in the United States: Incidence. *J. Am. Acad. Dermatol.* 1994;30:774–778. doi: 10.1016/S0190-9622(08)81509-5.
13. McDaniel B, Badri T, Steele RB. Basal Cell Carcinoma. [Updated 2021 Sep 20]. In: *StatPearls* [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK482439/>
14. J. Campisi, “Aging, cellular senescence, and cancer,” *Annual Review of Physiology*, vol. 75, no. 1, pp. 685–705, 2013.
15. U.S. Institute of Medicine, “Exploring the biological contributions to human health: does sex matter?,” *Journal of Women's Health & Gender-Based Medicine*, vol. 10, no. 5, pp. 433–439, 2001.

16. M. B. Cook, S. M. Dawsey, N. D. Freedman et al., "Sex disparities in cancer incidence by period and age," *Cancer Epidemiology Biomarkers & Prevention*, vol. 18, no. 4, pp. 1174–1182, 2009.
17. R. J. Biggar, A. W. Bergen, and G. N. Poulsen, "Impact of x chromosome genes in explaining the excess risk of cancer in males," *American Journal of Epidemiology*, vol. 170, no. 1, pp. 65–71, 2009.
18. TIFTIKCIOĞLU YÖ, Karaaslan O, Aksoy HM, Aksoy B, Koçer U. Basal cell carcinoma in Turkey. *The Journal of Dermatology*. 2006 Feb;33(2):91-5.
19. Janjua OS, Qureshi SM. Basal cell carcinoma of the head and neck region: an analysis of 171 cases. *Journal of skin cancer*. 2012 Dec 19;2012.
20. Oh CC, Jin A, Koh WP. Trends of cutaneous basal cell carcinoma, squamous cell carcinoma, and melanoma among the Chinese, Malays, and Indians in Singapore from 1968-2016. *JAAD international*. 2021 Sep 1;4:39-45.
21. Scrivener Y, Grosshans E, Cribier B. Variations of basal cell carcinomas according to gender, age, location and histopathological subtype. *British Journal of Dermatology*. 2002 Jul;147(1):41-7.
22. Metib NJ, Aboud AH. Histopathologic study for skin basal cell carcinoma. *Karbala J Med* 2018;11:3884-8.
23. Rishu gupta..6. Gupta R, Gordon SL, Council ML, Hurst EA. Clinical characteristics of basal cell carcinoma in African Americans: a 10-year retrospective review at a single Academic institution. *Dermatologic Surgery*. 2019 May 1;45(5):660-5.
24. Kikuchi A, Shimizu H, Nishikawa T. Clinical and histopathological characteristics of basal cell carcinoma in Japanese patients. *Archives of dermatology*. 1996 Mar 1;132(3):320-4.
25. Dowerah S, Naiding M. *International Journal of Research in Health Sciences*.
26. Moore MG, Bennett RG. Basal cell carcinoma in Asians: a retrospective analysis of ten patients. *Journal of Skin Cancer*. 2012 Jul 5;2012.
27. Al-Qarqaz F, Marji M, Bodoor K, Almomani R, Al Gargaz W, Alshiyab D, Muhaidat J, Alqudah M. Clinical and demographic features of basal cell carcinoma in north Jordan. *Journal of skin cancer*. 2018 Oct 4;2018.