

Original Research Article

Effectiveness of Standardized Nursing Care Protocol Post Cardiac Catheterization to Reduce Hematoma Development

Abstract

Background

Hematoma development is the most common vascular access complication among post-cardiac catheterization patients. It requires safe and standardized nursing care protocol, as well as qualified and skilled health providers to obtain good outcomes of management. The vascular complications were severe and increased the length of stay in the hospital. Timely recognition and management can diminish their burden.

Objectives

1. To evaluate the effectiveness of Standardized Nursing Care Protocol Post cardiac catheterization to reduce hematoma development.
2. To assess the association of hematoma development between procedure site (radial and femoral), comorbidities (diabetes, hypertension, and obesity) among post-cardiac catheterized patients.

Methodology

This quasi-experimental (post) study was conducted at Coronary Care Units at tertiary care hospitals in Karachi. In this study, a total of 108 post-cardiac catheterized patients participated. The patients were divided into two groups; the interventional group and the control group. In the control group, 54 post-cardiac catheterized patients were selected and

data were taken based on inclusion criteria who received routine holistic nursing care post-procedure and have been discharged. Besides 54 patients were participated in the interventional group who received Standardized Nursing Care Protocol post-cardiac catheterization including for proper patient assessment, compression techniques, interventions pre, during and post sheath and TR-band removal to discharge instruction by the principal investigator (PI). The data was collected through patients' assessment sheets and hematoma scales for the identification of hematoma development rate among post-cardiac catheterized patients. The non-probability consecutive sampling technique was accomplished for the collection of data.

Results

The rate of hematoma development in the (control) group was reflected 18 (33.3 %) while after implementation of the standardized nursing care protocol the rates of hematoma development was 9 (16.6%) in the (interventional) group out of 54 (50%) post-cardiac catheterized patients. There was a significant difference between the (control) and (interventional) groups regarding the rates of hematoma development post-cardiac catheterization i.e. the P values were $0.037 < 0.05$. Thus, the standardized nursing care protocol post-cardiac catheterization was effective for the reduction of hematoma development.

Furthermore, the radial site procedure reflected fewer rates of hematoma development 8 (14.81%) as compared to the femoral site procedure 19 (35.1%) among post-cardiac catheterized out of 54 (50%). The P-value was $0.058 > 0.05$ there was no association between hematoma development and procedure site in the (interventional and control) groups post-cardiac catheterization. Similarly, the P-value was greater than 0.05 hence there was no association

between hematoma development and comorbidities (diabetes, hypertension, and obesity) post-cardiac catheterization.

Conclusion

The implementation of standardized nursing care protocol post-cardiac catheterization was effective for the reduction of hematoma development in the (intervention) group while there was an association of hematoma development between the (control and interventional) groups. However, there was no association between hematoma development and site procedure (radial and femoral), comorbidities (diabetes, hypertension, and obesity) post-cardiac catheterization.

Keywords: Femoral arterial sheath and TR-band removal, nurses' knowledge and practice, standardized nursing care protocol, Cardiac Catheterization, and hematoma.

Introduction

Cardiac catheterization, including coronary angiography and angioplasty, is a standard diagnostic and therapeutic strategy for the evaluation of cardiovascular diseases¹. Coronary angiography is performed by cardiologists for evaluation of the extent of coronary artery disease or blockage². Even in the United States, more than one million patients undergo percutaneous coronary intervention (PCI) each year³. It can be performed via radial, ulnar, femoral, or brachial arteries through sheath⁴. The sheath is a small flexible catheter, introduced as a guide for wires, stents, and balloons during diagnostic and (PCI's)⁵. Similarly, the Trans radial (TR)-band is a compression device that maintains hemostasis of the radial artery after a Tran's radial procedures⁶. Despite being a popular procedure, vascular puncture access complications are experienced by MI (Myocardial infarction) patients after removals of arterial sheath and TR-band⁷. The most common complications are grade II or III hematoma⁸. It occurs when blood gets

trapped underneath the skin around the insertion site due to improper digital compression which requires 18-20 minutes⁹. Post-cardiac catheterization patients are at high risk of vascular site complications¹⁰. The prevalence of hematoma development ranges from small hematoma 30.6% to medium hematoma 37.6% and large hematoma 16.5% with the major bleeding occurrence at the rate of 1.7% post-cardiac catheterization^{11, 12}. Similarly, as per statistics, there is an association of hematoma development between procedure sites (radial, femoral) and comorbidities (diabetes, hypertension, and obesity) post-cardiac catheterization^{13,14}. As a result, patients face additional difficulties such as extended hospital stay and financial constraint¹⁵. These complications are best prevented by the development and implementation of standardized protocols for removal of femoral sheath and TR-band which are based on scientific evidence¹⁶. The protocols should be based on patients' assessment, compression techniques, interventions pre, during and post sheath and TR-band removal to discharge instruction¹⁷. The Manual compression method of sheath removal is considered the golden standard technique than vascular closure devices which obligate higher vascular site complications¹⁸. The nursing actions ensure an excess of blood return around the sheath¹⁹ especially severe vascular complications can increase illness and death 32% rate as well as expenses of the patients^{20, 21}. Recognition, as well as appropriate management, is dependent on the nurse's competence. Hence standardized nursing care protocols should be established to run nursing training where they are following certain strategies to be utilized^{22, 23}. The lack of universally recognized scientific guidelines or protocol may increase the complication rates if nurses are unaware of how to detect and respond to a hematoma, are unaware of the cause that placing patients at greater risk of complications²⁴. As per statistics, there is an inadequate 40% in the nurse's knowledge, and 87.1% of nurses were observed doing unsatisfactory practices which are related to post-cardiac catheterization at

tertiary care hospitals in Pakistan and Egypt^{25,26}. The nurses should be experts in the monitoring of warning signs, proper assessment, documentation, and proper evaluation of puncture site after removal of the femoral sheath and TR band²⁷. The post-cardiac catheterized patients are at a high risk of vascular site complications²⁸. In the United States of America (USA) the rate of large hematoma was 3% and bleeding was 1.7% post-cardiac catheterization^{29,30}. Additionally, in Jordan (38%) developed vascular complications out of 114 patients³¹. Comparatively in India forearm hematoma developed in 187 (10.7%) patients. Grade I hematoma was most common (3.53%), Grade II (3.08%), Grade III (2.83%) and Grade IV (1.25%) hematoma³².

Digital pressure is the best method to achieve hemostasis when a sheath is removed from the access site which requires specialized nursing care, training, and competency. As per statistics, the Trans radial access associated local hematoma rate is 1.2% to 2.6%, which is 60% lower than in the transfemoral puncture site^{33,34}. Tran's radial artery access is associated with a lower hematoma rate than Tran's femoral artery access³⁵. In USA Tran's radial approach, the rate of hematoma development was (2.9%) and in Tran's femoral approach (7.7%)³⁶. Similarly, in India forearm hematoma was developed in 187 (10.7%) patients³⁷. In elderly and hypertensive patients, there is a greater risk of hematoma or bleeding, which requires closer monitoring of the access site during and after sheath removal³⁸. As per evidence, the rate of hematoma development between comorbidities (hypertension, diabetes and obesity) are the factors of the vascular site complications. High blood pressure may increase the length of time to achieve hemostasis, antihypertensive is required before sheath removal^{39,40}. High blood pressure, 180 mmHg before sheath removal more likely to develop femoral complications⁴¹. The rate of hematoma development among hypertensive patients was 65 (73%) out of 118 after sheath removal post coronary interventions⁴². Similarly, diabetes is one of the factors which increase

the risk of access site complications among post-cardiac catheterization^{43, 44}. Additionally, it is revealed that diabetes was the significant variable of hematoma development 44 (49%) out of 118 patients⁴⁵. Moreover, obesity has been identified as a correlated factor with hematoma development post coronary intervention⁴⁶. Comparatively, the study revealed that obesity is associated with an increased risk of vascular complications post-cardiac catheterization^{47, 48}. Procedure protocols and staff education regarding complications not only ensure the patients' safety and comfort but also reduce the costs. At the same time, it also improves the effectiveness of cardiac catheterization⁴⁹.

Thus, it is recommended that the development and implementation of standardized protocols of arterial sheath and TR-Band can reduce the rate of hematoma development⁵⁰. Therefore, this study was conducted to evaluate the effectiveness of standardized nursing care protocol post-cardiac catheterization in reducing hematoma development and to identify the association of hematoma development between site procedure (radial and femoral) and comorbidities (diabetes, hypertension, and obesity) among post-cardiac catheterized patients.

Methodology

The study setting was conducted at cardiac units (Cath-lab, CCU, Cardiac Ward) at tertiary care hospital in Karachi, Pakistan. The study's target population was post coronary angiography and angioplasty male and female patients who were falling in the inclusion criteria were selected.

Quasi-experimental (post) design was employed to determine the effectiveness of the Standardized Nursing Care Protocol for the reduction of hematoma development and to evaluate the association of hematoma development between procedure site (radial and femoral), comorbidities (diabetes, hypertension, and obesity) among post cardiac catheterization. The whole study period was accomplished from March 2020 to July 2021. The sample size was

calculated by using OpenEpi software version 3 based on statistics. The rate of no hematoma post-cardiac catheterization patients was taken (Control group) 13(15.3%) and post-implementation of standardized nursing care protocol (Intervention group) 35(41.5%)⁵². It was calculated by taking a 95% confidence interval and 80% of the power. The total sample size was 108, 54 in each group. The non-probability consecutive sampling technique was used for the collection of data.

Sample Selection

Inclusion Criteria

- Post coronary angiography of both radial and femoral punctures site patients.
- Post coronary angioplasty only for radial puncture site patients.
- Both male and female patients between 18 years and above will be able to join and consent to take part in the study.
- Normal Prothrombin Time (PT) and Partial Thromboplastin Time (PTT) test.
- Single access puncture of the artery will be included.

Exclusion Criteria

- Post coronary angioplasty femoral puncture site patients were excluded due to the non-availability of the ACT machine.
- Patients with previous surgery of iliac, femoral, and radial arteries.
- Patients with peripheral vascular diseases.
- Hemodynamically unstable patients with bleeding disorders.
- Patients on thrombolytic therapy within 24 hours.

Pilot Study

The pilot study was conducted in March 2020 and April 2020 on 10% of the sample size on post-cardiac catheterization patients. The reliability of the tool according to Cronbach alpha value was ($r=0.722$) including patients' assessment sheet and hematoma scale post removal of TR band and femoral sheath.

Data Collection Procedure

The data were collected by utilizing a patient assessment sheet and hematoma scale post removal of the femoral, radial arterial sheath, and TR-band among post-cardiac catheterized patients. The data was collected among 108 post-cardiac catheterized patients 54 in each group. In this study, a total of 108 post-cardiac catheterized patients participated. The patients were divided into two groups; the interventional group and the control group. In the control group, 54 post-cardiac catheterized patients were selected and data were taken based on inclusion criteria who received routine holistic nursing care post-procedure and have been discharged. Besides 54 patients were participated in the interventional group who received Standardized Nursing Care Protocol post-cardiac catheterization including proper patient assessment, compression techniques, interventions pre, during and post sheath and TR-band removal to discharge instruction by the principal investigator (PI). She was experienced and well trained under interventional cardiologists and certified for sheath and Trans radial band removal. The development of hematoma was measured through a 12-hour follow-up post-procedure after that time has gone undetected among the interventional group. Then, the rate of hematoma development was assessed after the implementation of standardized nursing care protocol and a comparison was made between the interventional group and control group.

Ethical Consideration

The study was approved by the Clinical Research Committee (CRC), Ethics Research Committee (ERC), and Board of Advanced Study Research (BASR) of the Ziauddin University. Furthermore, permission from the hospital was obtained. To accomplish the participant's right of autonomy, informed consent was taken the background, purpose, possible risks of the study, and their right to refuse participation were discussed with the participants.

Result

In this study total of 108 male and female post-cardiac catheterized patients were selected. The patients were divided into two groups (control) group and an (interventional) group. The (control) group contained 54 patients and 54 in the (intervention) group.


The number of male patients was higher (62%) as compared to the number of females (38%) as shown in graph 2. Similarly, the highest number of participants 47 were falling in the 61-80 years age group, and the lowest number of participants 6 were falling 81-100 years as shown in graph 1. Moreover, there were (88%) of angiography and (12%) of angioplasty patients participated. Besides, the highest number of the patients did not receive the injection heparin according to different units, and the lowest number of participants have received heparin doses out of 108 participants. Further, standard 6F sheath was used in all patients and the normal values of PTT and APTT patients were selected.

Furthermore (54) radial puncture site procedures and (54) femoral site punctures were nominated from the study. The radial site procedure patients had fewer rates of hematoma developed eight (14.8%) than the femoral site procedure 19 (35.1%) among post-cardiac catheterized patients (n=108) as displayed in table 4. Besides, 78 were diabetic and 30 were nondiabetic patients participated. The rate of hematoma development identified in diabetic patients was 19 (24.3%) and in the non-diabetic, the hematoma ratio was eight (26.6%). The large established hematoma

development rate was 03 (3.8%) in diabetic patients while it was not developed in non-diabetic patients (table 5). Comparatively, 52 patients were hypertensive but 56 patients were not. The rate of hematoma developed among hypertensive patients was eight (15.3%) and 19 (33.9%) in non-hypertensive patients (table 6). Correspondingly the rate of hematoma development was recognized in 24 (24.7%) out of 97 obese patients. Similarly, in non-obese, the hematoma ratio was three (27.3%) out of 11 patients (table 7).

Distribution of Age Category among Study Participants


In this study 108 post-cardiac catheterization patients participated, the highest number of participants 47 were falling in the 61-80 years age group, Similarly, the lowest number of participants 6 were falling 81-100 years age group as shown in graph 1.



Graph: 1 Bar Chart Represents Age Distribution of Study Participants

Gender Distribution


The numbers of male patients were higher i.e. n= (62%) as compared to the female for the given study.



Graph: 2 Pie Chart Represents Gender Distribution of Study Participants


Distribution of Heparin

Graph: 3 shows the injection heparin received by the participants according to different units, the highest number of patients who have not received heparin and the lowest number of participants 7 received heparin dose out of 108 participants.



Graph: 3 Bar Chart represents Frequency of Heparin used for Study Participants.

Graph: 4. Reflecting most of the patients were treated by angiography procedure 88%, on the other hand, angioplasty patients were 12%.



Graph: 4 Pie Chart Represents Type of Procedures used for Study Participants

Table: 1 Association of Hematoma Development between Intervention and Control Group (n=108).

Hematoma	Groups		P-value
	Control Group n (%)	Interventional Group n (%)	
No Hematoma	36 (66.7)	45 (83.3)	0.037*
Small Hematoma	08 (14.8)	08 (14.8)	
Medium Hematoma	07 (13)	01 (1.9)	
Large Hematoma	03 (5.6)	00	
Total	54 (50)	54 (50)	

* Significant level at 0.05

Table 1. Shows a significant difference between the (Control) and (Interventional) groups regarding hematoma development post-cardiac catheterization i.e. the P values were $0.037 < 0.05$. The rate of hematoma development in (Control) group was indicated at 18 (33.3%) while after the implementation of standardized nursing care protocol the rate of hematoma development in the (Interventional) group was 9 (16.6%) out of 54 (50%) participants. Thus, there was an association of hematoma development between the (Control) group and the (Interventional) group. Furthermore, in the (Control) group, a large hematoma was developed 03 (5.6%) patients but after the implementation of standardized nursing care protocol, the large hematoma had not developed in the (Interventional) group. Moreover, the (Control) group reflected the highest percentage of 18 (33.3 %) of hematoma development as compared to the (Interventional) group 9 (16.6%).

Table: 2 Association between Hematoma Development and Site Procedures in Control

Group (n=54)

Hematoma	Site Procedures		P-value
	Radial n (%)	Femoral n (%)	
No Hematoma	22 (81.5)	14 (51.9)	0.087
Small Hematoma	03 (11.1)	05 (18.5)	
Medium Hematoma	02 (7.4)	05 (18.5)	
Large Hematoma	00	03 (11.1)	
Total	27 (50)	27 (50)	

Table: 2. This table illustrates that there was no significant difference between hematoma development and site procedures in the (Control) group post-cardiac catheterization i.e. P-value was $0.087 > 0.05$. So, there was no association between hematoma development and site

procedure in the (control) group among post-cardiac catheterized patients. Although the rate of hematoma development was represented higher in the femoral approach than the radial approach; such as in the femoral site procedure the highest number of patients 13 (48.1%) had developed hematomas. However, the rate of hematoma development was signifying 5 lowest (18.5%) in radial approach out of 27 (50%). Additionally, the large hematomas had developed three (11.1%) in femoral access, but did not develop in radial access among post-cardiac catheterized patients.

Table: 3 Association between Hematoma Development and Site Procedures in Interventional Group (n=54)

Hematoma	Site Procedures		P-value
	Radial n (%)	Femoral n (%)	
No Hematoma	24 (88.8)	21 (77.7)	0.087
Small Hematoma	03 (11.1)	05 (18.5)	
Medium Hematoma	00	01 (3.7)	
Total	27 (50)	27 (50)	

Table: 3. Clarifies that there was no significant difference between hematoma development and site procedure in (Interventional) group post-cardiac catheterization i.e. P-value was $0.087 > 0.05$. Hence, there was no association between hematoma development and site procedure in the (Interventional) group. The rate of hematoma development in radial site procedure represented 3 (11.1%) out of 27 (50%) while the medium and large hematoma did not develop in an (Interventional) group. However, in the femoral approach, 6 (22.2%) patients had developed hematomas and large hematoma had not developed. The table revealed that in the radial site procedure 24 (88.8%) patients were free from hematoma as compared to the femoral site 21 (77.7%) patients were free from hematoma development post cardiac catheterization.

Table: 4 Association between Hematoma Development and Site Procedures of study

Subjects (n=108)

Hematoma	Site Procedures		P-value
	Radial n (%)	Femoral n (%)	
No Hematoma	46 (85.2)	35 (64.8)	0.058
Small Hematoma	06 (11.1)	10 (18.5)	
Medium Hematoma	02 (3.7)	06 (11.1)	
Large Hematoma	00	03 (5.5)	
Total	54 (50)	54 (50)	

Table: 4. Points out that there was no significant difference between hematoma development and site procedure in (interventional and Control) groups i.e. P-value was $0.058 > 0.05$. The radial site procedure reflected fewer rates of hematoma development 8 (14.81%) as compared to the femoral site procedure 19 (35.1%) out of 54 (50%) post-cardiac catheterized patients. So, there was no association of hematoma development between control and interventional groups. Apart from that in the radial approach 46 (85.2%) of patients were free of hematoma while in the femoral approach 35 (64.8%) patients were no hematoma out of 54 (50%). Similarly, the large hematoma did not develop in radial site procedure while 3 (5.5%) developed in femoral access.

Table: 5 Association between Hematoma Development and Comorbidity (Diabetes

Mellitus) of Study Subjects (n=108)

Hematoma	Diabetes Mellitus		P-value
	Yes n (%)	No n (%)	
No Hematoma	59 (75.6)	22 (73.3)	0.582
Small Hematoma	10 (12.8)	06 (20)	
Medium Hematoma	06 (7.7)	02 (6.7)	
Large Hematoma	03 (3.8)	00	

Total	78	30	
--------------	----	----	--

Table: 5. This table elucidates that there was no significant difference between hematoma development and co-morbidity (diabetes mellitus) post-cardiac catheterization i.e. P-value was $0.582 > 0.05$. Therefore, there was no association between hematoma development and co-morbidity (diabetes mellitus). The rate of hematoma development was established 19 (24.3%) out of 78 diabetes patients while 8 (26.6%) out of 30 non-diabetic patients. The large hematoma 03 (3.8%) was established in a diabetic while wasn't developed in the non-diabetic patients.

Table: 6 Association between Hematoma Development and Comorbidity (hypertension) of Study Subjects (n=108)

Hematoma	Hypertension		P-value
	Yes	No	
	n (%)	n (%)	
No Hematoma	44 (84.6)	37 (66.1)	0.102
Small Hematoma	05 (9.6)	11 (19.6)	
Medium Hematoma	03 (5.8)	05 (8.9)	
Large Hematoma	00	03 (5.4)	
Total	52	56	

Table: 6, this table shows that there was no significant difference between hematoma development and co-morbidity (hypertension) post-cardiac catheterization i.e. P-value was $0.102 > 0.05$. Hence, there was no association between hematoma development and comorbidity (hypertension). The rate of hematoma development among fifty-two hypertensive patients was eight (15.38%) and 19 (33.9%) in non-hypertensive fifty-six patients. The highest score of small hematomas development appeared in 11 (19.6%) in non-hypertensive patients while the small hematomas development rate was identified five (9.6%) among hypertensive patients post cardiac catheterization.

Table: 7. Association between Hematoma Development and Comorbidity (obesity) of Study Subjects (n=108)

Hematoma	Obesity		P-value
	Yes n (%)	No n (%)	
No Hematoma	73 (75.3%)	08 (72.7%)	0.468
Small Hematoma	13 (13.4%)	03 (27.3%)	
Medium Hematoma	08 (8.2%)	00	
Large Hematoma	03 (3.1%)	00	
Total	97	11	

Table:7. This table presents that there was no significant difference between hematoma development and co-morbidity (obesity) post-cardiac catheterization i.e. P-value was $0.468 > 0.05$. Thus, there was no association between hematoma development and comorbidity (obesity). The rate of hematoma development was recognized 24 (24.7%) out of 97 obese patients. Similarly, in non-obese patients, the hematoma development mentioned three (27.3%) out of 11 patients. The highest ratio of small hematoma 13 (13.4%) was established among obese patients although 03 (27.3%) was recognized among non-obese patients post cardiac catheterization.

Discussion

Table 1. This study revealed that there was a significant difference between the hematoma development in (Control) group and after the implementation of standardized nursing care protocol (Interventional) group. The rate of hematoma development in (Control) group was indicated 18 (33.3%) while after implementation of standardized nursing care protocol the rate of hematoma development in the (Interventional) group was 9 (16.6%) out of 54 (50%) participants which support the alternative hypothesis. Thus, there was an association of hematoma

development between the (Control) group and the (Interventional) group. It's proved by the statistic that implementation of standardized nursing care protocol post-cardiac catheterization is an effective role in reducing bleeding, hematoma as well as improving patient's safety, outcomes, saving treatment time and cost ⁵¹. Furthermore, there was a positive association between nurses' knowledge about the practice and lower the rate of vascular site complications among patients in the intervention group which requires safe and standardized care, as well as qualified and skillful health providers ⁵². The rate of hematoma development post coronary intervention was reduced after implementation of good nursing care ⁵³. Additionally, another study was in Lahore Pakistan (Feroze et, al) found a positive correlation between the skills and knowledge regarding patient's safety post-cardiac catheterization, adequate skills, and knowledge which reflected on enhanced patient safety ⁵⁴.

Table 2. This study has revealed that there was no significant difference between hematoma development and site procedures in the (control) group post cardiac catheterization. Although the rate of hematoma development was represented higher in the femoral approach than the radial approach; such as in the femoral site procedure the highest number of patients 13 (48.1%) had developed hematomas. However, the rate of hematoma development was signifying 5 lowest (18.5%) in radial approach out of 27 (50%). So, there was no association between hematoma development and site procedure in the (control) group post-cardiac catheterization. Similarly, as per statistics, Tran's radial artery access is associated with lower hematoma than Tran's femoral artery access ⁵⁵. Such as evidenced by small hematoma 60 (20%) then large hematoma 28 (9.3%) in femoral approach ⁵⁶. Additionally, in this study, the large hematoma had developed three (11.1%) in femoral access but had not developed in radial access besides 77% reduction in major vascular complication by using Trans radial approach ⁵⁷. Moreover, the radial hematoma

development is preventable and easy to treat by the utilization of proper techniques that could reduce the occurrence and severity of complications^{58,59}.

Table 3. It clarifies that there was no significant difference between the rate of hematoma development and site procedure in (interventional group) post-cardiac catheterization. Hence, there was no association between hematoma development and site procedure in the (Interventional) group. The rate of hematoma development in radial site procedure represented 3 (11.1%) while in femoral site 6 (22.2%) out of 27 (50%). The Trans radial access associated local hematoma rate is 1.2% to 2.6%, which is 60% lower than in Transfemoral puncture sites^{60,61}.

Table 4. Points out that there was no significant difference between hematoma development site procedure in (Control and Interventional) groups post-cardiac catheterization. The radial site procedure reflected fewer rates of hematoma development 8 (14.81%) as compared to the femoral site procedure 19 (35.1%) out of 54 (50%) post-cardiac catheterized patients. So, there was no association of hematoma development between control and interventional groups. As per evidence, the radial access has shown a less vascular access complication rate of 27% versus femoral route 45% as well as the large hematomas were significantly higher in the femoral approach^{62,63}. In the USA the hematoma development rate was in Tran's radial approach (2.9%) Tran's femoral approach (7.7%)⁶⁴. In India forearm hematoma developed in 187 (10.7%) patients⁶⁵. Similarly, this study identified that the large hematoma was not developed in the radial site procedure while 3 (5.5%) had developed in femoral access. As per contradicting findings radial artery spasm, radial artery occlusion, are the common complications while radial artery perforation and compartment syndrome are rare complications in TRA⁶⁶.

Table 5: interpret that there was no significant difference between hematoma development and co-morbidity (diabetes mellitus) post-cardiac catheterization. The rate of hematoma development was established 19 (24.3%) out of 78 diabetes patients while 8 (26.6%) out of 30 non-diabetic patients. Therefore, there was no association between hematoma development and co-morbidity (diabetes mellitus). As per the investigation, diabetes is one of the factors which increases the risk of access site complications among post-cardiac catheterization ^{67, 68}. As per statistics, those patients who had diabetes mellitus were prone to groin complications development post coronary intervention ^{69, 70}. Another study has shown that diabetes mellitus was not a significant predictor for hematoma development ⁷¹. Diabetes is a significant variable of hematoma 44 (49%) out of 118 patients.

Table 6: shows that there was no significant difference between hematoma development and co-morbidity (hypertension) post-cardiac catheterization. The rate of hematoma development among fifty-two hypertensive patients was eight (15.38%) and 19 (33.9%) in non-hypertensive fifty-six patients. Hence, there was no association between hematoma development and comorbidity (hypertension). As per statistics hypertension was found to be a not significant variable of vascular complications ^{72, 73}. Correspondingly, hypertension is not a predictor for hematoma formation ⁷⁴. The investigator proved that high blood pressure may increase the length of time to achieve hemostasis, as medically antihypertensive is required before sheath removable ^{75, 76}. It's proved by investigators that a history of hypertension and diabetes mellitus is an associated factor of hematoma development ⁷⁷⁻⁸³. Additionally, it's concluded that high blood pressure, 180 mmHg before sheath removal more likely to develop femoral complications ⁸⁴. In hypertensive patients, the rate of hematoma development was 65 (73%) out of 118 post coronary interventions

Table 7: Presented that there was no significant difference between hematoma development and comorbidity (obesity) post-cardiac catheterization. The rate of hematoma development was recognized 24 (24.7%) out of 97 obese patients. Similarly, in non-obese patients, the hematoma development mentioned three (27.3%) out of 11 patients. Thus, there was no association between hematoma development and comorbidity (obesity). As per statistics obesity is not a significant factor of vascular complications post-cardiac catheterization⁸⁶. As per previous studies, obesity was associated with an increased risk of vascular complications post-cardiac catheterization^{87, 88}. Additionally, obesity was associated with hematoma development⁸⁹. Furthermore, obesity has been identified as a correlated factor with hematoma development post coronary intervention⁹⁰.

Strength of the Study

The current study has various strengths, which are as follow:

- The essential feature of the study was its design, quasi-experimental (post), which facilitated in carrying out interventional research.
- The implementation of the standardized nursing care protocol and data collection carried out by the researcher herself, she was experienced dealing with post-cardiac catheterized patients besides well trained under interventional cardiologists and certified for sheath and Trans radial band removal.
- Standardized nursing care protocol was effective for the reduction of hematoma post-cardiac catheterization, which has not been studied earlier in Pakistan.
- In this study the manual digital method was used for the removal of the femoral arterial sheath thus its evidence that the digital approach is an effective and safe method for the removal of femoral arterial sheaths.

- The study adopted a valid data collection tool which was further validated by experts and pilot tested on 10% of sample size on post-cardiac catheterization patients.

Limitations

- It is a single-center study with a small sample size hence it cannot be generalized.
- Data for the current study were collected through a 12-hour follow-up post-cardiac catheterization within this time was recorded but any hematoma development after that time has gone undetected.

Recommendation for Nurses

- Standardized Nursing Care protocol on post-cardiac catheterization associated training and certification should be necessary for nurses.
- In this study, the digital approach is an effective and safe method for the removal of femoral arterial sheaths thus nurses should implement the manual digital compression method on post-cardiac catheterized patients instead of vascular close devices (VCD).
- Nurses should be identified patients' health-related issues besides gaps then fill the gap through advanced-based literature.

Recommendation for Hospital Management

- The finding of the study addressed that the hospital management should provide simulation for hospital site nurses to up to date their knowledge and practice related to new research.
- Refresher courses follow-up to check nurses' knowledge and displayed checklist protocol in areas.
- It is recommended to the hospital management or policy creator to implement the standardized nursing care protocol post-cardiac catheterization for the reduction of

hematoma development in the hospital, as a result, can patient free from hematoma, diminish hospitalization, reduce financial constraints, release patient's discomfort, increase patient satisfaction and also nurses can manage these patients easily.

Recommendation for Research

- Randomized control trial (RCT), with a large sample size to confirm the effectiveness of standardized nursing care protocol on post-cardiac catheterized patients and also to reassess the association of hematoma development between procedure site (radial and femoral), comorbidities (diabetes, hypertension, and obesity) on large sample size.

Conclusion

The implementation of standardized nursing care protocol post-cardiac catheterization was effective for the reduction of hematoma development in the (intervention) group while there was an association of hematoma development between (control and interventional) groups. However, there was no association of hematoma development between procedure site (radial and femoral), comorbidities (diabetes, hypertension, and obesity) post-cardiac catheterization.

References

1. Tranberg T, Lippert FK, Christensen EF, Stengaard C, Hjort J, Lassen JF, Petersen F, Jensen JS, Bäck C, Jensen LO, Ravkilde J. Distance to invasive heart center, the performance of acute coronary angiography, and angioplasty and associated outcome in

out-of-hospital cardiac arrest: a nationwide study. *European heart journal*. 2017 Jun 1;38(21):1645-52.

2. Wasfy JH, Hidrue MK, Yeh RW, Armstrong K, Dec Jr GW, Pomerantsev EV, Fifer MA, Ferris TG. Differences among cardiologists in rates of positive coronary angiograms. *Journal of the American Heart Association*. 2015 Oct 16;4(10):e002393.
3. Shuvy M, Ko DT. Bleeding after percutaneous coronary intervention: can we still ignore the obvious?.
4. Archbold RA, Robinson NM, Schilling RJ. Radial artery access for coronary angiography and percutaneous coronary intervention. *BMJ*. 2004 Aug 19; 329 (7463):443-6.
5. Ray S. *Tips & Tricks in Interventional Cardiology*. Jaypee Brothers, Medical Publishers Pvt. Limited; 2017 Sep 30.
6. Wang Y, Tang J, Ni J, Chen X, Zhang R. A comparative study of TR Band and a new hemostatic compression device after trans-radial coronary catheterization. *Journal of Interventional Medicine*. 2018 Nov 1;1(4):221-8.
7. Manda YR, Baradhi KM. Cardiac catheterization, risks, and complications. *InStatPearls [Internet]* 2018 Sep 26. StatPearls Publishing.
8. Granchelli M. Development and Evaluation of an Evidence-Informed Protocol for the Nursing Care of Post-Cardiac Catheterization Patients.
9. <http://www.bing.com/search?q=vascular+access+site+sheath+removal&src=IE-SearchBox&Form=IE8SRC>
10. Portal MR. Care of the patient post cardiac catheterization.

11. Ali HA, Ali MM. Effect of Designed Teaching Protocol Regarding Patients' Safety after Cardiac Catheterization on Nurses' Performance and Patients' Incidence of Vascular Complications. *International Journal of Studies in Nursing*. 2019 Jan 7;4(1):107.
12. Chhatriwalla AK, Amin AP, Kennedy KF, et al. Association between bleeding events and in-hospital mortality after percutaneous coronary intervention. *JAMA* 2013;309:1022–9
13. Petrie JR, Guzik TJ, Touyz RM. Diabetes, hypertension, and cardiovascular disease: clinical insights and vascular mechanisms. *Canadian Journal of Cardiology*. 2018 May 1;34(5):575-84.
14. Cardiac Cath Pro. Vascular access. [Internet]. [Cited 2018 November]. Available from: <http://cardiaccathpro.com/VascularAccess.html>
15. Ebeed, M.E., Khalil, N.S., & Ismaeel, M.S. (2017). Vascular Complications and Risk Factors among Patients Undergoing Cardiac Catheterization. *Egypt Nurse J*, 14, 259-68. <https://doi.org/10.4103/2090-6021.233668>.
16. Granchelli M. Development and Evaluation of an Evidence-Informed Protocol for the Nursing Care of Post-Cardiac Catheterization Patients.
17. Tuttle MK, Haroian NQ, Gavin LF, Esposito CA, Ho KK. Expedited Removal of a Radial Hemostatic Compression Device Following Cardiac Catheterization Is Safe and Associated With Reduced Time to Discharge. *Cardiology Research*. 2019 Dec;10(6):331.
18. Monitoring & Managing Patients with Arterial & Venous Sheaths. (2018, January 29). Retrieved from <https://study.com/academy/lesson/monitoring-managing-patients-with-arterial-venoussheaths.html>.
19. LEARNING MODULE FOR LATE ARTERIAL & VENOUS SHEATH REMOVAL
Post Cardiac Catheterization and Percutaneous Coronary Intervention (PCI)

(DELEGATED MEDICAL FUNCTION) CC 10-060 APPROVED BY: Karen MacRury-Sweet.

20. Dvoretzky LI. TRAVEL TO COUNTRY IATROGENIC. YATROGENIYA DIAGNOSTIC PROCEDURES (Message 3). The Russian Archives of Internal Medicine No 2018. 2011 Jul 26:5.
21. [https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-\(DVDs\)](https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(DVDs)) 11 June 2021.
22. Granchelli M. Development and Evaluation of an Evidence-Informed Protocol for the Nursing Care of Post-Cardiac Catheterization Patients February 1, 2018.
23. Dols JD, Muñoz LR, Martinez SS, Mathers N, Miller PS, Pomerleau TA, Timmons A, White S. Developing policies and protocols in the age of evidence-based practice. The Journal of Continuing Education in Nursing. 2017 Feb 1;48(2):87-92.
24. Granchelli M. Development and Evaluation of an Evidence-Informed Protocol for the Nursing Care of Post-Cardiac Catheterization Patients February 1, 2018.
25. Adnan Yaqoob, M. R.-B. (n.d.). Knowledge and Practices among Nurses regarding Patient Care, following Cardiac Catheterization, At a Tertiary Care Hospital in Karachi, Pakistan.
26. Thabet OF, Ghanem HM, Ahmed AA, Abd-ElMouhsen SA. Assessment of Nurse's knowledge and practice for patients undergoing Cardiac Catheterization. Assiut Scientific Nursing Journal. 2019 Jun 1;7(17):95-101.
27. Costa GF, Sardinha DM, Oliveira VM, Barata FS, da Silva PH, Mafra IN, de Oliveira LC, da Silva Lima TF, Nunes MC. Nursing Care for Patients Submitted to Coronary

Transluminal Angioplasty: An Integrative Review. *Cardiology and Angiology: An International Journal*. 2020 Jan 9:18-26.

28. Bernat I, Horak D, Stasek J, et al. ST-segment elevation myocardial infarction treated by radial or femoral approach in a multicenter randomized clinical trial: the STEMI-RADIAL trial. *J Am Coll Cardiol* 2014;63:964–72.
29. Mason PJ, Shah B, Tamis-Holland JE, et al. An update on radial artery access and best practices for transradial coronary angiography and intervention in acute coronary syndrome: a scientific statement from the American Heart Association. *Circ Cardiovasc Interv* 2018;11:e000035. [Crossref](#) [PubMed](#).
30. Mason PJ, Shah B, Tamis-Holland JE, Bittl JA, Cohen MG, Safirstein J, Drachman DE, Valle JA, Rhodes D, Gilchrist IC, American Heart Association Interventional Cardiovascular Care Committee of the Council on Clinical Cardiology; Council on Cardiovascular and Stroke Nursing; Council on Peripheral Vascular Disease; and Council on Genomic and Precision Medicine. An update on radial artery access and best practices for transradial coronary angiography and intervention in acute coronary syndrome: a scientific statement from the American Heart Association. *Circulation: Cardiovascular Interventions*. 2018 Sep;11(9):e000035.
31. Jolly SS, Yusuf S, Cairns J, *et al* Radial versus femoral access for coronary angiography and intervention in patients with acute coronary syndromes (RIVAL): a randomized, parallel-group, multicentre trial. *Lancet* 2011;377:1409–20.
32. Dwivedi SK, Nayak G, Sharma AK, Chaudhary GK, Chandra S, Bhandari M, Vishwakarma P, Pradhan A, Sethi R, Narain VS. Forearm hematoma as a complication of

trans-radial coronary intervention: an Indian single-center experience. *American Journal of Cardiovascular Disease*. 2021;11(4):462.

33. Bernat I, Horak D, Stasek J, et al. ST-segment elevation myocardial infarction treated by radial or femoral approach in a multicenter randomized clinical trial: the STEMI-RADIAL trial. *J Am Coll Cardiol* 2014;63:964–72.
34. Mason PJ, Shah B, Tamis-Holland JE, et al. An update on radial artery access and best practices for transradial coronary angiography and intervention in acute coronary syndrome: a scientific statement from the American Heart Association. *Circ Cardiovasc Interv* 2018;11:e000035. [Crossref](#) [PubMed](#).
35. Mason PJ, Shah B, Tamis-Holland JE, Bittl JA, Cohen MG, Safirstein J, Drachman DE, Valle JA, Rhodes D, Gilchrist IC, American Heart Association Interventional Cardiovascular Care Committee of the Council on Clinical Cardiology; Council on Cardiovascular and Stroke Nursing; Council on Peripheral Vascular Disease; and Council on Genomic and Precision Medicine. An update on radial artery access and best practices for transradial coronary angiography and intervention in acute coronary syndrome: a scientific statement from the American Heart Association. *Circulation: Cardiovascular Interventions*. 2018 Sep;11(9):e000035.
36. Jolly SS, Yusuf S, Cairns J, *et al* Radial versus femoral access for coronary angiography and intervention in patients with acute coronary syndromes (RIVAL): a randomized, parallel-group, multicentre trial. *Lancet* 2011;377:1409–20.
37. Dwivedi SK, Nayak G, Sharma AK, Chaudhary GK, Chandra S, Bhandari M, Vishwakarma P, Pradhan A, Sethi R, Narain VS. Forearm hematoma as a complication of

- trans-radial coronary intervention: an Indian single-center experience. *American Journal of Cardiovascular Disease*. 2021;11(4):462.
38. Cheng KY, Chair SY, Choi KC. Access site complications and puncture site pain following trans-radial coronary procedures: A correlational study. *International journal of nursing studies*. 2013 Oct 1;50(10):1304-13.
39. Batiha AM, Abu-Shaikha HS, Alhalaiqa FN, Jarrad RA, Ramadan HJ. Predictors of complications after sheath removal post transfemoral percutaneous coronary interventions. *Open Journal of Nursing*. 2016;6(06):497.
40. Al-Makhamreh HK, Alqsous N, Ammari Z, Husari K, Nimri S, Ya'aqoub L, et al. Vascular complications following cardiac catheterization at Jordan University Hospital. *Jordan Med J*. 2014;48(4).
41. Erol F, Arslan Ş, Yüksel İÖ, Üreyen ÇM, Serdar S, İnci S, et al. Determinants of iatrogenic femoral pseudoaneurysm after cardiac catheterization or percutaneous coronary intervention via the femoral artery. *Turk Kardiyol Dern Ars*. 2015;43(6):513–9.
42. Cox N, Resnic FS, Popma JJ, Simon DI, Eisenhauer AC, Rogers CA. *Am J Cardiol*. 2004 Nov 1; 94(9):1174-7. [[PubMed](#)] [[Ref list](#)]
43. Dwivedi SK, Nayak G, Sharma AK, Chaudhary GK, Chandra S, Bhandari M, Vishwakarma P, Pradhan A, Sethi R, Narain VS. Forearm hematoma as a complication of trans-radial coronary intervention: an Indian single-center experience. *American Journal of Cardiovascular Disease*. 2021;11(4):462.
44. Sandoval Y, Bell MR, Gulati R. Transradial artery access complications. *Circulation: Cardiovascular Interventions*. 2019 Nov;12(11):e007386.

45. Cox N, Resnic FS, Popma JJ, Simon DI, Eisenhauer AC, Rogers CA. *Am J Cardiol.* 2004 Nov 1; 94(9):1174-7. [[PubMed](#)] [[Ref list](#)]
46. Al-Momani MS, AbuRuz ME. Incidence and predictors of groin complications early after coronary artery intervention: a prospective observational study. *BMC nursing.* 2019 Dec;18(1):1-0.
47. Kassem HH, Elmahdy MF, Ewis EB, Mahdy SG. Incidence and predictors of post-catheterization femoral artery pseudoaneurysms. *Egyptian Heart J.* 2013;65(3):213–21
48. Dumont CJ, Keeling AW, Bourguignon C, Sarembock IJ, Turner M. Predictors of vascular complications post diagnostic cardiac catheterization and percutaneous coronary interventions. *Dimens Crit Care Nurs.* 2006;25(3):137–42
49. Finke M, Norton D, Gassler J. Evaluation and Management of Chest Pain and Acute Coronary Syndrome (ACS) in the Emergency Department. In *Cardiology Consult Manual 2018* (pp. 15-31). Springer, Cham.
50. Deuling JH, Vermeulen RP, van den Heuvel AF, Schurer RA, van der Harst P. A randomized controlled study of standard versus accelerated deflation of the Terumo radial band hemostasis device after transradial diagnostic cardiac catheterization. *European Journal of Cardiovascular Nursing.* 2017 Apr;16(4):344-51.
51. Parach A, Sadeghi-Ghahroudi M, Saeid Y, Ebadi A. The effect of evidence-based care guidelines on outcomes after removal of the arterial sheath in patients undergoing angioplasty. *ARYA atherosclerosis.* 2018 Nov;14(6):237.
52. Ali HA, Ali MM. Effect of Designed Teaching Protocol Regarding Patients' Safety after Cardiac Catheterization on Nurses' Performance and Patients' Incidence of Vascular Complications. *International Journal of Studies in Nursing.* 2019 Jan 7;4 (1):107.

53. Rezaei-Adaryani, M., Ahmadi, F., & Asghari-Jafarabadi, M. (2009). The Effect of Changing Position and Early Ambulation after Cardiac Catheterization on Patients' Outcomes: A single-Blind Randomized Controlled Trial. *Int J Nurs Stud*, 46(8), 1047-53. <https://doi.org/10.1016/j.ijnurstu.2009.02.004>.
54. Feroze M, Afzal M, Sarwar H, Galani A, Afshan S. Knowledge and Practice of Registered Nurses about Patient Safety after Cardiac Catheterization in Punjab Institute of Cardiology Hospital in Lahore, Pakistan. *International Journal of Musculoskeletal Pain Prevention*. 2017 Jun 1;2(2):233-8.
55. Mason PJ, Shah B, Tamis-Holland JE, Bittl JA, Cohen MG, Safirstein J, Drachman DE, Valle JA, Rhodes D, Gilchrist IC, American Heart Association Interventional Cardiovascular Care Committee of the Council on Clinical Cardiology; Council on Cardiovascular and Stroke Nursing; Council on Peripheral Vascular Disease; and Council on Genomic and Precision Medicine. An update on radial artery access and best practices for transradial coronary angiography and intervention in acute coronary syndrome: a scientific statement from the American Heart Association. *Circulation: Cardiovascular Interventions*. 2018 Sep;11(9):e000035.
56. Al-Momani MS, AbuRuz ME. Incidence and predictors of groin complications early after coronary artery intervention: a prospective observational study. *BMC nursing*. 2019 Dec;18(1):1-0.
57. Ferrante G, Rao SV, Juni P, et al. Radial versus femoral access for coronary interventions across the entire spectrum of patients with coronary artery disease: A meta-analysis of randomized trials. *JACC Cardiovasc Interv* 2016;9:1419–34.

58. Bernat I, Horak D, Stasek J, et al. ST-segment elevation myocardial infarction treated by radial or femoral approach in a multicenter randomized clinical trial: the STEMI-RA Chugh SK, Chugh Y, Chugh S. How to tackle complications in radial procedures: tip and tricks. Indian heart journal. 2015 May 1;67(3):275-81.
59. Sandoval Y, Bell MR, Gulati R. Transradial artery access complications. Circulation: Cardiovascular Interventions. 2019 Nov;12(11):e007386.DIAL trial. J Am Coll Cardiol 2014;63:964–72.
60. Bernat I, Horak D, Stasek J, et al. ST-segment elevation myocardial infarction treated by radial or femoral approach in a multicenter randomized clinical trial: the STEMI-RADIAL trial. J Am Coll Cardiol 2014;63:964–72.
61. Mason PJ, Shah B, Tamis-Holland JE, et al. An update on radial artery access and best practices for transradial coronary angiography and intervention in acute coronary syndrome: a scientific statement from the American Heart Association. Circ Cardiovasc Interv 2018;11:e000035.[Crossref](#) [PubMed](#).
62. Woods SL. Sivarajan Froelicher ES. Underhill Motzer S. Bridges EJ. Cardiac nursing 6th ed. Lippincott Williams & Wilkins. 2010
63. Cardiac Cath Pro. Vascular access. [Internet]. [Cited 2018 November]. Available from: <http://cardiaccathpro.com/VascularAccess.html>
64. Dwivedi SK, Nayak G, Sharma AK, Chaudhary GK, Chandra S, Bhandari M, Vishwakarma P, Pradhan A, Sethi R, Narain VS. Forearm hematoma as a complication of trans-radial coronary intervention: an Indian single-center experience. American Journal of Cardiovascular Disease. 2021;11(4):462.

65. Sandoval Y, Bell MR, Gulati R. Transradial artery access complications. *Circulation: Cardiovascular Interventions*. 2019 Nov;12(11):e007386.
66. Femoral Arterial Sheath removal CPG Peninsula Health 2017. Available from: <http://prompt.phcn.vic.gov.au/Search/download.aspx?filename=17855073\178>
67. Woods SL, Sivarajan Froelicher ES, Underhill Motzer S, Bridges EJ. *Cardiac nursing* 6th ed. Lippincott Williams & Wilkins. 2010
68. Al-Sadi AKA, Omeish AFY, Al-Zaru IM. Timing and predictors of femoral hematoma development after manual compression of femoral access sites. *JPMA J Pakistan Med Assoc*. 2010;60(8):620
69. Batiha AM, Abu-Shaikha HS, Alhalaiqa FN, Jarrad RA, Ramadan HJ. Predictors of complications after sheath removal post transfemoral percutaneous coronary interventions. *Open Journal of Nursing*. 2016;6(06):497.
70. Al-Makhamreh HK, Alqsous N, Ammari Z, Husari K, Nimri S, Ya'aqoub L, et al. Vascular complications following cardiac catheterization at Jordan University Hospital. *Jordan Med J*. 2014;48(4).
71. Al-Sadi AKA, Omeish AFY, Al-Zaru IM. Timing and predictors of femoral hematoma development after manual compression of femoral access sites. *JPMA J Pakistan Med Assoc*. 2010;60(8):620
72. Al-Momani MS, AbuRuz ME. Incidence and predictors of groin complications early after coronary artery intervention: a prospective observational study. *BMC nursing*. 2019 Dec;18(1):1-0.
73. Woods SL, Sivarajan Froelicher ES, Underhill Motzer S, Bridges EJ. *Cardiac nursing* 6th ed. Lippincott Williams & Wilkins. 2010

74. Femoral Arterial Sheath removal CPG Peninsula Health 2017. Available from:
<http://prompt.phcn.vic.gov.au/Search/download.aspx?filename=17855073\178>
75. Al-Momani MS, AbuRuz ME. Incidence and predictors of groin complications early after coronary artery intervention: a prospective observational study. *BMC nursing*. 2019 Dec;18(1):1-0.
76. Erol F, Arslan Ş, Yüksel İÖ, Üreyen ÇM, Serdar S, İnci S, et al. Determinants of iatrogenic femoral pseudoaneurysm after cardiac catheterization or percutaneous coronary intervention via the femoral artery. *Turk Kardiyol Dern Ars*. 2015;43(6):513–9.
77. Kassem HH, Elmahdy MF, Ewis EB, Mahdy SG. Incidence and predictors of post-catheterization femoral artery pseudoaneurysms. *Egyptian Heart J*. 2013;65(3):213–21
78. Dumont CJ, Keeling AW, Bourguignon C, Sarembock IJ, Turner M. Predictors of vascular complications post diagnostic cardiac catheterization and percutaneous coronary interventions. *Dimens Crit Care Nurs*. 2006;25(3):137–42
79. Lee GA. Determinants of quality of life five years after coronary artery bypass graft surgery. *Heart Lung*. 2009;38(2):91–9.
80. Al-Momani MS, AbuRuz ME. Incidence and predictors of groin complications early after coronary artery intervention: a prospective observational study. *BMC nursing*. 2019 Dec;18(1):1-0.
81. Batiha AM, Abu-Shaikha HS, Alhalaiqa FN, Jarrad RA, Ramadan HJ. Predictors of complications after sheath removal post transfemoral percutaneous coronary interventions. *Open Journal of Nursing*. 2016;6(06):497.
82. Gregory AB, Lester KK, Gregory DM, Twells LK, Midodzi WK, Pearce NJ. Impact of body mass index on short-term outcomes in patients undergoing percutaneous coronary

intervention in Newfoundland and Labrador, Canada. *Cardiology research and practice*. 2016 Jan 1;2016.

83. Cardiac Cath Pro. Vascular access. [Internet]. [Cited 2018 November]. Available from: <http://cardiaccathpro.com/VascularAccess.html>

84. Al-Momani MS, AbuRuz ME. Incidence and predictors of groin complications early after coronary artery intervention: a prospective observational study. *BMC nursing*. 2019 Dec;18(1):1-0.

85. Batiha AM, Abu-Shaikha HS, Alhalaiqa FN, Jarrad RA, Ramadan HJ. Predictors of complications after sheath removal post transfemoral percutaneous coronary interventions. *Open Journal of Nursing*. 2016;6(06):497.

86. Gregory AB, Lester KK, Gregory DM, Twells LK, Midodzi WK, Pearce NJ. Impact of body mass index on short-term outcomes in patients undergoing percutaneous coronary intervention in Newfoundland and Labrador, Canada. *Cardiology research and practice*. 2016 Jan 1;2016.

87. Cardiac Cath Pro. Vascular access. [Internet]. [Cited 2018 November]. Available from: <http://cardiaccathpro.com/VascularAccess.html>

88. Cox N, Resnic FS, Popma JJ, Simon DI, Eisenhauer AC, Rogers CAM *J Cardiol*. 2004 Nov 1; 94(9):1174-7.[PubMed] [Ref list]

89. Al-Momani MS, AbuRuz ME. Incidence and predictors of groin complications early after coronary artery intervention: a prospective observational study. *BMC nursing*. 2019 Dec;18(1):1-0.

90. Al-Sadi AKA, Omeish AFY, Al-Zaru IM. Timing and predictors of femoral hematoma development after manual compression of femoral access sites. JPMA J Pakistan Med Assoc. 2010;60(8):620

UNDER PEER REVIEW