

# **Pediatric Anthropometrics: An observational study from Indus Medical college, Tando Muhammad Khan, Sindh, Pakistan**

---

## **Abstract:**

**Background:** Children constitute a big proportion of human population and they are the future of any nation so their health is of prime importance. Child health care has the special focus in the health care policy of any system. **Study place, duration and design:** The current research was an observation study conducted during July to December 2021 in Pediatric outpatient department at Indus Medical College Hospital Tando Muhammad Khan. **Methods:** Consent was taken from parents and demographic data and anthropometric measurements were carried out and recorded on prepared proforma, frequency and percentage was calculated for various parameters **Results:** There were 956 children evaluated for anthropometric parameters out of which male were 637(66.63%) and females were 319(43.37%). The weight was normal in 603(63.10%), Low in 332(34.73%) and above normal in 21(2.20%) children. The height was seen normal 713(74.5%), short stature was seen in 201(21%) and 42(4.5%) children were taller.

**Conclusion:** Various abnormalities in child anthropometric parameters were observed.

**Key Words:** Anthropometrics, Low weight, Obesity, Short Stature

## **Introduction:**

Normal anthropometric measurements especially weight and height child health are so essential for a prosperous society as these depict the health, nutritional, social and economic status of any country or region on the globe [1]. Adolphe Quetelet, a Belgian anthropologist was the 1<sup>st</sup> to describe the impact of socioeconomic status on the anthropometric parameters in children in 18<sup>th</sup> century [2]. Quetelet's work on pediatric anthropometry was followed by various researchers around the globe in 19<sup>th</sup> century with much progress in this field [3]. Child growth and development is deeply affected by physical activity and the

environment around that is why physical education and training has been a part of child education globally [4]. Many researchers have agreed on certain factors to influence growth and development out of which the socioeconomic status and the lifestyle of an individual are the two most important ones [5]. The height of children from Poland was highly affected due socioeconomic change in the 20<sup>th</sup> century [6]. Similarly the prevalence of obesity in the children is on rise globally and reported to be 20 % in school age children in some countries [7]. Poor economic status of a family leads to malnutrition while a rich economic status of a family leads to weight gain and obesity [8]. There was lack of data on this topic from our part of land so we manage to investigate the anthropometric parameters in children visiting our pediatric outpatient department at Indus medical college hospital.

### **Methodology:**


The current research was an observation study conducted during July to December 2021 in Pediatric outpatient department at Indus Medical College Hospital Tando Muhammad Khan. Consent was taken in written from parents of children. Patients were selected through non-probability consecutive sampling technique and demographic data and anthropometric measurements were carried out and recorded on prepared proforma for the data collection. The frequency and percentage was calculated for various parameters and results were compiled and represented in the form of various tables and charts.

### **Results:**

There were 956 children evaluated for anthropometric parameters out of which 637(66.63%) were males and 319(43.37%) were females [Table-1, Fig-1]. Majority of children were 668(69.87%) above 5 years where as 288 (30.13%) children in our study participants were below 5years of age. The weight was normal in 603(63.10%), Low in 332(34.73%) and above normal in 21(2.20%)

children. The height was seen normal 713(74.5%), short stature was seen in 201(21%) and 42(4.5%) children were taller [Table-2, Fig-2].

UNDER PEER REVIEW




**Fig-1: Pie chart representation of gender distribution**

**Table-1: Demographic data of the patients**

Age	Below 5years	Above 5 years	Total patients
	288 (30.13%)	668(69.87%)	
Sex	Male	Female	956(100%)
	637(66.63%)	319(43.37%)	

**Table-2: Weight and height of distribution in study patients**

Weight	Normal	Low weight	Obese
	603(63.10%)	332(34.73%)	21(2.20%)
Height	Normal	Short Stature	Taller
	713(74.5%)	201(21%)	42(4.5%)



**Fig-2: Bar chart distribution of anthropometric parameters**

### **Discussion:**

Physical development is reported to be modified by social factors with poor socioeconomic status associated with growth retardation and lower weight and a rich socioeconomic status with obesity and weight gain [9]. The previously published data shows that the prevalence of stunting was 14% and the obesity was 5% in Pakistani children which is consistent with our findings [10]. There are approximately 156 million stunted children on the globe out of which 89 million belong to south Asia [11]. Obesity in children is assumed to result from increase intake high caloric diet, fatty meal, intake of excessive soft drinks, low physical activity affecting the physical, social and mental health of children profoundly along with unsatisfactory performance in academic career. Obesity in children may lead to many diseases like metabolic, cardiovascular, hepatic, kidney, neurological, orthopedic and pulmonary [12]. Short stature which is defined as a child's height above two standard deviations below mean age, or <3rd percentile and the same is known as Idiopathic short stature when the etiology unknown.

Majority of children presenting with short stature have usually constitutional growth delay, familial short stature or short stature of idiopathic nature only 5% children have some identified cause whereas 95% are of idiopathic nature[13]. The identifiable etiologies for short stature include celiac disease, growth hormone deficiency, genetic causes, hepatic failure, hypothyroidism, renal diseases and Turner syndromes [14]. We could not work on finding etiological factors responsible for various abnormal anthropometric parameters in our study children and we could not compare the ratio or proportion between the male and female children which were our study limitations. We are further planning a long term study project on this topic in the near future covering wider range of parameters and we also recommend similar sort of the studies from other districts of Sindh province especially the remote areas. We also recommend studies from other provinces of Pakistan so that a clear picture of the pediatric community may be drawn and the policy maker may be compelled to design proper programs for child health care.

### **Conclusion:**

Abnormalities in anthropometric parameters like weight above and below normal and height as taller as well as short stature were observed in our study children

### **References:**

1. Tanner JM. Growth as a mirror of the condition of society: secular trend and class distinction. In: Demirjian A, Brault-Dubuc M (eds) Human growth: a multidisciplinary review. Taylor and Francis, London, 1986; 3–34 25
2. Quetelet A. Sur l'homme et le développement des facultés, ou essai de physique sociale. Paris 1835;2:34–62.
3. Ward WP. The biological standard of living in comparative perspective. In: Komlos J, Baten J (eds) Birth weight and the history of modern biological living standards. Franz Steiner, Stuttgart, 1998; 303–305.
4. Śniadecki J On the physical education of a child. Dziennik Wileński T. 1805/1806; 2 nr 41.
5. Wang Y. Cross-national comparison of childhood obesity: the epidemic and the relationship between obesity and socioeconomic status. Int J Epidemiol 1910;30:1129–1136

6. Bielecki EM, Haas JD, Hulanicka B. Secular changes in the height of Polish schoolboys from 1955 to 1988. *Econ Hum Biol* 2012; 10:310–317.
7. Kułaga Z, Litwin M, Tkaczyk M, Palczewska I, Zajączkowska M, Zwolińska D. Polish 2010 growth references for school-aged children and adolescents. *Eur J Pediatr* 2011;170:599–609
8. Andegiorgish AK, Wang J, Zhang X, Liu X, Zhu H. Prevalence of overweight, obesity, and associated risk factors among school children and adolescents in Tianjin, China. *Eur J Pediatr* 2012;171:697–703.
9. Beata Gurskowska , Zbigniew Kułaga , Mieczysław Litwin , Aneta Grajda , Anna Świąder , Katarzyna Kułaga et al. The relationship between selected socioeconomic factors and basic anthropometric parameters of school-aged children and adolescents in Poland *Eur J Pediatr* 2014;173:45–52
10. Sina Aziz, Wajeeha Noor-ul-Ain, Rukhsana Majeed, M. Amanullah Khan, Iftikhar Qayum, Intisar Ahmed Growth centile charts (anthropometric measurement) of Pakistani pediatric population *J Pak Med Assoc* 2012; 62(4):367-377.
11. Grantham-McGregor S, Cheung YB, Cueto S, Glewwe P, Richter L, Strupp B. International Child Development Steering Group Developmental potential in the first 5 years for children in developing countries. *Lancet* 2007; 369: 60-70.
12. Krushnapriya Sahoo, Bishnupriya Sahoo, Ashok Kumar Choudhury, Nighat Yasin Sofi, Raman Kumar, Ajeet Singh Bhadoria. Childhood obesity: causes and consequences *Journal of Family Medicine and Primary Care* 2015; 4(2):187-192.
13. Lindsay R, Feldkamp M, Harris D, Robertson J, Rallison M. Utah Growth Study: growth standards and the prevalence of growth hormone deficiency. *J Pediatr*. 1994;125(1):29–35.
14. Lashari SK, Korejo HB, Memon YM. To determine frequency of etiological factors in short statured patients presenting at an endocrine clinic of a tertiary care hospital. *Pak J Med Sci*. 2014; 30(4):858–861.