

# Severity of splenic trauma and its surgical outcome at Liaquat University Hospital

## ABSTARCT

**Objective:** To determine the severity of splenic trauma and its surgical outcome among patients presented at surgical emergency with abdominal blunt trauma at Liaquat University Hospital Jamshoro/Hyderabad.

**Material and methods:** This prospective case series study was conducted in the department of Surgery at Liaquat University Hospital Jamshoro, from December 2017 to 5th November 2017. Patients having diagnosis abdominal trauma with diagnosis of splenic injury, age above 18 years and both genders were included. All the patients underwent surgical treatment as per indications. Patients were monitored during Hospital stay to assess the outcome. Study proforma was used for the data collection and SPSS version 26 was used for the data analysis.

**Results:** The study included 60 patients; their mean age was  $28.3 \pm 12.4$  years. Most of the cases 51(84.9%) were between 18 to 40 years of age groups. Males were predominant with a male to female ratio of 3.1:1. Most of the cases had Grade III and Grade IV of injury as 48.3% and 33.3% respectively and none of our patients was diagnosed as Grade V by the scan. Among 30 cases developed complications, particularly as 10 cases had pulmonary infection, 08 cases had intra-abdominal bleeding, wound infection was in 9 and one had developed over-whelming post splenectomy (OPSS). Overall mortality rate was 5.0%.

**Conclusion:** Splenic trauma and its severity in patients having blunt abdominal trauma was observed to be the most frequent. Better outcome was observed and it depends on the timely arrival, proper diagnosis and skill an experience of the surgeons.

**Key words:** Splenic trauma, complications, surgical management.

## INTRODUCTION

In blunt abdominal trauma, the spleen is one of the most commonly injured organs.<sup>1</sup> The treatment of individuals with blunt splenic damage has evolved from surgical and to non-surgical care in recent decades.<sup>1</sup> Trauma is the leading cause of death and morbidity among people under the age of 50 worldwide and in Pakistan recently reported that that the out of 130 blunt trauma cases, 65% had spleen injuries.<sup>2</sup> Hemodynamic instability with a rising pulse rate and lowering blood pressure is the most dependable indicator of splenic injuries. Splenic damage can cause pain in the left hypochondrium, as well as pain in the left shoulder tip and widespread abdominal pain. Left lower chest damage has been observed in 43 percent of individuals with splenic injuries.<sup>3,4</sup> Whether there are other organs injuries, a higher injury severity score, and increased age, more likely to die from a spleen injury. Mortality rate following splenic damage is 6 percent to 10%, according to trauma facilities in North America.<sup>3,5</sup> The traumatic incident remains one of the leading causes of the mortality in people under the age of 40, and it reflects a long-standing clinical challenge with social and economic consequences, like the direct cost to society (treatment) and indirect cost to society (rehabilitation and loss of productivity) both influence the community and the person.<sup>6</sup> The surgical therapy at the turn of the 19th century was splenectomy, which was done to avoid traumatic amputation and had a postoperative death rate of 40% compared to non-intervention, which might reach 90%.<sup>1</sup> Splenectomy may be needed in individuals with splenomegaly (e.g., tropical, lymphoma, portal hypertension, auto-immunity, ) or in any scenario where the trauma surgeon believes non-operating management is not safe or appropriate.<sup>7,8</sup> Splenic trauma can be formerly classified based on anatomical lesions. Patients' circumstances, on the other hand, may necessitate an emergency transfer to the operating room (OR) without the chance to determine the grade of splenic lesions prior to surgical investigation.<sup>9</sup> The most frequent method of splenectomy is still open surgery. In addition, several centers use laparoscopic and robotic procedures.<sup>10,11</sup> Splenectomy can result

in infectious, pulmonary, pancreatic, hemorrhagic and thromboembolic consequences. Most prevalent consequences are atelectasis and pneumonia in the lower lobes. Hemorrhages due to insufficient local hemostasis, wound infection, subphrenic abscess, infective morbidity caused by encapsulated bacterial involvement, pseudocyst and fistula caused by traumatic injuries to the pancreas at the time of splenic hilus dissection, and thromboembolic complications caused by the hematological alteration are all possible complications.<sup>10,12,13</sup> However the mortality rate following splenectomy varies depending on the indications, however it is 0.97-6.04 times that of the general population.<sup>10</sup> However this study has been conducted to evaluate the severity of splenic trauma and its surgical outcome among patients presented at surgical emergency with abdominal blunt trauma at Liaquat University Hospital Jamshoro/Hyderabad

## MATERIALS AND METHODS

This prospective case series study was conducted in the department of Surgery at Liaquat University Hospital Jamshoro, from December 2017 to November 2017. Patients having diagnosis abdominal trauma with diagnosis of splenic injury, age above 18 years and both genders were included. Patients with multi organ injuries and those who did not agree to participate in the study were excluded. Non probability consecutive sampling technique was used. Complete history, general physical examination, systemic examination and base line investigations such as Blood CP, X-ray abdomen and Chest Ultrasound, abdomen and chest, Urine DR, Random Blood Sugar, Virology Test, Urea & Creatinine and Electrolytes as well as Specific investigation such as CT scanning of abdomen and chest and peritoneal lavage were done. All the patients underwent surgical treatment as per indications. Patients were monitored during Hospital stay to assess the outcome. Study proforma was used for the data collection and SPSS version 26 was used for the data analysis.

## RESULTS

The study included 60 patients, their mean age was  $28.3 \pm 12.4$  years, ranging from 18 to 60 years. In the present study, males were predominant 45 (75.0%), with a male to female ratio of 3.1:1.

According to the severity of the splenic trauma, the Grade I was seen in 1(1.6%) patient and Grade II was observed in 10(16.6%) patients by CT scan whereas Grade III and Grade IV were seen in 48.3% and 33.3% respectively and none of our patients was diagnosed as Grade V by the scan. Table No. 2

The morbidity in this study occurred in 30 patients (50.0%), 10 cases of pulmonary infection, Intra-abdominal bleeding was seen in 08(13.3%) cases, wound infection was present in 9(15.0%) and one (1.6%) developed over-whelming post splenectomy (OPSS), There were three deaths (5.0%) in this study. Table.3

Table no. 1. Discriptive statistics of age and gender n=60

| Varibales   |         | Statistics      |       |
|-------------|---------|-----------------|-------|
| Age (years) | Averag  | 28.3±12.4 years |       |
|             | Minimum | 18 years        |       |
|             | Maximum | 60 years        |       |
| Gender      | Male    | 45              | 75.0% |
|             | Female  | 15              | 25.0% |

Table no. 2 Distribution of patients according to sevrity of splenic injury and surgical outcomes n=60

| Variables                 |           | Frequency | Percentage |
|---------------------------|-----------|-----------|------------|
| Sevrity of splenic taruma | Grade-I   | 01        | 1.6%       |
|                           | Grade-II  | 10        | 16.6%      |
|                           | Garde-III | 29        | 48.3%      |

|                  |  |    |       |
|------------------|--|----|-------|
|                  | Grade-IV                               | 20 | 33.3% |
| <b>Morbidity</b> | Intra-abdominal bleeding               | 08 | 13.3% |
|                  | Pulmonary infection                    | 10 | 16.6% |
|                  | Wound infection                        | 09 | 15.0% |
|                  | Sub phrenic abscess                    | 02 | 3.33% |
|                  | Thrombocytosis                         | 01 | 1.66% |
|                  | Portal vein thrombosis                 | 02 | 3.33% |
|                  | Pseudo cyst formation                  | 01 | 1.66% |
|                  | Overwhelming postsplenectomy infection | 01 | 1.66% |
|                  | <b>Mortality</b>                       |    | 03    |

## DISCUSSION

The spleen is the most usually injured solid organ in the blunt trauma of the abdomen, as well as the most commonly injured abdominal structure after blunt trauma. This study also has been done to assess the severity of splenic trauma and its surgical outcome among patients presented at surgical emergency with abdominal blunt trauma. In this study mean age of the participants was  $28.3 \pm 12.4$  years, and males were predominant 45 (75.0%), with a male to female ratio of 3.1:1. Consistently Mehboob A et al<sup>14</sup> reported that the most of the cases 56%, were found with age group of 21-40 years and in the line of this study, they found males in majority 93% and only 7% were females. On other hand Emre A et al<sup>10</sup> reported that the average age of the cases was  $50.51 \pm 22.63$  years and inconsistently they found females in majority as 55.4% and remaining 4.5% were males. Above difference according to age and gender with international study may because in our societies males are mostly involved in outdoor activities, therefore they are on high risk on these types' traumas.

In this study according to the severity of the splenic trauma, the Grade I was seen in 1(1.6%) patient and Grade II was observed in 10(16.6%) patients by CT scan whereas Grade III and Grade IV were seen in 48.3% and 33.3% respectively and none of our patients was diagnosed as Grade V by the scan. Similarly Mehboob A et al<sup>14</sup> reported that the splenic injuries were graded as 1(4%) case had grade I, among 3 (12%) cases grade II, in the 11 (44%) of the cases the injury grade was III, grade IV injury was seen in 10 (40%) of the cases and grade V injury was in only one case. In another study of twenty patients had Grade IV injuries, and eight cases had Grade V injuries, out of a total 28 cases having high-grade injuries. On other hand Chalya PL et al<sup>16</sup> also found similar findings. In the line of this study Gangat SA et al<sup>17</sup> reported that the grade III injuries were the most prevalent form of splenic injury 47%, and Grade IV injuries were 34%.

In this study the morbidity in this study occurred in 30 patients (50.0%), 10 cases of pulmonary infection, Intra-abdominal bleeding was seen in 08(13.3%) cases, wound infection was present in 9(15.0%) and one (1.6%) developed over-whelming post splenectomy (OPSS). The most prevalent post-splenectomy consequences are lower lobe atelectasis and pneumonia, in addition, hemorrhages due to insufficient local hemostasis, infective complications from a wound infection, subphrenic abscess, and encapsulated bacteria, pseudocyst and fistula formation due to traumatic injuries of the pancreas during the dissection of the splenic hilus and coagulation abnormalities from hematological variations could occur in individuals.<sup>10,18</sup> Consistently Gangat SA et al<sup>17</sup> demonstrated that the 15.9% pulmonary infection, 11.3% septicemia, 6.8% pancreatic fistula, 2.2% subphrenic abscess, 13.6% wound infection and 2.2% pneumococcal pneumonia were post operative complications. In this study mortality rate was (5.0%), while Gangat SA et al<sup>17</sup> demonstrated that the overall mortality rate was seven (15.9%). Splenic injuries account for 25% of all solid abdominal organ injuries worldwide, and the death rate linked with splenic trauma is estimated to be between 7 and 18%.<sup>16,19</sup> The duration of hospital stay amongst trauma individuals has been demonstrated to be a significant marker of morbidity. Prolonged hospitalization places an unacceptably high demand on health-care resources and reduces the population's productive potential due to time wasted in the hospital and impairment.<sup>16</sup> In order to provide appropriate trauma treatment for victims of splenic injuries, the challenges found in the management of the individuals having splenic injuries in our environment must be addressed.<sup>16</sup>

## CONCLUSION

Splenic trauma and its severity in patients having blunt abdominal trauma was observed to be the most frequent. Better outcome was observed and it depends on the timely arrival, proper diagnosis and skill an experience of the surgeons.

## REFERENCES

1. Petrone P, Peña MF, Staffolani MJ, Brathwaite C, Axelrad A, Esparragón JC. Evolution of the treatment of splenic injuries: from surgery to non-operative management. *Cirugía Española (English Edition)*. 2017 Oct 1;95(8):420-7.
2. Latif A, Alim M, Khan B, Khan KA, Kumar J. Study of Spleen and Liver Injuries in Patients of Blunt Trauma Abdomen. *Pakistan Journal of Medical & Health Sciences*. 2022 Apr 12;16(03):320-.
3. Al-Busaidi A, Al-Shafei T, Al-Moqbali H, Al-Kindi S, Al-Saadi M, Al-Busaidi N, Al-Qadhi H. The Incidence of Splenic Injury Following Blunt Abdominal Trauma (BAT), Sultan Qaboos University Hospital Experience. *Surgical Science*, 2017, 8, 312-18
4. Hildebrand DR, Ben-Sassi A, Ross NP, Macvicar R, Frizelle FA, Watson AJ. Modern management of splenic trauma. *Bmj*. 2014 Apr 2;348.
5. Brady RR, Bandari M, Kerssens JJ, Paterson-Brown S, Parks RW. Splenic trauma in Scotland: demographics and outcomes. *World journal of surgery*. 2007 Nov;31(11):2111-6.
6. Perrotta G, Guerrieri E, Guerrieri M. Splenic trauma: Definition, classifications, clinical profiles and best treatments. *Open Journal of Trauma*. 2021 Oct 21;5(1):019-36.
7. Birindelli A, Martin M, Khan M, Gallo G, Segalini E, Gori A, Yetasook A, Podda M, Giuliani A, Tugnoli G, Lim R. Laparoscopic splenectomy as a definitive management option for high-grade traumatic splenic injury when non operative management is not feasible or failed: a 5-year experience from a level one trauma center with minimally invasive surgery expertise. *Updates in Surgery*. 2021 Aug;73(4):1515-31.
8. Lopez Jr JM, McGonagill PW, Gross JL, Hoth JJ, Chang MC, Parker K, Requarth JA, Miller PR. Subcapsular hematoma in blunt splenic injury: A significant predictor of failure of nonoperative management. *Journal of Trauma and Acute Care Surgery*. 2015 Dec 1;79(6):957-60.
9. Coccolini F, Montori G, Catena F, Kluger Y, Biffl W, Moore EE, Reva V, Bing C, Bala M, Fugazzola P, Bahouth H. Splenic trauma: WSES classification and guidelines for adult and pediatric patients. *World Journal of Emergency Surgery*. 2017 Dec;12(1):1-26.
10. Emre A, Sanli AN. Splenectomy indications and postoperative follow-up results of a department of general surgery. [Annals of Medical Research](#). 2018;25(2):241-5
11. Berelavichus SV, Smirnov AV, Ionkin DA, Kriger AG, Dugarova RS. [Robot-assisted and laparoscopic partial splenectomy for nonparasitic cysts]. *Khirurgiia (Mosk)* 2015;(7):41-8
12. Tsutsumi N, Tomikawa M, Akahoshi T, Kawanaka H, Ota M, Sakaguchi Y, et al. Pancreatic fistula after laparoscopic splenectomy in patients with hypersplenism due to liver cirrhosis: effect of fibrin glue and polyglycolic acid felt on prophylaxis of postoperative complications. *Am J Surg* 2016;212(5):882-8.
13. Wang X, Li Y, Crook N, Peng B, Niu T. Laparoscopic splenectomy: a surgeon's experience of 302 patients with analysis of postoperative complications. *Surg Endosc* 2013;27(10):3564-71
14. Mehboob A. Non-operative Management of Blunt Splenic Trauma. *Journal of Surgery Pakistan*. 2019 Oct 31;24(3):136-9.

15. Nijdam TM, Spijkerman R, Hesselink L, Leenen LP, Hietbrink F. Predictors of surgical management of high grade blunt splenic injuries in adult trauma patients: a 5-year retrospective cohort study from an academic level I trauma center. *Patient safety in surgery*. 2020 Dec;14(1):1-9.
16. Chalya PL, Mabula JB, Giiti G, Chandika AB, Dass RM, Mchembe MD, Gilyoma JM. Splenic injuries at Bugando Medical Centre in northwestern Tanzania: a tertiary hospital experience. *BMC Research notes*. 2012 Dec;5(1):1-9.
17. Gangat SA, Rehman AN, Khaskhali A, AYUB M, FATIMA S, MEMON IA. Splenic trauma—management in relation to mode and grade. *Pak. J. Surgery*. 2008 Jan 15;24(1):2-4.
18. Merchant RH, Shah AR, Ahmad J, Karnik A, Rai N. Post splenectomy outcome in  $\beta$ -thalassemia. *Indian J Pediatr* 2015;82(12):1097-100.
19. Alli N: Management of blunt abdominal trauma in Maiduguri: a retrospective study. *Niger J Med* 2005, 14(1):17-22.

UNDER PEER REVIEW