

**Ecological risk assessments of Heavy metal contamination in water, Sediment and Biota of Otamiri and Imo Rivers, Nigeria**

**ABSTRACT**

Heavy metals pollution of the environment is a serious concern because of the hazardous effects they pose to human health, various ecosystems and other biological receptors. The aim of this study is to assess the ecological risk of exposure to heavy metals and physicochemical parameters around Otamiri and Imo rivers in Etche and Oyibo Local Government Areas of Rivers state. Water, Sediment, Arrow head plant (*Sagittaria latifolia*) and Christmas bush Plant (*Ceratopetalum gummiferum*) and Tilapia fish samples (*Oreochromis niloticus*) and Cat Fish (*Clarias gariepinus*) were collected at different points from the five sample stations for wet and dry seasons. These samples were analyzed quantitatively to assess the levels of contamination in the samples using the ecological risk assessment models to assess the risks associated with exposure to these contaminants. The results of the heavy metal analysis show that the Estimated Daily Intake (EDI) for biota samples ranged from 0.000 – 1.254 mg/kg/day while water ranged from 0.000 -0.322 mg/kg/day. The target hazard quotient (THQ) ranged from 0.000 – 69.228 for both biota and river water samples for Otamiri and Imo rivers. The hazard index values indicated significant public health risk with values of 1.42 was recorded for Tilapia and 33.14 for Christmas bush plant. Cancer risk due to heavy metals exposure had a peak value of  $1.04 \times 10^{-1}$  for wet season. The bioaccumulation of the metals showed that the pollution load index values ranged from 2.467- 5.822 in wet season and 9.708 – 99.853 in dry season. The presence of these heavy metals above threshold values suggest that the aquatic life that inhabit such rivers risk bioaccumulation which may affect the human population that depends on Otamiri/Imo rivers for fishing, drinking or irrigation. These poses a danger if bioaccumulation was to take place over a long period of time.

**Keywords: Ecological risks, Heavy metal, Bioaccumulation, Aquatic plants, fishes, sediment, Rivers.**

**Introduction**

Most water pollutants originate from human activity, while a small percentage of them have their sources in natural activities such as volcanic eruptions. Human activities in urban areas render surface waters including streams, river, lagoons, etc. and ground water bodies susceptible to contamination from organic pollutants such as heavy metals, industrial chemicals, sewage discharged from sources like urban runoffs, industrial activities, subsurface infiltration, or atmosphere precipitation effluents, sewage treatment

plants, chemical fishing activities, leachate from decomposing refuse dumps, agricultural fertilizer applications (Nwankwoala and Ekpewerechi, 2017). Once pollutants are introduced into receiving surface water and groundwater bodies via discharge processes, they are transported within the water cycle. Of more serious concern is the fact that heavy metals can cause serious health effects with different symptoms depending on the nature and quality of the metal ingested (Nwankwoala and Ekpewerechi, 2017). Most of these organic pollutants bind to cellular proteins and DNA resulting in biochemical

disruptions and cell damage leading to mutations, developmental malformations, tumors, and cancer in humans (Ramesh *et al.*, 2011). Lipmann, (2009). Some other pollutants come primarily from occupational hazards on workers exposed to mixtures of Polycyclic aromatic hydrocarbons (PAHs) such as benz[a]anthracene, benzo[b]fluoranthene, indeno[1,2,3-cd]pyrene, benzo[k]fluoranthene, dibenzo[a,e]pyrene which are known to cause increased risk skin, lung, bladder and gastrointestinal cancers. Cancer Index (2019) show that ingestion of these contaminants can increase current cancers burden in Nigeria with 102,100 people diagnosed yearly while about 71,600 dying from cancer yearly.

The impact of anthropogenic activities on Otamiri and Imo Rivers as a result of the increasing rate of urbanization and other industrial activities because of the closeness of the vicinity and linkage to Trans-Amadi industrial layout shows that Otamiri River is a major tributary that receives effluents from the activities and operations of companies operating within the flanks of the rivers in Rivers State. Worrying is the fact that water from these rivers serves as a source of water distributed for public consumption and domestic uses. These rivers are vulnerable to surface water pollution because effluents from industries, municipal wastes, agricultural and urban run-off are discharged into it thereby deteriorating the water quality (Ogbonna *et al.*, 2021). Heavy metals and polycyclic

aromatic hydrocarbons (PAHs) come from industrial and urban activities. As pollutants, these compounds are toxic and many are identified

as carcinogenic, mutagenic and teratogenic. They stress the ecosystem, lowering its resilience so that the health of the ecosystem is weakened and becomes less productive, causing increased mortality of aquatic life and poor water quality. Eventually the compounds enter the food web as pollutants through the filter-feeders such as bivalves and worms which then bioaccumulate, threatening the health of those higher up in the food web – a major concern for people who eat seafoods as a major diet.

Other human impacts affecting water resources include Oil spills through accidental discharges from ships at sea, ballast water from ships, reclaiming land by drainage, building bridges (structures such as sea walls, bridges, drains, jetties, marinas, levy banks, rock walls and breakwaters can all affect the passage of water and can change the conditions), sand extraction for construction aggregates, rubbish/ocean dumping (litter on beaches or rivers and floating in the salt and fresh water pollutes) contribute grossly to water pollution. These impacts on estuaries contribute to the reduction of habitats for estuarine animals and plants and spoil recreational activity and the beauty of our estuarine coastline. They also jeopardize the role estuaries play in

maintaining the health of coastal waters, including the marine organisms within them, which affects the fishing industry. Industries like fishing and oyster farming can impact the water quality and also the movement of water in such rivers. Over-fishing and habitat loss reduces the amount of fish in the ocean and disrupts the food chain.


However, this fragile ecosystem has been degraded seriously because of environmental changes such as global climate change and environmental pollution (Ogbonna *et al.*, 2021). Among various pollutants, heavy metals with persistence, non-biodegradation, toxicity and bioavailability pose a major threat to rivers biodiversity and human health. Unfortunately, the fragile ecosystem of Otamiri and Imo Rivers are increasingly being threatened by these anthropogenic activities. Therefore the aim of this study was to assess the Ecological risks associated with contamination of heavy metals in water resources as well as its impact on the biota of the ecosystems. Ecological risk assessment focuses on evaluating the impacts of human activities on ecological systems and the services they provide. Ecological risk

assessment has its roots in estimating exposure and risk or hazard from chemicals in the environment, with range of stressors facing ecological receptors (McIntosh and Pontius, 2017).

## **Materials and Methods**

### **Description of Study Area**

The Imo River is one of the major rivers in the Southern part of Nigeria. It originates in the vicinity of Okigwe and takes a Southerly course until it is joined by the Otamiri River which flows from the neighborhood of Owerri. It then proceeds eastward meandering close to Akwete from where it moves towards the Imo tidal basin as a smooth straight river. Imo River is considered as part of Niger Delta River basin. It is a deep freshwater river which cuts across Imo State, Abia State and Rivers State. The river flows through coastal plain, alluvium and mangrove swamp and empties into the Atlantic Ocean through the Opobo creek at the Bight of Benin (Akoma, 2008). The study area lies within the Etche and Oyibo LGA axis of Otamiri and Imo River in Rivers State. The dimension of the study area is approximately 94665.46 km<sup>2</sup> as shown in Figure 1.



**Figure 1: Map of the study area showing Sampling Stations.**  
 Source: NDDC GIS laboratory.

2

4

5

**Table 1: Description of sampling stations**

S/N	Sampling Stations	Location	Coordinates	
			Latitude	Longitude
1	Station 1	Otamiri River at Umuechem Etche LGA, Rivers State	5° 0' 38.1" N	7° 3' 22.7" E
2	Station 2	Otamiri River at Chokocho Etche LGA, Rivers State.	4° 59' 37.0" N	7° 3' 22.7" E
3	Station 3	Imo River at Odagwa Etche LGA, Rivers State.	4° 58' 54.1" N	7° 10' 16.7" E
4	Station 4	Imo River at Umuebulu Oyibo LGA, Rivers State.	4° 53' 37.0" N	7° 8' 36.8" E
5	Station 5	Imo River at Imo Gate, Oyibo LGA, Rivers State.	4° 53' 16.3" N	7° 8' 42.1" E

## Sampling Technique

The purposive sampling frame was adopted in this study. Five sampling stations were established along the Imo River and its adjoining tributary along Otamiri River covering a distance of approximately five kilometers. Notably, samples were collected for both wet and dry seasons in this study. The description of the sampling stations using the coordinates is shown in Table 1.

## Sample Collection and Preparation

Water, sediment and biota (fish and plant) samples were collected from Otamiri and Imo rivers during the wet and dry seasons. Selection of organisms was based on the different niches in which they normally thrive within the river ecosystem. The fish samples were identified by a fishery expert and a hydrobiologist and eventually assigned an aquarium number. The biota (fish and plants) samples, collected in this study consisted of two species namely, tilapia and catfish while the Arrow head plant (*Sagittaria latifolia*) and Christmas bush plant (*Ceratopetalum gummiferum*) were the plants collected from the sampling stations.

The fish and plant samples were preserved in a dark bottle away from sunlight to guard against sunlight induced photo-degradation. Samples were preserved at  $-4^{\circ}\text{C}$  using an ice pack. Sediment samples were collected from the top 10-cm layer of the river bed.

Sediment samples were air-dried for 14 days to reduce moisture and sieved using 2mm sieve. The sieved samples were mashed to powder and stored in the refrigerator until analysis.

Water samples were collected in transparent and brown glass bottles prewashed with detergent, rinsed with water and acetone (99.9%) and dried before sample collection. Samples were taken from 0.1 m below the water surface and transported directly to the laboratory. All samples were sent to laboratory and biota samples stored in refrigerator at  $-4^{\circ}\text{C}$  until the analysis.

## Physicochemical characteristics

The physico-chemical parameters of the water sample namely, pH, temperature, turbidity, conductivity, total dissolved solids, total suspended solids, total alkalinity, biological oxygen demand, chemical oxygen demand, dissolved oxygen, total organic carbon, sulphate, nitrate, phosphate, salinity, chlorine, bromine, and conductivity were all determined in this study using standard procedures as listed in Table 2

- i. The conductivity was determined using conductivity meter – model SM 3030 manufactured by surgifield medicals, England.
- ii. Turbidity was determined using turbidity meter model - SGZ –

- 200BS made by the American brewing industry turbidity unit.
- iii. Dissolved oxygen was determined using dissolved oxygen meter – model DO 5509, made in Taiwan.
  - iv. Salinity was determined by salinity meter HI98203 made by Hanna Instruments, Romania.
  - v. The pH was determined using pocket series waterproof digital pH meter.
  - vi. Conductivity was measured using an electrical conductivity meter (EC meter) from Mettler Toledo, People's Republic of China.
  - vii. TDS tester model TDS SCAN 30 manufactured by ASBL America was used in determining TDS
  - viii. Chlorine and bromine were determined using a 3-way pool and SPA test kit manufactured by ASTRAL –US, China.
  - ix. Temperature was determined using a Liquid Crystal Thermometer
  - x. The BOD meter OxiDirect was used in determining the biochemical oxygen demand.

### **Heavy metal Determination**

The EPA vigorous digestion method described by Ekeanyanwu *et al* (2010) was adopted. For the water sample, 10ml of concentrated HNO<sub>3</sub> was added to 100ml of each of the

representative water samples in a Pyrex beaker. The samples were boiled slowly and then evaporated on a hot plate to the lowest possible volume (20ml). The beakers were allowed to cool and another 5ml of concentrated HNO<sub>3</sub> was added and heated until digestion was complete. The samples were evaporated again to dryness (but not baked) and the beakers were cooled, followed by the addition of 5ml of HCl solution (1:1 v/v). The solutions were then warmed and 5ml of 5M NaOH was added, then filtered. The filtrates were transferred to 100ml volumetric flasks and diluted to the mark with distilled water. These solutions were then used for the elemental analysis using atomic absorption spectrophotometer (UNICAM 929 AAS).

Heavy metals (Pb, Cd, Ni, V, Cr), and metalloid (As) in both the sediment and biota was determined using Solar Thermo Elemental Flame Absorption Spectrometer (S4 710). To accurately determine the presence of the heavy metals of interest, sediment and biota samples were air dried to reduce moisture and 9 ml of 65% concentrated HNO<sub>3</sub> and 3ml perchloric acid was used

to digest 5g of each biota and sediment samples. The solution was then transferred to a hot plate at 110°C for 5 hours. Afterwards, the samples were introduced into an oven under a temperature that was gradually increased in 100°C every 60 minutes until the desired final temperature of 450°C was reached eighteen hours later; white ashes were obtained. Following this, samples were left to cool. The white ashes were then dissolved with 1.5% HNO<sub>3</sub> (5ml) and a final volume of 25ml was made by adding deionized water. The resulting solution was filtered using a Whatman filter paper (number 42) fitted into a Bucher funnel into a beaker before it was transferred into a tightly sealed plastic container.

### **Heavy Metal Risk Assessment Method**

Significant carcinogenic and non-carcinogenic human health risk associated with heavy metals (Cd, Hg and Pb) exposure from samples was determined using estimated daily intakes (EDI), target hazard quotient (THQ), hazard index (HI) and Carcinogenic risk.

### **Estimated Daily Intakes (EDI) of Metals**

The daily intake of metals depends on the metal concentration in sample, the daily food consumption rate and the body weight. The estimated daily intake (EDI) of metals is a concept introduced to take into account these factors. The EDI was calculated based on the following formula by Singh *et al* (2010).

$$EDI = \frac{C_{\text{metal}} \times D_{\text{food intake}}}{BW_{\text{average}}}$$

Where *C* is the metal concentration in food in mg/kg, *D* is the daily intake of food in kg person<sup>-1</sup>, and BW is average body weights for children (4 to 9 years old), adolescents (10 to 19 years old), male adults (20 to 65 years old) are 24kg, 54.5kg, 70.0kg, respectively (Falco *et al.*, 2003).

### **Target Hazard Quotient (THQ)**

Non-carcinogenic risk estimation of heavy metals consumption was determined using THQ values and Hazard Index Values. THQ is a ratio of the determined dose of a pollutant to a reference level considered harmful. THQ values were determined based on

the following formula by Singh *et al* (2010).

$$THQ = \frac{Efr \times ED \times FIR \times C}{RfDo \times Baverage \text{ wt} \times ATn \times 10^{-3}}$$

where Efr is exposure frequency assumed to be 365 days year<sup>-1</sup>, ED is exposure duration in 56 years equivalent to an average lifetime, FIR is average daily consumption in kg person<sup>-1</sup>day<sup>-1</sup>, (0.116 g/day, 0.109 g/day and 0.097 g/day for adults, adolescents and children respectively (Ihedioha *et al.*, 2014), C is concentration of metal in food sample in mg/kg, RfDo is reference dose in mg/kg day<sup>-1</sup>, and ATn is average exposure time for non-carcinogens in days. Reference doses of 0.003, 0.001, 1.5, 0.0371, 0.0035, 0.0016, 0.03 and 0.007 mg/kg/day for As, Cd, Cr, Cu, Pb, Hg, Zn and Fe respectively were used (Kamunda *et al.*, 2016)

### Ecological Risk Assessment

The determination of ecological Risk was determined using two different models namely; Bioaccumulation factor and pollution load index.

### Determination of Bioaccumulation Factor (BAF)

The bioaccumulation factor (BAF) is determined using the formula below

$$BAF = \frac{\text{Conc of metal in plants collected}}{\text{Conc of analyzed metal in soil (mg/kg)}}$$

BAF value greater than 1 indicates high accumulation potential of the biota (Mendil *et al.*, 2005).

### Pollution Load Index (PLI)

Pollution Load Index PLI using the Tomlinson's approach (Tomlinson *et al.*, 1980). The PLI represents how much a metal concentration exceeds average and normal background levels. It also provides a summation of the overall heavy metal toxicity found in an examined sample. The control samples were taken to represent natural background. The summation of the BAF gives the PLI (Adamo *et al.*, 2005).

This was achieved using Equations (7);

$$PLI = (BAF_1 \times BAF_2 \times BAF_3 \times \dots \times BAF_n)^{1/n}$$

where,

BAF: Bioaccumulation factor, n = number of metals = 7;

C<sub>sample</sub>: Mean metal concentrate in the plants;

$C_{\text{background}}$ : Mean metal concentration in the soil/background/substrate.

## RESULTS

### Physico-Chemical Analysis of Water Samples

The physico-chemical characteristics of the water samples is presented in Table 3. pH is one of the vital environmental characteristics that decides the survival, metabolism, physiology and growth of aquatic organisms. The pH values range of water sample across the stations was between 5.56 and 5.80 as compared to the WHO limit of 6.5-8.5. The pH showed significant variations across the sample stations at  $p\text{-value} < 0.05$ . The pH is not less than 4 ( $<4$ ) hence is suitable for aquatic life. Notably, Odagwa had the highest pH value of 5.80 while Umuechem and Umuebulu 5.56. Electrical Conductivity which is a measure of the total ionic composition of the water samples had mean values of  $73.00 \mu\text{S/cm}$  and  $113.50 \mu\text{S/cm}$ , which were below the WHO permissible limit of  $500 \mu\text{S/cm}$ . There were remarkable variations in the electrical conductivities of water in the sample stations. Imo Gate  $92.50 \mu\text{S/cm} < \text{Umuechem}$ ,  $73.00 \mu\text{S/cm} < \text{Chokocho}$ ,  $94.50 \mu\text{S/cm} < \text{Umuebulu}$ ,

$104.50 \mu\text{S/cm} < \text{Odagwa}$   $113.50 \mu\text{S/cm}$ . Odagwa had the highest conductivity value of  $113.50 \mu\text{S/cm}$  while Umuechem had  $73 \mu\text{S/cm}$ . Statistical test a  $p\text{-value} < 0.05$  showed that there was a significant difference.

Turbidity gives an expression of the scattering and absorption of light through water. Turbidity level is increased by the presence of clay, silt, fine organic and inorganic matter, soluble coloured organic compounds, plankton and other microscopic organisms. Turbidity values for water samples was recorded as 3.00 NTU to 27.50 NTU

The Total Dissolved Solids (TDS) value of  $62.50 \text{mg/l}$  was obtained from Odagwa while Umuechem had  $39.00 \text{mg/l}$  as against the permissible limit of  $500 \text{mg/l}$  Total Suspended Solids (TSS) is the dry-weight of suspended particles that are not dissolved in a water sample that can be trapped by a filter and analyzed. Table 3 and figure 2 TSS values for water samples across stations ranged from  $2.00 \text{mg/l}$  to  $37.00 \text{mg/l}$ . Umuechem and Chokocho had  $6.50 \text{mg/l}$  and  $2.00 \text{mg/l}$  respectively, while Odagwa, Umuebulu and Imo Gate Stations had  $37.00 \text{mg/l}$ ;  $31.50 \text{mg/l}$  and  $30.50 \text{mg/l}$  respectively. Biochemical Oxygen Demand ( $\text{BOD}_5$ ) BOD is a measure of the amount of oxygen that bacteria will require to decompose organic matter under aerobic condition. The values for BOD ranged from

2.31 mg/l to 2.76 mg/l. Chemical Oxygen Demand (COD) is a measure of the total quantity of oxygen required to oxidize all

organic material into carbon IV oxide (CO<sub>2</sub>) and water. The water samples had 4.56mg/l and 5.12mg/l.

**Table 3: Physico-chemical Characteristics of Surface water**

Parameters	Sampling Stations					WHO Limits	P- value
	Chokocho	Odagwa	Imo River	Umuechem	Umuebulu		
pH	5.71±0.23	5.80±0.22	5.70±0.50	5.56±0.69	5.56±0.55	6.5-8.5	0.000
Temperature (°C)	23.35±3.61	24.25±2.90	24.25±2.33	25.55±0.78	25.40±0.57	26.5-32	0.033
EC (µS/cm)	94.50±19.09	113.50±17.68	92.50±23.33	73.00±12.73	104.50±14.85	500	0.000
TDS (mg/l)	52.00±7.07	62.50±6.36	49.00±12.73	39.00±5.66	56.50±6.36	500	0.000
Turbidity (NTU)	3.00±2.83	31.00±11.31	27.50±9.19	6.50±2.12	26.00±11.31	5	0.269
TSS (mg/l)	2.00±1.41	37.00±31.11	30.50±28.99	6.50±2.12	31.50±27.58	25	0.000
BOD (mg/l)	2.46±0.51	2.43±0.32	2.76±0.20	2.40±0.13	2.31±0.29	4	0.002
COD (mg/l)	5.12±0.13	4.78±0.18	4.56±0.08	5.00±0.01	4.73±0.17	120	0.000

**Keys:** EC- Electrical Conductivity, TDS- Total Dissolved Solids (mg/l), TSS- Total Suspended Solids, BOD-Biological Oxygen Demand, and COD-Chemical Oxygen Demand

**Note:** Values represents mean±SD

### Heavy Metals concentration in Water

Table 4 shows the mean concentration of heavy metals (Arsenic, Cadmium, Chromium, Copper, Lead, Mercury, Zinc and Iron) in water across the sample stations. The concentration of Arsenic in water from the sample stations was below detectable limits (BDL). From the Umuechem station, the Cd mean value was 0.34 mg/kg while Chokocho, had 0.29mg/kg, Odagwa 0.30mg/kg, Umuebulu

0.32mg/kg and Imo Gate 0.23mg/kg. Umuechem had for Copper 0.02mg/kg, Odagwa 0.05 mg/kg, Chokocho 0.04mg/kg, Umuebulu 0.05mg/kg while Imo Gate 0.02mg/kg. Umuechem, Chokocho, Odagwa and Umuebulu had same Pb value of 0.12mg/kg respectively while Imo Gate had 0.52mg/kg. Hg in water was absent below detectable limit. Umuechem recorded for Zn 0.24 mg/kg, Chokocho 0.23mg/kg, Odagwa 0.22mg/kg and Umuebulu 0.17 mg/kg while Imo Gate had 0.18 mg/kg. It

differed significantly at p-value <0.05. The value of Fe at Umuechem was 4.68mg/kg, while Chokocho 3.39mg/kg and Odagwa

had 0.94mg/kg, also Umuebulu station had 1.40 mg/kg while Imo Gate recorded 1.12mg/kg.

**Table 4: Heavy metal Concentrations in Surface Water**

Parameters (mg/kg)	Sampling Stations					WHO Limits	P-value
	Chokocho	Odagwa	Imo River	Umuechem	Umuebulu		
Arsenic	BDL	BDL	BDL	BDL	BDL	0.01	-
Cadmium	0.29±0.22	0.30±0.21	0.23±0.15	0.32±0.19	0.34±0.16	0.003	0.003
Chromium	BDL	BDL	BDL	BDL	BDL	0.05	0.695
Copper	0.05±0.04	0.04±0.02	0.02±0.01	0.02±0.01	0.05±0.04	2	0.002
Lead	0.12±0.17	0.12±0.17	0.52±0.73	0.12±0.17	0.12±0.17	0.01	0.180
Mercury	BDL	BDL	BDL	BDL	BDL	0.001	-
Zinc	0.23±0.10	0.22±0.05	0.18±0.04	0.24±0.09	0.17±0.07	3	0.000
Iron	0.39±0.02	0.94±0.97	1.12±0.21	4.68±4.89	1.40±0.98	0.3	0.152
Calcium	5.06±0.84	4.88±1.44	3.78±0.79	2.69±0.80	4.49±0.68	200	0.000
Magnesium	1.08±0.08	1.06±0.04	1.13±0.08	1.11±0.13	1.36±0.20	50	0.000
Sodium	0.57±0.61	1.17±0.08	1.46±0.19	1.07±0.07	1.26±0.07	200	0.000
Potassium	1.71±0.81	0.21±0.10	0.82±0.23	1.23±0.15	0.36±0.15	NIL	0.132
Chloride	15.03±13.68	19.36±21.55	15.24±13.39	12.38±10.50	18.13±18.63	250	0.000
Nitrate	2.45±1.51	2.31±1.12	2.24±0.74	1.82±0.93	2.33±1.03	50	0.000
Sulphate	2.05±0.35	6.36±5.01	6.12±4.53	3.00±2.91	6.04±5.14	200	0.000
Ammonia	0.28±0.06	0.59±0.19	0.57±0.12	0.48±0.16	0.66±0.13	1.5	0.004
Phosphate	0.12±0.02	0.17±0.06	0.17±0.04	0.10±0.04	0.19±0.04	0.1	0.127
Hardness	13.00±4.24	16.00±11.31	13.00±4.24	8.00±5.66	15.00±7.07	200	0.000

### Heavy Metal concentration in Fishes and Plants

Table 5 show the concentration of heavy metals in Tilapia and Catfish samples. Concentration of Cadmium ranged from 0.32 to 0.42 mg/kg in Tilapia fish, Lead ranged from 0.2mg/kg in Tilapia fish to 0.9 mg/kg in Catfish, Zinc recorded 0.32 mg/kg

(Catfish) to 0.54 mg/kg (Tilapia) and Iron ranged from 0.88 mg/kg in Catfish to 1.56 mg/kg in Tilapia fish. Table 6 shows heavy metal

concentrations in two plant species namely Arrow Head Plant (*Sagittaria latifolia*) and Christmas Bush Plant (*Ceratopetalum*


*gummiferum*). Generally, it was observed that all heavy metals analysed except Iron was significantly high in Christmas Bush

Plant(*Ceratopetalum gummiferum*) compared to *Sagittaria latifolia*.

**Table 5: Heavy metals in fishes**

Parameters	Tilapia	Catfish	WHO Limit	P-value
<b>Arsenic</b>	BDL	BDL	1.4	-
Cadmium	0.42±0.03	0.32±0.09	0.05	0.176
Chromium	BDL	BDL	1	-
Copper	0.49±0.63	0.04±0.02	3	0.111
Lead	0.2±0.01	0.9±1.18	0.5	0.67
Mercury	BDL	BDL	0.5	-
Zinc	0.54±0.14	0.32±0.10	30	0.013**
Iron	1.56±1.13	0.88±0.40	100	0.013**

*Note: Values represents mean±SD; BDL represents Below Detectable Limit; \*\* Significant*



**Figure 2: Heavy metal mean concentrations in plants**

### Target Hazard Quotient and Hazard Index of Heavy Metals

The Target Hazard Quotient (THQ) and Hazard Index (HI) from the heavy metals in water and biota for adults and children for both seasons are shown in Table 7 to 10. Table 7 shows the seasonal variation in the target hazard quotient and hazard index of heavy metals in surface water for adults in dry season. Arsenic, Chromium and Mercury recorded zero concentration across the five sampling stations, The target hazard quotient and hazard index of heavy metals in surface water for Cadmium ranged from 6.50mg/kg to 11.68mg/kg, Copper 0.02mg/kg to 0.04mg/kg, Lead ranged from 3.59mg/kg to 15.75mg/kg, Zinc ranged from

0.02mg/kg to 0.03mg/kg, Iron ranged from 1.95 mg/kg to 61.44mg/kg. Table 8 shows the seasonal variation in the target hazard quotient and hazard index of heavy metals in surface water for adults in wet season. Arsenic, Chromium and Mercury recorded also recorded zero across the five sampling station, Concentration target hazard quotient and hazard index of heavy metals in surface water for Cadmium ranged from 17.81mg/kg to 23.73mg/kg, Copper 0.05mg/kg to 0.09mg/kg, Lead recorded 0.02mg/kg across the five stations, Zinc ranged from 0.04mg/kg to 0.05mg/kg, Iron ranged from 2.87 mg/kg to 15.81 mg/kg.

**Table 6: Seasonal Variation in the Target Hazard Quotient and Hazard Index of Heavy Metals in Biota for adults (Dry Season)**

Metals	Arrow Head Plant ( <i>Sagittaria latifolia</i> )	Christmas Bush Plant ( <i>Ceratopetalum gummiferum</i> )	Tilapia ( <i>Oreochromis niloticus</i> )	Cat Fish ( <i>Clarias gariiepinus</i> )
As	7.54	9.24	0.00	0.00
Cd	0.81	0.89	0.72	0.42
Cr	0.00	0.00	0.00	0.00
Cu	0.02	0.07	0.04	0.00
Pb	0.36	0.78	0.09	0.82
Hg	0.00	0.00	0.00	0.00
Zn	0.01	0.01	0.00	0.00

Fe	0.42	0.37	0.56	0.28
HI	9.15	11.36	1.42	1.53

**Table 7: Seasonal Variation in the Target Hazard Quotient and Hazard Index of Heavy metals in Biota for adults (Wet Season)**

Metals	Arrow Head Plant ( <i>Sagittaria latifolia</i> )	Christmas Bush Plant ( <i>Ceratopetalum gummiferum</i> )	Tilapia ( <i>Oreochromis niloticus</i> )	Cat Fish ( <i>Clarias gariepinus</i> )
As	0.00	0.00	7.05	7.05
Cd	0.66	0.65	0.65	0.68
Cr	0.00	0.00	0.00	0.00
Cu	0.00	0.00	0.00	0.00
Pb	0.09	0.03	0.09	0.09
Hg	0.00	0.00	0.00	0.00
Zn	0.00	0.00	0.00	0.00
Fe	0.18	0.14	0.32	0.19
HI	0.93	0.83	8.12	8.02

## DISCUSSION

The safety of water and biota regarding contaminants such as heavy metals, is a major concern due to their deleterious impacts on human health. More attention is gained when these contaminants are found in drinking water which is most frequently consumed by the most vulnerable population. Heavy metals persist naturally in the environment, however elevated concentrations in the food chain are primarily linked to man-made environmental pollution. Cadmium and Lead are reported

as the most toxic heavy metals, while over consumption of some essential mineral elements also can result to poisonous effects (Unak *et al.*, 2007).

Heavy metals are a threat for human health due to their potential damage to human cells and the skeletal system, osteoporosis, kidney failure and lung and blood cancer (Kumar *et al.*, 2007). The routes of entry for these heavy metals are ingestion or inhalation (Mendil, 2006). The toxicity of heavy metals is dependent on a number of factors, the most critical of which are

ingestion rate and route, bioavailability, gender, age, excretion rate and the chemical state of the heavy metals (Enb *et al.*, 2009). Among environmental pollutants, heavy metals are of particular concern because they are known for their persistent behaviour in the environment and are harmful, because of their ability to bioaccumulate in the environment causing damage to plants and animal tissues. (Censi *et al.*, 2006; Zheng *et al.*, 2013). Heavy metals such as lead (Pb), cadmium (Cd), mercury (Hg), zinc (Zn), and chromium (Cr), are introduced into the environment from geogenic (weathering) and anthropogenic sources. It is difficult to measure the input of metals in the environment as a result of anthropogenic activities. Concentration of heavy metals in aquatic ecosystem has increased considerably due to the inputs of industrial waste, sewage, runoff and agricultural wastes (Li *et al.*, 2010). In this study the following heavy metals; Arsenic, Cadmium, Chromium, Copper, Lead, Mercury, Zinc and Iron were analyzed in surface water and Biota samples. Result obtained reviewed that Arsenic, Chromium and Mercury were below detectable limit across the sample stations which was far below the permissible limit of 0.01mg/kg by WHO. Similar trend was also recorded in the surface water samples with Umuechem recording the highest concentration than the

Imo Gate station. The high concentrations of these heavy metal especially Lead, Cadmium etc. recorded in this study can result to exposures of aquatic flora and fauna to unacceptable effects apart from its vital role in carbohydrate metabolism (i.e., glucose tolerance and glycogen synthesis). Among the hazardous heavy metals, Cadmium and Lead recorded values above the WHO permissible limits. However, Sediments are an important sink and long-term store of a variety of pollutants, particularly heavy metals, and may serve as an enriched source of food for benthic organisms in estuarine ecosystems because they are in constant flux with the overlying water column (Bai *et al.*, 2010; Deng *et al.*, 2010; Ayejuyo *et al.*, 2010). The occurrence of increased concentrations of heavy metals in sediments can be a good indicator of man-induced pollution rather than natural enrichment of the sediment by geological weathering (Adebowale *et al.*, 2008; Wang *et al.*, 2010). However, metals cannot always be fixed by sediments permanently. Some of the sediment-bound metals may be remobilized and released into water as a result of changes in environmental conditions that leads to acidification and reduction/oxidation and impose adverse effects on living organisms (Liu 2009).

The presence of heavy metals in two fish species (Tilapia and Catfish) showed that all heavy metals analysed except Lead was higher in Tilapia when compared to Catfish. Cadmium in Tilapia has 0.42mg/kg while the Catfish recorded 0.32mg/kg. Lead concentration in Tilapia fish was 0.2mg/kg while Catfish had 0.9mg/kg. The mean concentration of Zinc in Tilapia was 0.54mg/kg while the Catfish had a mean of 0.32mg/kg. Iron concentration in Tilapia fish was 1.56 mg/kg while Catfish had 0.88mg/kg. The result of heavy metal concentrations recorded in this study are similar to those obtained by Ogbonna and Origbe (2021), who carried out a research on heavy metal concentration of surface water, sediment and fishes impacted by crude oil pollution in Bodo/Bonny River, Nigeria, which revealed that the concentration of metals like Iron, Zinc, Chromium, Copper, Cadmium and Lead in sediment indicating pollution. It also shows that the accumulation of heavy metals is predominant in sediments rather than of water and organisms, which can be linked to the fact that sediments act as an important host for all toxic metals, contaminants and dead organic matter descending from the ecosystem above. Fishes ingest heavy metals from the surrounding waters, planktons, other feeding diets and sediments resulting to their accumulation in reasonable amounts

(Olawusi-Peters and Akinola 2017). Metals such as Copper and Zinc are essential for metabolism in fish at low concentrations while some others such as Lead and Cadmium are toxic to living organisms (Virha et al., 2011). When present at high concentrations, these metals impose serious damage to metabolic, physiological and structural systems of organisms in the aquatic environment. These therefore shows the impact of anthropogenic and industrial activities to the concentration of heavy metal. Although, low concentration of heavy metals such as zinc and copper can cause stress in fish which can lead to smaller body weight, thereby reducing their ability to compete for food and habitat. This phenomenon suggests that metal levels in the surrounding biota are very low and are not interfering with the normal metabolic processes of the fishes. This finding showed that the fishes can bioaccumulate these metals and therefore may not be safe for human consumption.

### **Physico-Chemical Analysis of Water Samples**

Physicochemical properties in water serve as measure of water contamination and are usually evaluated to determine the quality of the water body. Management of water is done to ensure that contaminants that gets into it does not exceed the set or permissible limits, thus, the quality of water is related to the expected use of the water

for fishing, recreation, or wild life (Amaku and Akani, 2016). In this study, certain physicochemical parameters such as: pH, Temperature, Electrical Conductivity, Total Dissolved Solids (TDS), Turbidity, Total Suspended Solids (TSS) Biochemical Oxygen Demand (BOD) and Chemical Oxygen Demand (COD) of the surface water from Otamiri and Imo River in Etche LGA, Rivers State were evaluated.

pH is important in water quality assessment as it influences many biological and chemical processes within a water body (Meregini-Ikechukwu *et al.*, 2020). The results of pH showed that the mean values varied in all the sample stations. The mean pH values recorded in this study were below the acceptable range of 6.5 - 8.5 as prescribed by the regulatory agency (NSDWQ, 2008) and 7.0 to 8.5 by WHO. The pH values recorded ranged from  $5.56 \pm 0.69$  to  $5.80 \pm 0.22$  which were slightly acidic. The pH range for optimum growth of most aquatic bacteria is pH 6.5 and 8.5 (Rheinheimer, 1974), which were in the range of values reported in this study. The variations in pH observed is in consonance with Obunwo and Chukwudi (2015), who reported that variations in pH in surface water were due to the inherent geochemical properties of the formations of chemicals being carried. The pH of water is a property that plays vital role in different biological functions since it has the tendency to impact

on the enzymes, hormonal balance, proteins, growth as well as control the metabolism of biota and dissolution of minerals (Yao and Byme, 2001).

Temperature values recorded in the water samples of Otamiri and Imo River in Etche were reasonably warm with consistent temperatures, value within the mesophilic range. The overall mean temperature value for each station was highest in Umuechem and Chokocho station had the least. The range of values were normal for water in the tropics, and are attributed to weather conditions of the study area which is characterized by hot dry season and cold wet season (Akpan *et al.*, 2015). The difference in the water temperature between the stations were expectedly due to the various activities at the sampling stations, but statistically at  $p\text{-value} < 0.05$  was obtained when tested against WHO temperature standards, showed a significant difference between the means across the sample stations.

Electrical conductivity (EC) in natural waters is the normalized measure of the ability of the water to conduct electricity. Water with EC values between 500 and  $1000 \mu\text{s/cm}$  are not usually recommended for human consumption, and such waters are also not suitable for irrigation except for high salt tolerant crops with special techniques of management (Kuyeli *et al.*, 2009). The mean electrical conductivity (EC) values

measured in the water sample from the five sampled stations of Otamiri and Imo River were generally low. EC values measured in Odagwa was the highest followed by Umuebulu while Umuechem recorded the least value ( $73.00 \pm 12.73$ ). Akpan *et al* (2015) explained that precipitation, fresh water discharge and low temperature conditions do not favour high concentration of ionized substances in water. Ephraim and Ajayi (2015), considered waters with EC values less than  $250 \mu\text{s}/\text{cm}$  as excellent waters. Olawusi-Peters(2010) stated that the distribution of fish generally follows environmental quality parameters such as conductivity, which is an index of the total ionic content of water, and indicates the freshness or otherwise of the water. Conductivity values from this study show that the sample stations contains appreciable amount of dissolved ions thus forming a saline barrier for the survival of sensitive organisms. It is important in fish spawning habitat, as many species are sensitive to abrupt changes in the salinity of their environment.

Total Dissolved Solids (TDS) are the inorganic matters and small amounts of organic matter, which are present as solution in water. The TDS values for the five (5) sample stations were below the standard or allowable value of the TDS by WHO. The values found in the water samples were all below the maximum limit

of  $500 \text{mg}/\text{l}$ . The highest TDS value of  $62.50 \text{mg}/\text{l}$  was obtained from Odagwa and the least TDS values of  $39.00 \text{mg}/\text{l}$  was recorded for Umuechem.

Total suspended solids (TSS) are the dry-weight of suspended particles that are not dissolved, in a water sample that can be trapped by a filter and analyzed. TSS values for the sample stations ranged from  $2.00 \text{mg}/\text{l}$  to  $37.00 \text{mg}/\text{l}$ . Umuechem and Chokocho had mean values of  $6.50 \text{mg}/\text{l}$  and  $2.00 \text{mg}/\text{l}$  respectively, lower than the WHO permissible limit while Odagwa, Umuebulu and Imo Gate Stations had higher values of  $37.00 \text{mg}/\text{l}$ ,  $31.50 \text{mg}/\text{l}$  and  $30.50 \text{mg}/\text{l}$  respectively. High concentration of suspended solids can lower water quality by absorbing light hence warmer water and lesser potential of water to hold oxygen necessary for aquatic life.

Biological Oxygen Demand (BOD), is a measure of the biological activities in a water body, which gives an indication of the organic load of water bodies, especially those receiving organic effluent. Biochemical Oxygen Demand ( $\text{BOD}_5$ ) is used as an index to determine the amount of dissolved oxygen required by aerobic biological organisms (microorganisms) to decompose organic materials and also biological activity in the water. Hence high concentrations of BOD is an indication of organic pollution (Ngah *et al.*, 2017). The

values for BOD ranged from 2.31 mg/l to 2.76 mg/l..

Chemical Oxygen Demand (COD) is a measure of the total quantity of oxygen required to oxidize all organic material into carbon IV oxide (CO<sub>2</sub>) and water. The sampled water had mean value range between 4.56mg/l and 5.12mg/l. All sample stations had mean values below the WHO permissible limit. When subjected to statistical test the values differed significantly at p-value of 0.000

### **Estimated Daily Intakes of Heavy Metals (EDI)**

Heavy metals are considered as one of the most important constituents of food contamination from the environment due to its ability to persist, accumulate, and become toxic to living organism through consumption. Human exposure to heavy metals occurs via various routes such as oral ingestion, dermal contact or inhalation. The degree of toxicity of heavy metals to human health is directly related to their daily intake. The first step in the non-carcinogenic analysis is the computation of the estimated daily intake (EDI) values.

### **Bioaccumulation Factors and Pollution Load Index**

The values obtained for bioaccumulation factor, pollution load index and modified contamination degree respectively in biota (fishes and plants) for both seasons show

that the sample stations were moderately polluted by Cadmium, Copper, Lead and Iron, while Zinc pollution was severe in the plants and fish species in the wet season. For dry season the biota from the sample stations were very severely polluted by Cadmium, Lead and Iron while Copper and Zinc excessively polluted the biota. Aquatic organisms have the ability to accumulate heavy metals from various sources including sediments, soil erosion and run off, air depositions of dust and aerosol and discharges of waste water (Asaolu, 1998). For this reason, monitoring of fish tissue contamination may serve as early warning indicator of contamination or related water quality problems (Kpobari *et al.*, 2013). Several studies have indicated that fish and shellfishes are able to accumulate and retain heavy metals from their environment depending upon exposure concentration and duration as well as salinity, temperature, hardness and metabolism of the animals (Karthikeyan *et al.*, 2007, Olawusi-Peters *et al.*, 2014a & 2015). Fishes absorb heavy metals into their tissue directly from the water across the surface of the gills and ingestion of sediment particles containing heavy metals as part of their feeding process (Lorenzo *et al.*, 2003, Cruz-Rodriguez and Chu, 2003). Also, phytoplankton can take up high levels of heavy metals from contaminated waters and when shrimps consumed planktons

containing elevated levels of heavy metals, the heavy metals become concentrated in them (Janssen and Scholz, 1979). Also, their levels of contamination are of great interest as they pose health risk to those humans who consume them

### **Conclusion**

This study has shown clearly that significant risks are inherent in both biota and water samples obtained from the different sample stations. This is attributed to contamination from organic pollutants such as heavy metals, industrial chemicals, sewage discharged from sources like urban runoffs, industrial activities, subsurface infiltration, or atmosphere precipitation effluents, sewage

### **COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

### **References**

Adamo, P., Arienzo, M., Imperato, M., Naimo, D., Nardi, G., and Stanzione, D. (2005). Distribution and partition of heavy metals in surface and sub-surface sediments of Naples city port. *Chemosphere*, 61(6), 800–809.

Adebowale, K.O., Agunbiade F.O. and Olu-Owolabi, B.I. 2008. Impacts of natural and anthropogenic multiple source of pollution on the environmental conditions of Ondo

Akoma, O. C. (2008). Phytoplankton and nutrient dynamics of a tropical estuarine system, Imo River Estuary,

treatment plants, chemical fishing activities, leachate from decomposing refuse dumps, agricultural fertilizer applications. The cancer risk from the present study showed that the highest burden of likely cancer cases was from lead and arsenic. Notably, the cancer risk due to exposure to cadmium was relatively lower in this study. The total cancer risk for both biota and water samples from Umuechem, Chokocho, Odagwa, Umuebulu and Imo gate were all above the recommended safety margin thereby indicating the risk of cancer on the general population and exposed age groups of children and adults.

Nigeria. *African Research Review*, 2(2), 253–264.

Asaolu, S.S, ( 1998). Chemical pollution studies of coastal waters of Ondo State. Ph.D Thesis, Federal University of Technology, Akure, , 15-17

Ayejuyo, O.O., Omojole, M.O. and Ojoniyi, O.J. 2010. Sediment load and bioconcentration of heavy metals by shrimp (*Peaneus notalis*) from Epe Lagoon, Nigeria. *Environmental Monitoring Assessment*. 16(3): 295-301

Bai, J., Cui, B., Chen, B., Zhang, K., Deng, W., Gao, H. and Xiao, R. 2010. Spatial

- Boonyatumanond, R. M., Murakami, G., Wattayakorn, A., Togo, and H. Takada, "Sources of polycyclic aromatic hydrocarbons (PAHs) in street dust in a tropical Asian megacity, Bangkok, Thailand," *Science of the Total Environment*, vol. 384, no. 1–3, pp. 420–432, 2007.
- Censi, P., Spoto, S. E., Saiano, F., Sprovieri, M., Mazzola, S., Nardone, G., Di Geronimo, S. I., Punturo, R., and Ottonello, D. (2006). Heavy metals in coastal water systems. A case study from the northwestern Gulf of Thailand. *Chemosphere*, 64(7), 1167–1176.
- Cruz-Rodriguez, L.A. and Chu, F.L.E. (2003). HSP70 levels in oyster *Crassostrea virginica* exposed to cadmium sorbed to algal food and suspended clay particles. Intergovernmental Oceanographic Commission of UNESCO, Santiago de Compostela, Spain. Pp. 555-567.
- Deng, H.A., Zhan, J., Wang, D., Chen, Z. and Zu, C. 2010. Heavy metal pollution and distribution and ecological risk assessment of heavy metals in surface sediments
- Enb, A., Abu Donia, M. A., Abd-Rabou, N. S., Arab, A. A. K. and El-Senaity, M. H. (2009). Chemical composition of raw milk and heavy metals behavior during processing of milk products, *Global Veterinaria* 3(3): 268-275
- EPA. Load-based Licensing: Draft operational plan, environmental economic series, EPA Publication 96/38. Environmental Protection Authority, Sydney; 1996.
- Essam Nasher,<sup>1</sup> Lee Yook Heng,<sup>1,2</sup> Zuriati Zakaria,<sup>3</sup> and Salmijah Surif (2013). Assessing the Ecological Risk of Polycyclic Aromatic Hydrocarbons in Sediments at Langkawi Island, Malaysia *The Scientific World Journal* Vol. 2013 1-13
- Falco, G., Domingo, J. L., Llobet, J. M., Teixido, A., Casas, C., and Müller, L. (2003). Polycyclic aromatic hydrocarbons in foods: Human exposure through the diet in Catalonia, Spain. *Journal of Food Protection*, 66(12), 2325–2331.
- Ihedioha, J. N., Okoye, C. O., and Onyechi, U. A. (2014). Health risk assessment of zinc, chromium, and nickel from cow meat consumption in an urban Nigerian population. *International Journal of Occupational and Environmental Health*, 20(4), 281–288.
- Kamunda, C., Mathuthu, M., and Madhuku, M. (2016). Health risk assessment of heavy metals in soils from Witwatersrand gold mining basin, South Africa. *International Journal of Environmental Research and Public Health*, 13(7), 663.
- Karthikeyan. S, Palaniappan. P.L.R.M, Sabhanayakan. S. (2007). Influence of pH and water hardness upon nickel accumulation in edible fish *Cirrhinus mrigala*. *Journal Environmental Biology* 28: 484-492
- Kpobari, W. N. Mathew, O.W. and Eka B. E. (2013) Heavy metals concentrations in four selected seafood from crude oil polluted waters of Ogoniland, Rivers State, Nigeria. *Arch. Appl. Sci. Res.*, 5 (4):97-104.
- Kumar, K.S. Kumar, P.S. and Babu, M.J.R. (2007), Performance evaluation of

- Waste water treatment plant. *IJEST* 2, 7785–7796.
- Li, H. Mugabe, R. H. and Yen-Hsun, C., (2010). Accumulation of heavy metals and trace elements in fluvial River received effluents from traditional and semiconductor industries. *Scientific Reports* 6(8), 34-50.
- Lipmann, M. (2009). *Environmental Toxicants: Human Exposures and Their Health Effect*. John Wiley Sons, Inc.
- Lorenzo, J.I., Aierbe, E., Mubiana, V.K., Blust, R. & Beiras, R. (2003). Indications of regulation on copper accumulation in the blue mussel *Mytilus edulis*. In: *Molluscan Shellfish Safety*,
- McIntosh, A., and Pontius, J. (2017). Chapter 1—Tools and Skills. In A. McIntosh and J. Pontius (Eds.), *Science and the Global Environment* 1–112.
- Mendil, D., Uluözlü, Ö. D., Tüzen, M., Hasdemir, E., and Sarı, H. (2005). Trace metal levels in mushroom samples from Ordu, Turkey. *Food Chemistry*, 91(3), 463–467.
- National Research Council, *Oil in the Sea III: Inputs, Fates, and Effects*, National Academies Press, Washington, DC, USA, 2003.
- Neff, J. M. *Polycyclic Aromatic Hydrocarbons in the Aquatic Environment: Sources, Fates, and Biological Effects*, Applied Science, 1979.
- Nwankwoala, P. O., and Ekpewerechi, P. O. (2017). Human Activities and Heavy Metal Concentrations in Aba River, Abia State, Nigeria. *British J Earth Sciences Research*, 5(1), 26–36.
- Ogbonna DN, Ideriah TJ. Effect of abattoir on the physico-chemical characteristics of soil and sediment in Southern Nigeria. *Journal of Scientific Research and Reports*. 2014;3(12):1612-1632.
- Ogbonna, D.N Origbe, M. Kpormon .L.B (2021). Effect of abattoir on the physico-chemical characteristics of soil and sediment in Southern Nigeria. *Journal of Scientific Research and Reports*, 3 (12),1612-1632.
- Olawusi-Peters, O. O. (2010). Physicochemical parameters and fish richness in Agboyi Creek, Lagos state, Nigeria. *Applied Tropical Agriculture*, 15 (1&2);111-119.
- Olawusi-Peters, O.O., Ayo-Olalus, C. I. Adeyemi, T. V (2014a). Bioaccumulation of some trace element in the gills and tissues of *Clarias gariepinus* and *Oreochromis niloticus* in river Ogbese, Ondo State, Nigeria. *Journal of Environmental Chemistry and Ecotoxicology* 6(2):13-19
- Olawusi-Peters, O.O., O.E. Aguda, F.O. Okoye (2015). Heavy metals level in Shellfish (*Etheria elliptica*), water and sediments of river Ogbese, Ondo State, South West, Nigeria. *International Science Index* 1 (5): 686-689.
- Ramesh, A., Archibong, A. E., Hood, D. B., Guo, Z., and Loganathan, B. G. (2011). Global environmental distribution and human health effects of polycyclic aromatic hydrocarbons. *Global Contamination Trends of*

- Persistent Organic Chemicals*, 95–124.
- Ristola T, Pellinen J, Ruokolainen MA, Jussi VK. Effect of sedimentary type feeding level and larval density on growth and development of a midge (*Chironomus riparius*). *Solar Journal, Environmental Toxicology and Chemistry*. 1999;18:756- 764.
- Singh, A., Sharma, R. K., Agrawal, M., and Marshall, F. M. (2010). Health risk assessment of heavy metals via dietary intake of foodstuffs from the wastewater irrigated site of a dry tropical area of India. *Food and Chemical Toxicology*, 48(2), 611–619.
- Tam, N. F. Y. L. Ke, X.H.Wang, and Y. S.Wong, “Contamination of polycyclic aromatic hydrocarbons in surface sediments of mangrove swamps,” *Environmental Pollution*, vol. 114, no. 2, pp. 255–263, 2001.
- Tomlinson, D. L., Wilson, J. G., Harris, C. R., and Jeffrey, D. W. (1980). Problems in the assessment of heavy-metal levels in estuaries and the formation of a pollution index. *Helgoländer Meeresuntersuchungen*, 33(1), 566–575.
- Unak P, Lambrecht FY, Biber FZ, Darcan S (2007). Iodine measurements by isotope dilution analysis in drinking water in Western Turkey. *J. Radioanalytical Nuclear Chem.* 273:649-651.
- Villalba, A., Reguera, B., Romalde, J. L., & Beiras, R. (Eds metals in estuarine sediment cores and implications for the use of sediment quality
- Wei M.-C. and J.-F. Jen, “Determination of polycyclic aromatic hydrocarbons in aqueous samples by microwave assisted headspace solid-phase microextraction and gas chromatography/ flameionization detection,” *Talanta*, vol. 72,no. 4, pp. 1269– 1274, 2007.
- Zeng E. Y. and C. L. Vista, “Organic pollutants in the coastal environment off San Diego, California. 1. Source identification and assessment by compositional indices of polycyclic aromatic hydrocarbons,” *Environmental Toxicology and Chemistry*, vol. 16, no. 2, pp. 179–188, 1997
- Zhang, Z., Huang J, Yu, G. and Hong, H., “Occurrence of PAHs, PCBs and organochlorine pesticides in the tonghui river of Beijing, China,” *Environmental Pollution*, vol. 130, no. 2, pp.249–261, 2004.