

***In Vitro* Antisalmonellal and Antioxidant Activity of *Bauhinia rufescens* (Fabaceae) Leaf and Stem Bark Extracts**

ABSTRACT

In developing countries, enteric fever remains a major public health problem where it is endemic due to the precariousness of life hygiene combined with the abusive and inappropriate use of antibiotics. This public health problem calls for the development of new antibacterial substances that are effective, inexpensive and non-toxic for patients. *Bauhinia rufescens* is a plant used in traditional medicine for the treatment of typhoid fever by the Chadian population. **Objective:** This work aimed to evaluate the *in vitro* antisalmonellal activity of *Bauhinia rufescens* leaf and stem bark extracts against five *Salmonella* species and to study the *in vitro* antioxidant activity of leaf and stem bark extracts. **Method:** The antisalmonellal activity was evaluated using the liquid microdilution method. Antioxidant activity of these extracts was determined by investigating their DPPH[•] antiradical and iron reducing capacities. Phytochemical Screening of these extracts was performed by standard methods to justify the observed activities. **Result:** Antimicrobial activities revealed that all isolates and *salmonella* Typhi strain (ATCC6539) were sensitive to the hydroethanolic extracts of leaves and stem barks with Minimum Inhibitory Concentrations (MIC) ranging from 256 to 1024 µg/ml. Concerning, antioxidant tests, the 95% hydroethanolic extract of the leaves showed higher DPPH[•] free radical scavenging activity than all extracts. The hydroethanolic and aqueous extracts of leaves and stem barks of *Bauhinia rufescens* exhibited IC₅₀s ≤ 20 µg/ml for all extracts tested. The iron reducing power of the 95% hydroethanolic extract was higher than all other extracts at 200 µg/ml. The total phenol content of the 95% hydroethanolic extract and the flavonoid content of the 50% hydroethanolic extract of the leaves and stem barks were higher than that of all other extracts. **Conclusion:** These results showed that the 95% hydroethanolic extract of *Bauhinia rufescens* leaves possess *in vitro* antisalmonellal and antioxidant activities and could be used for *in vivo* antisalmonellal and toxicity studies.

Key words: Typhoid fevers, *Bauhinia rufescens*, antisalmonellal, antioxidant, phytochemical screening.

1. INTRODUCTION

Typhoid and paratyphoid salmonellosis are usually found in areas with poor hygiene and are a serious public health problem worldwide [1]. They mainly affect Asia, Africa and Latin America. Estimates made by the World Health Organization in 2008 report 22 million cases and 600,000 deaths per year worldwide [2]. The high costs, as well as the toxicity of some of these antibiotics are commonly encountered, following the example of chloramphenicol whose use is limited due to its bone marrow toxicity [3]. All this confronts medicine with problems of anti-infective therapy. Furthermore, typhoid salmonellosis is often exacerbated by oxidative stress, which can be a consequence of microbial invasion or militate for its progression [4, 5]. These *Salmonella* infections produce superoxide ion and nitric oxide, which react together to form peroxynitrite, which is a potent biological oxidant [6]. It then becomes necessary to find a new, effective, low-cost therapy that in addition to anti-salmonella activity can reduce the level of free radicals produced during *Salmonella* infection. The traditional use of *Bauhinia rufescens* for the treatment of typhoid fever by the Chadian population and a retrospective study on this plant revealed that it is used in traditional medicine for the treatment of several pathologies such as diabetes, diarrhea, dysentery, fungus, fibrosis, jaundice and inflammations [7, 8]. Given its traditional use by the Chadian population in the treatment of typhoid fever and the results of numerous previous scientific researches on the plant species *Bauhinia rufescens*, its extracts could constitute a non-toxic alternative against typhoid fever. Hence the objective of this work which aims to provide concrete scientific evidence on the therapeutic efficacy of *Bauhinia rufescens* against typhoid fever through the *in vitro* antisalmonellal and antioxidant study.

2. MATERIALS AND METHODS

2.1. Plant material

The plant material used for this work consisted the leaves and stem bark of *Bauhinia rufescens*, collected in December 2017 in Abeche, eastern Chad, and identified at the Unité Botanique de l'Institut de Recherche en Élevage pour le Développement (UBIRED) in N'djaména, Chad, under the reference IRED/LRVZ 1325.

2.2. Preparation of the extracts

The leaves and stem bark of *Bauhinia rufescens* were harvested and dried at room temperature away from the sun and ground using a Moulinex brand Zaiba (Super Blender, China). The powder obtained was stored in a cardboard box at room temperature, in a dry place and protected from humidity and light until use. The obtained powders were used for the preparation of different aqueous (infusion, maceration and decoction) and hydroethanolic (95%, 75% and 50%) extracts. The protocol described by Kamsu et al. [9] was scrupulously

followed for the preparation of the extracts using 250 g of powder for 2500 ml of solvent each time. The extraction yield of the different extracts was calculated according to the following formula [10].

$$R = (m \times 100) / m^{\circ}$$

R: yield of the crude extract in percentage (%), m: the mass of the crude extract obtained after extraction (g), m° : the mass of the dry plant material (g).

The different extracts obtained were stored at -20°C until their use.

2.3. Microorganisms and culture media

The microorganisms used for the determination of the antibacterial activities of the extracts were Gram-negative bacteria: *Salmonella enterica* serovar Typhi (ST), *Salmonella enterica* serovar Paratyphi A (SPA), *Salmonella enterica* serovar Paratyphi B (SPB) and *Salmonella enterica* serovar Typhimurium (STM) obtained at the Centre Pasteur du Cameroun and a *Salmonella enterica* serovar Typhi ATCC 6539 strain. Microorganisms were stored at -20°C on Salmonella-Shigella agar (SSA) (Liofilchem, Italia) and subcultures were freshly prepared before use. Mueller Hinton broth (MHB) (Liofilchem, Italia) was used as a basic enrichment medium for aerobic culture at 37°C with agitation at 150 rpm in the different tests.

2.4. Reference antibiotic and bacterial growth indicator reagent

Ciprofloxacin was used as the reference drug, while para-iodonitrotetrazolium chloride (INT) was used as an indicator of bacterial growth.

2.5. In vitro evaluation of the antisalmonellal activity of Bauhinia rufescens leaf and stem bark extracts

The bacterial growth inhibitory potential of *Bauhinia rufescens* extracts was determined by the microdilution method as described by Mativandlela et al. [11]. In each well of a 96-well microplate, 100 μl of culture broth (MHB) was introduced. Then, 100 μl of each extract was introduced to obtain an initial concentration (4096 $\mu\text{g}/\text{ml}$) respectively in the first three (03) wells of the first row; subsequently serial dilutions were performed to obtain final concentrations ranging from 2048 to 16 $\mu\text{g}/\text{ml}$. A volume of 100 μl of broth plus inoculums at the concentration of 1.5×10^6 CFU/ml bacterial was introduced into each well. Plates were brought to incubation at 37°C for 18 hours. Wells containing the inoculums as well as those containing only the culture media were made and constituted the negative controls and the positive control with the antibiotic. After this incubation time, 40 μl of a 0.2% aqueous para-iodonitrotetrazolium chloride (INT) solution was added to these wells and incubated at 37°C

for 30 min. Thus, wells that turn pink after addition of INT indicate bacterial growth [11]. All concentrations that prevented the appearance of pink color were taken as the inhibitory concentrations and the smallest was scored as Minimal Inhibition Concentration (MIC). For each extract, three columns were made and the revelation was done on two columns.

The third column was used to determine the Minimum Bactericidal Concentrations (MBC). After reading the different MICs, 150 µl of newly prepared Mueller-Hinton Broth was introduced into the wells of the new plates, and then 50 µl of the contents of each well where there was inhibition of bacterial growth (absence of pink staining) was withdrawn with a micropipette and introduced into the corresponding wells of the new plate. These plates were again covered with a sterile lid. Negative control wells, containing only Mueller-Hinton broth and those containing the inoculums without extract or antibiotics were made. The new incubation was also done at 37°C, for 48 hours. The revelation was done as for MIC determination (40 µl of an aqueous solution of INT was added to each well). All extract concentrations for which the absence of bacterial growth was noted (no appearance of pink coloration) were considered as bactericidal concentrations and the smallest was noted as MBC. This test was repeated three (03) times.

2.6. In vitro evaluation of the antioxidant activity of extracts from the leaves and stem bark of *Bauhinia rufescens*

2.6.1. DPPH[•] radical assay

DPPH[•] radical scavenging by the DPPH[•] (1, 1-diphenyl-2-picrylhydrazyl) assay was used to investigate the radical scavenging activities of *Bauhinia rufescens* extracts according to the technique described by Mensor et al. [12]. The extract (2000 µg/ml) was serially diluted twice with methanol. 100 µl of diluted extract was mixed with 900 µl of 0.3 mM DPPH methanol solution, for a final extract concentration of 12.5 to 200 µg/ml (12.5, 25, 50, 100 and 200 µg/ml). The absorbance at 517 nm was measured using a spectrophotometer, after 30 min of incubation at room temperature in the dark. Ascorbic acid (vitamin C) was used as a control.

$$\text{Radical reduction percentage} = \frac{[\text{Absorbance of DPPH} - \text{Absorbance of sample}]}{\text{Absorbance of DPPH}} \times 100$$

The IC₅₀ (amount of sample required to inhibit 50% of the free radical DPPH) was determined by plotting the percentages of radical scavenging activity against the log values of the test sample concentration.

2.6.2. Ferric reduction/antioxidant power assay (FRAP)

The iron (Fe³⁺) reducing power of the extracts was determined according to the method described by Padmaja et al. [13]. For this purpose, one milliliter of extract at different concentrations (200; 100; 50; 25 and 12.5 µg/ml) was mixed with 2.5 ml of 0.2 M phosphate buffer solution (pH 6.6) and 2.5 ml of 1% potassium ferricyanide K₃Fe (CN)₆ solution. The whole set was incubated in a water bath at 50°C for 20 min; then 2.5 ml of 10% trichloroacetic acid was added to stop the reaction and the tubes were centrifuged at 3000 rpm for 10 min. An aliquot (2.5 ml) of supernatant was combined with 2.5 ml of distilled water and 0.5 ml of a 0.1% FeCl₃ ethanol solution. The absorbance of the reaction medium was read at 700 nm against a similarly prepared blank, where the plant extract was replaced with distilled water.

The positive control was represented by a solution of a standard antioxidant (L- ascorbic acid or vitamin C) whose absorbance was measured under the same conditions as the samples. An increase in absorbance corresponds to an increase in the reducing power of the tested extracts [14, 5].

2.6.3. Determination of total phenols

The content of total phenols was determined by the method described by Ramde-Tiendrebeogo et al. [15]. The reagent was a mixture of phosphotungstic acid (H₃PW₁₂O₄₀) and phosphomolybdic acid (H₃PMO₁₂O₄₀). It was reduced upon oxidation of the phenols to a mixture of blue oxides of tungsten and molybdenum. These blue pigments have a maximum absorption depending on the qualitative and or quantitative composition of phenolic mixtures in addition to the pH of the solutions, usually obtained by adding sodium carbonate [16]. The reaction mixture in this assay consisted of 20 µl of extract (2 mg/ml), 20 µl 2N of Folin-Ciocalteu reagent and 40 µl of a 20% sodium carbonate solution. The mixture was stirred and incubated in a water bath at 40 °C for 20 min, and then the absorbance was measured at 760 nm. The extract was replaced with distilled water in the blank tubes. A calibration curve was plotted using gallic acid (0-2 mg/ml); results were expressed as milligrams of gallic acid equivalent per gram of extract (mgGAE/g).

2.6.4. Determination of flavonoid content

The flavonoid content of the extracts was determined using the aluminum trichloride colorimetric method of Padmaja et al. [13]. 100 µl of the extract was mixed with 1.49 ml of

distilled water, and then 30 µl of a 5% sodium nitrite NaNO₂ solution was added. After 5 min, 30 µl of a 10% aluminum chloride solution AlCl₃ was added. The mixture was left to stand for 6 min, and then 200 µl of 1M NaOH solution and 240 µl of distilled water were added. The whole mixture was homogenized with a vortex and the absorbance was measured at 510 nm. The total flavonoid content was calculated using a calibration curve with catechin and the results were expressed as milligrams of catechin equivalent per gram of extract (mgCE/g).

2.7. Phytochemical Screening

The determination of the different classes of potential bioactive compounds present in the extracts of *Bauhinia rufescens* was carried out following the standard method of Harbone [17].

2.8. Statistical Analysis

Statistical analyses were done using SPSS (22.0) for Windows software by ANOVA analysis of variance followed by Waller-Duncan test for comparison between the parameters of the control groups and those of the test groups. All results obtained were expressed as means ± standard deviation. Probability values P < 0.05 were considered significant.

3. RESULTS

3.1. The extraction yield of the different extracts

Table 1 shows the extraction yield of hydroethanolic and aqueous extracts of *Bauhinia rufescens* leaves and stem bark. The extraction yield varies with the solvent used. The yields of hydroethanolic extracts were higher than those of aqueous extracts. Regardless of the plant part, the best extraction yield was obtained with the 95% hydroethanolic solvent system.

3.2. In vitro antisalmonellal activities of Bauhinia rufescens leaf and stem bark extracts

The values of the Minimum Inhibitory Concentrations (MICs) and Minimum Bactericidal Concentrations (MBCs) obtained in the evaluation of the *in vitro* antisalmonellal activity of the extracts of the leaves and stem barks of *Bauhinia rufescens* are represented in Table 2. From this Table, it appears that all the hydroethanolic extracts showed MICs between 256 and 1024 µg/ml. The macerated and decocted leaves showed no activity up to the threshold tested (1024 µg/ml). The 95% hydroethanolic extract of the leaves exhibited a MIC of 512 µg/ml on *Salmonella* Typhi (ST), *Salmonella* Typhi (ATCC6539), *Salmonella* Typhimurium (STM) and *Salmonella* Paratyphi A (SPA). The hydroethanolic extracts (95%, 75% and 50%) and aqueous extracts (decocted, macerated and infused) of stem barks

inhibited the growth of *Salmonella* Typhi, *Salmonella* Paratyphi B, and *Salmonella* Paratyphi A with concentrations ranging from 256 to 512 µg/ml. Regarding the hydroethanolic extracts of leaves and stem barks, MBCs/MICs ratios < 4 were observed on all extracts that presented MBC value.

3.3. Study of in vitro antioxidant activities of extracts from leaves and barks of *Bauhinia rufescens*

3.3.1. Effects of the extracts on the DPPH[•] radical

The free radical scavenging activity of the different extracts from the leaves of *Bauhinia rufescens* was determined after 30 minutes of reaction in the presence of DPPH[•] radical. The ability of the extracts of *Bauhinia rufescens* leaves and vitamin C (reference antioxidant) to inhibit the DPPH[•] radical is presented in Table 3. From this Table, it can be seen that all these extracts possess significant activity against the DPPH[•] radical. The most active extracts are the 95% hydroethanolic extract of leaves and the macerated stem bark which inhibit more than 90% of the DPPH[•] radical at the concentration of 200 µg/ml. All extracts significantly ($p < 0.05$) inhibited DPPH more than vitamin C at all concentrations tested. In general, the activity of L-ascorbic acid was significantly ($p < 0.05$) higher than that of all extracts at all concentration.

3.3.2. IC₅₀ of the extracts of the leaves and stem barks of *Bauhinia rufescens*

The IC₅₀ of the different extracts of the tested leaves and barks of *Bauhinia rufescens* in relation to that of ascorbic acid are presented in Table 4. We find that all the extracts present IC₅₀ values between 9,780 and 19,531 µg/ml. The lowest IC₅₀ value of the extracts was obtained with the 95% hydroethanolic extract of the leaves (IC₅₀ = 9.780±0.125). However, the 50% hydroethanolic extract of the leaves showed the highest IC₅₀ (IC₅₀ = 19,531±2,502). Vitamin C was the most active substance with an IC₅₀ (8,570 mg/ml) significantly lower ($p < 0.05$) than that of all extracts (Table 4).

3.3.3. Iron reducing capacity (FRAP) of *Bauhinia rufescens* leaf and stem bark extracts

The results of iron reducing power of *Bauhinia rufescens* extracts are presented in Table 5. From this table, it can be seen that the 95% hydroethanolic extract of *Bauhinia rufescens* leaves and stem bark showed the highest reducing power ($p < 0.05$) compared to all extracts followed by the 50% hydroethanolic extract of stem bark from the concentration 50 µg/ml. In general, the activity of L-ascorbic acid was significantly ($p < 0.05$) higher than that of all extracts at all concentrations.

3.3.4. Total phenol and flavonoids content of *Bauhinia rufescens*

The analysis of total phenols and flavonoids content revealed that most of these types of metabolites are present in each extract of *Bauhinia rufescens* as presented in Table 6. From this table, it can be seen that the total phenol content of the 95% hydroethanolic extract of the leaves and stem barks were significantly ($p < 0.05$) higher than that all of extracts. However, the flavonoid content of the 50% hydroethanolic extract of leaves and stem bark were significantly ($p < 0.05$) higher than all extracts. The leaf infusions showed the lowest phenol and flavonoid contents.

3.4. Phytochemical study of *Bauhinia rufescens* extracts

The qualitative phytochemical screening of the extracts of the leaves and stem barks of *Bauhinia rufescens* revealed the presence of several classes of secondary metabolites (Table 7). From this table, it appears that tannins, alkaloids, anthraquinones, phenols and flavonoids are present in all extracts tested. All the secondary metabolisms tested are present in the 50%, 75% hydroethanolic extracts and the decoction of the stem barks. The triterpenes and saponins are present in the 95% and 75% hydroethanolic extracts of the leaves; they are also present in the decoction, the infusion and the 50% and 75% hydroethanolic extracts of the stem barks.

Table 1. Extraction yield of leaves and stem bark of *Bauhinia rufescens*.

Plant extracts		Sample mass in (g)		Mass of extracts (g)		Yields %	
		leaves	Barks	leaves	Barks	leaves	Barks
Hydro Ethanolic extracts	95%	250		45.96	39.23	18.387	15.695
	75%			43.83	35.49	17.544	14.196
	50%			41.82	28.45	16.728	11.38
Aqueous extracts	Infused			25.13	25.97	10.052	10.39
	Decocted			27.66	27.50	11.064	11.00
	Macerated			26.25	23.75	10.50	9.50

Table 2. MICs values, MBCs and MBCs/MICs ratios of *Bauhinia rufescens* leaf and stem bark extracts on the four *Salmonella* isolates and *Salmonella Typhi* (ATCC 6539).

Extracts	Concentrations in (µg/ml)	Leaves					Stem bark				
		STS	ST	STM	SPB	SPA	STS	ST	STM	SPB	SPA
95% Hydroethanolic extract	CMI	512	512	512	1024	512	1024	512	512	512	512
	CMB	1024	1024	512	—	1024	1024	1024	512	1024	1024
	CMB/CMI	2	2	1	—	2	1	2	1	2	2
75% Hydroethanolic extract	CMI	1024	1024	—	1024	—	512	512	256	512	256
	CMB	—	1024	—	—	—	512	1024	512	512	512
	CMB/CMI	—	1	—	—	—	1	2	2	1	2
50% Hydroethanolic extract	CMI	1024	1024	1024	1024	1024	512	512	1024	512	512
	CMB	1024	1024	—	—	—	—	—	—	512	1024
	CMB/CMI	1	1	—	—	—	—	—	—	1	2
infused	CMI	1024	1024	1024	1024	—	1024	512	512	512	256
	CMB	—	1024	—	—	—	—	—	1024	—	1024
	CMB/CMI	—	1	—	—	—	—	—	2	—	4
Decocted	CMI	—	—	—	—	—	1024	256	512	512	254
	CMB	—	—	—	—	—	—	1024	—	512	512
	CMB/CMI	—	—	—	—	—	—	4	—	1	2
Macerated	CMI	—	—	—	—	—	—	—	—	512	512
	CMB	—	—	—	—	—	—	—	—	1024	1024
	CMB/CMI	—	—	—	—	—	—	—	—	2	2
Ciprofloxacin	CMI	0,5	0,5	1	0,25	2	0,5	0,5	1	0,25	2
	CMB	2	2	4	1	4	2	2	4	1	4
	CMB/CMI	4	4	4	4	2	4	4	4	4	2

STs: *Salmonella Typhi* strain (ATCC 6539), ST: *Salmonella Typhi*, STM: *Salmonella Typhimurium*, SPB: *Salmonella Paratyphi B* and SPA: *Salmonella Paratyphi A*.
MIC= Minimum Inhibitory Concentration. MBC= Minimum Bactericidal Concentration, the line (-) = No concentration.

Table 3. Percentage of DPPH- free radical inhibition for hydroethanolic and aqueous extracts of *Bauhinia rufescens* leaves and stem bark

Extracts	Concentrations ($\mu\text{g} / \text{ml}$)	Percentage of inhibition \pm Standard deviation						
		EtOH 95%	EtOH 75%	EtOH 50%	Decocted	Macerated	Infused	Vit C
Leaves	12.5	82.007 \pm 3.8249 ^{cd}	63.620 \pm 4.920 ^b	39.425 \pm 11.792 ^a	79.554 \pm 3.899 ^c	82.788 \pm 11.063 ^{cd}	71.084 \pm 12.698 ^{bc}	95.734 \pm 0.223 ^d
	25	89.713 \pm 0.508 ^{ef}	86.701 \pm 6.356 ^{de}	37.551 \pm 5.121 ^a	88.517 \pm 0.365 ^{de}	86.689 \pm 1.106 ^{de}	82.271 \pm 4.846 ^{de}	96.666 \pm 0.654 ^f
	50	91.612 \pm 0.215 ^{ef}	87.639 \pm 1.706 ^{cde}	85.940 \pm 2.811 ^{bcde}	84.358 \pm 6.080 ^{bcd}	84.222 \pm 1.146 ^{bcd}	82.386 \pm 3.963 ^{bc}	96.738 \pm 0.062 ^f
	100	93.010 \pm 1.516 ^{fgh}	91.505 \pm 1.777 ^{efgh}	76.332 \pm 1.196 ^a	90.099 \pm 1.473 ^{efg}	87.378 \pm 0.358 ^{cde}	86.058 \pm 0.172 ^{cd}	96.845 \pm 0.124 ⁱ
	200	94.516 \pm 0.284 ^{cd}	95.957 \pm 0.878 ^{cd}	88.810 \pm 1.330 ^{ab}	84.534 \pm 6.696 ^a	86.861 \pm 1.599 ^a	94.664 \pm 1.204 ^{cd}	97.276 \pm 0.328 ^d
Stem bark	12.5	82.776 \pm 3.801 ^{cd}	39.127 \pm 7.831 ^b	72.461 \pm 1.641 ^{bc}	72.544 \pm 4.737 ^{bc}	78.888 \pm 7.862 ^c	46.559 \pm 4.190 ^a	95.734 \pm 0.223 ^d
	25	86.291 \pm 1.522 ^{de}	67.756 \pm 8.504 ^c	79.403 \pm 7.612 ^d	81.505 \pm 3.182 ^{de}	86.881 \pm 0.654 ^{de}	56.236 \pm 0.558 ^b	96.666 \pm 0.654 ^f
	50	85.3544 \pm 2.661 ^{bcd}	80.091 \pm 3.184 ^{ab}	85.197 \pm 5.844 ^{bcd}	89.569 \pm 0.107 ^{de}	89.856 \pm 0.407 ^{de}	75.304 \pm 1.894 ^a	96.738 \pm 0.062 ^f
	100	90.158 \pm 2.324 ^{ef}	84.509 \pm 1.984 ^{bc}	88.755 \pm 3.791 ^{def}	92.222 \pm 0.345 ^f	94.767 \pm 0.620 ^{hi}	81.254 \pm 2.740 ^b	96.845 \pm 0.124 ⁱ
	200	93.555 \pm 1.129 ^{cd}	91.623 \pm 1.037 ^{bc}	93.631 \pm 2.257 ^{cd}	97.096 \pm 0.492 ^d	96.523 \pm 0.124 ^d	84.623 \pm 1.182 ^a	97.276 \pm 0.328 ^d

EtOH: Ethanol; a, b,c.... : The numbers with different letters are significantly different at the 5% level ($p < 0.05$). The underlying values are of the form of means \pm Standard deviation.

Table 4. Antiradical activities of *Bauhinia rufescens* extracts expressed in term of IC₅₀.

Extracts	IC ₅₀ in µg / ml	
	Leaves	Stem bark
EtOH 95%	9.780±0.125 ^{ab}	10.469±0.051 ^b
EtOH 75%	10.754±0.331 ^b	14.940±0.754 ^c
EtOH 50%	19.531±2.502 ^c	11.347±0.866 ^b
Decocted	11.285±0.482 ^b	10.414±0.049 ^b
Macerated	11.255±0.383 ^b	9.845±0.220 ^{ab}
Infused	11.575±1.175 ^b	17.701±0.798 ^d
Vitamin C	8.570±0.016 ^a	

EtOH = Ethanol; The numbers bearing the letters a, b, c.... are significantly different at the p<0.05.

UNDER PEER REVIEW

Table 5. Iron reducing power (FRAP) of hydroethanolic and aqueous extracts of leaves and stem bark of *Bauhinia rufescens*.

Extracts	Concentration (µg/ml)	Optical density ± Standard deviation						
		EtOH 95%	EtOH 75%	EtOH 50%	Decocted	Macerated	Infused	VitC
Leaves	12.5	0.180±0.002 ^{abc}	0.142±0.019 ^a	0.197±0.009 ^{bc}	0.178±0.020 ^{ab}	0.132±0.041 ^a	0.176±0.001 ^{bc}	0.204±0.006 ^{bc}
	25	0.467±0.050 ^e	0.223±0.023 ^a	0.226±0.002 ^a	0.228±0.013 ^a	0.229±0.008 ^a	0.240±0.021 ^a	0.452±0.025 ^{de}
	50	0.686±0.028 ^{gh}	0.353±0.007 ^{cd}	0.299±0.013 ^{ab}	0.279±0.033 ^a	0.315±0.018 ^{abc}	0.410±0.002 ^{ef}	0.992±0.026 ⁱ
	100	0.763±0.016 ^d	0.472±0.014 ^{ab}	0.745±0.000 ^c	0.424±0.019 ^a	0.508±0.066 ^{ab}	0.444±0.004 ^{ab}	1.813±0.005 ^e
	200	0.938±0.002 ^{cd}	0.810±0.026 ^b	0.857±0.025 ^{bc}	0.802±0.052 ^b	0.781±0.009 ^b	0.663±0.016 ^a	1.829±0.003 ^e
Stem bark	12.5	0.230±0.007 ^{cde}	0.271±0.014 ^{de}	0.280±0.021 ^e	0.265±0.045 ^{de}	0.224±0.010 ^{bcd}	0.193±0.008 ^{bc}	0.204±0.006 ^{bc}
	25	0.286±0.003 ^{ab}	0.345±0.070 ^{bc}	0.453±0.018 ^{de}	0.392±0.009 ^{cd}	0.371±0.022 ^c	0.221±0.029 ^a	0.452±0.025 ^{bc}
	50	0.395±0.017 ^{de}	0.642±0.009 ^{fg}	0.784±0.014 ^h	0.446±0.003 ^{fg}	0.652±0.037 ^g	0.333±0.015 ^{bc}	0.992±0.026 ⁱ
	100	0.855±0.072 ^{cd}	0.762±0.126 ^c	0.920±0.067 ^d	0.859±0.043 ^{cd}	0.970±0.012 ^d	0.554±0.001 ^b	1.813±0.005 ^e
	200	1.004±0.022 ^d	0.939±0.013 ^{cd}	0.967±0.017 ^d	0.926±0.092 ^b	0.965±0.026 ^d	0.804±0.067 ^b	1.829±0.003 ^e

ETOH = Ethanol; VitC = Vitamin C; The numbers bearing the letters a, b, c.... are significantly different at p <0.05.

Table 6. Total phenol and flavonoid content of hydroethanolic and aqueous extracts of *Bauhinia rufescens* leaves and stem bark.

Extracts		Concentrations of total phenols (mgGAE/g) ± Standard deviation	Concentrations of total flavonoids (mgCE/g) ± Standard deviation
Leaves	EtOH 95%	2.895±0.081 ^d	2.335±0.265 ^{ab}
	EtOH 75%	1.215±0.098 ^a	2.505±0.605 ^{ab}
	EtOH 50%	2.081±0.488 ^c	2.590±0.469 ^{ab}
	Decocted	1.203±0.168 ^b ^a	2.717±0.447 ^{ab}
	Macerated	1.377±0.005 ^{ab}	2.335±0.383 ^{ab}
	Infused	0.994±0.145 ^a	1.337±0.220 ^a
Stem bark	EtOH 95%	4.000±0.011 ^f	2.165±0.168 ^a
	EtOH 75%	3.500±0.058 ^{ef}	2.462±0.715 ^{ab}
	EtOH 50%	3.075±0.180 ^{de}	4.225±0.574 ^b
	Decocted	1.779±0.453 ^{bc}	1.231±0.447 ^a
	Macerated	0.877±0.122 ^a	2.760±0.669 ^{ab}
	Infused	1.232±0.533 ^a	2.144±0.715 ^a

EtOH = Ethanol; The numbers bearing the letters a, b, c.... are significantly different at p < 0.05.

UNDER PEEL

Table 7. Phytochemical composition of leaf and bark extracts of *Bauhinia rufescens*.

Extracts	Chemical classes	Hydroethanolic extracts			Aqueous extracts		
		95%	75%	50%	Decocted	infused	Macerated
Leaves	Alkaloids	+	+	+	+	+	+
	Phenols	+	+	+	+	+	+
	Flavonoids	+	+	+	+	+	+
	Anthocyanins	-	-	-	-	-	-
	Anthraquinones	+	+	+	+	+	+
	Tannins	+	+	+	+	+	+
	Steroids	+	+	-	-	-	-
	Triterpenes	+	+	-	-	-	-
	Saponins	+	+	-	-	-	-
Stem barks	Alkaloids	+	+	+	+	+	+
	Phenols	+	+	+	+	+	+
	Flavonoids	+	+	+	+	+	+
	Anthocyanins	-	+	+	+	-	+
	Anthraquinones	+	+	+	+	+	+
	Tannins	+	+	+	+	+	+
	Steroids	-	+	+	+	+	-
	Triterpenes	-	+	+	+	+	-
	Saponins	-	+	+	+	+	-

-: Absence; +: presence

4. DISCUSSION

The hydroethanolic extracts (95%, 75% and 50%) showed the highest yields compared to the aqueous extracts. In addition, the addition of water to ethanol results in alcohol dilution and reduces the yield instead. These results are in agreement with those of Mohammedi and Atik [18] who revealed that mixed solvents are very efficient to extract. In addition, the yield of the decoction was higher than that of the other aqueous extracts. The superiority of this yield over the maceration and infusion suggests that temperature would play a positive effect, improving extraction [19]. The results of antimicrobial tests (MIC and MBC) on the leaves and stem barks of *Bauhinia rufescens* show that they contain substances with antisalmonellal activity. Indeed, many plant extracts and isolated compounds from the Fabaceae family have been presented as possessing antibacterial activity [20]. Hydroethanolic and aqueous extracts of *Bauhinia rufescens* leaves and stem bark exhibited significant to low Minimum Inhibitory Concentrations (MICs) (256 to 1024 µg/ml).

According to Kuete [21], the antibacterial activity of plant extracts is considered significant when $MIC < 100 \mu\text{g/ml}$, moderate when $100 \mu\text{g/ml} \leq MIC \leq 625 \mu\text{g/ml}$ and low when $MIC > 625 \mu\text{g/ml}$. The 95% hydroethanolic extract of the leaves exhibited the moderate activity on three isolates and strain of *Salmonella* (*Salmonella* Typhi, *Salmonella* Paratyphi A, *Salmonella* Typhimurium and *Salmonella* Typhi ATCC6539) with concentrations of 512

µg/ml. The hydroethanolic extracts and the aqueous extract of the stem bark exhibited moderate activity on isolates (*Salmonella* Typhi, *Salmonella* Typhi B and *Salmonella* Paratyphi A) with concentrations ranging from 256 to 512 µg/ml. These results corroborate those of Muhammad and Sirat [17] who showed that the methanolic extract of this plant inhibited the growth of some pathogenic bacteria like *P. aeruginosa*. The differences in inhibitory activities observed with the same extract towards the different isolates and the *Salmonella* strain could be due to the difference in molecular structure between the tested pathogens [22, 5].

The null activity observed with the aqueous extracts (macerated and decocted) of the leaves could be related to the absence of anthocyanins and saponins in these extracts. The inhibition of *Salmonella* growth by the different extracts of *Bauhinia rufescens* may be due to the presence of phenolic compounds. This result corroborates the work of Sokoudjou et al. [23] who showed that phenolic compounds such as gallic acid and scopoletin were active (MIC ranging from 16 to 128 µg/ml) against ST, STM and *S. Typhi* 6539. Antibacterial substances can be classified as bactericidal, when the BMC/MIC ratio ≤ 4 or bacteriostatic, when the MBC/MIC ratio > 4 [24]. Based on these criteria, the majority of the hydroethanolic extracts were bactericidal. The solvent system and the extraction methods used influenced the different results obtained. Indeed, several works have already shown that the solvent system used for an extraction can strongly influence its secondary metabolite content [25, 5, 26] and that the method of extraction of plant extracts can influence their activities [27, 28].

The difference in the observed activity between the different extracts on the one hand and the isolates on the other hand may be due either to the constitutional or structural variability of the germs tested, or to the difference in solubility of the active substances present in each extract. It could also be due to the difference in the chemical composition of the genetic elements of transferable resistance between strains [24], or to the difference in the composition of secondary metabolites found in each extract.

It is likely that their antimicrobial activity is not attributable to a single mechanism, but their simultaneous action at different bacterial sites. This corroborates the work of Kobanski et al; Kodjio [29, 30] and Sokoudjou et al. [31] who suggest that some bacteriostatic or bactericidal substances act by interfering with some essential structures of the bacteria such as the membrane wall, genome and proteins. The results of antimicrobial tests (MIC and MBC) of the leaves and stem bark of *Bauhinia rufescens* showed that this plant contains substances with antisalmonellal activity that could be used in the treatment of typhoid and paratyphoid fevers.

Antioxidant compounds have various mechanisms of action and a single method would not be sufficient to evaluate the total capacity of an antioxidant [32, 5]; for this reason, several approaches were used to evaluate the antioxidant activity of the different extracts of the leaves and stem bark of *Bauhinia rufescens*. The results of this study showed that the 95% hydroethanolic extract of leaves and the macerated stem bark exhibited higher antiradical activity than all extracts at the concentration of 100 µg/ml. The antioxidant activity of the extracts could be explained by their richness in polyphenolic substances, tannins and more particularly in total flavonoids. These results are in agreement with those of several authors, who reported a positive correlation between all phenolic content and antioxidant activity [33, 5]. Hydroethanolic and aqueous extracts of *Bauhinia rufescens* leaves and stem barks showed IC₅₀ values between 9.780 and 19.531 µg/ml. These results corroborate those of Promprom and Chatan [34] who showed that the ethanolic leaf extract of *Bauhinia nakhonphanomensis* presented an IC₅₀ value of 17.07 ± 0.24 µg/ml. According to Soury et al. [35] the antioxidant potential of a plant is divided into three groups: high when IC₅₀ < 20 µg/ml, moderate when 20 µg/ml ≤ IC₅₀ ≤ 75 µg/ml and low when IC₅₀ > 75 µg/ml. The IC₅₀ values showed that the hydroethanolic and aqueous extracts of the leaves and stem barks of *Bauhinia rufescens* exhibited high antiradical activities because all these extracts have IC₅₀ < 20 µg/ml. The high antiradical activity of the different extracts of the leaves and stem barks of *Bauhinia rufescens* could be explained by the high presence of polyphenolic compounds (total phenols, anthraquinones, and flavonoids). The antioxidant activity of other secondary metabolites is directly related to their hydroxyl group.

The iron reduction test is one of the methods for assessing antioxidant activity. The 95% hydroethanolic extract of the leaves showed the highest reducing power (p<0.05) compared to all extracts followed by the 50% hydroethanolic extract of the stem bark from the concentration 50 µg/ml. The antioxidant potential of *Bauhinia rufescens* leaf and bark extracts could be related to the presence of total phenol and flavonoids which were detected during the quantitative assay. Indeed, total phenols and flavonoids are powerful antioxidants [34]. These results suggest the reducing power of *Bauhinia rufescens* is likely due to the presence of hydroxyl group in phenolic compounds that can serve as proton donors. Therefore, antioxidants are considered to be reductants and inactivators of oxidants [36].

Phenolic compounds are important plant constituents, as they act against lipid peroxidation and inactivate free radicals [15]. The concentration of phenolic compounds is very high in the 95% hydroethanolic extract of leaves and stem bark compared to other extracts. In addition, flavonoids are powerful antioxidants, which possess redox properties that allow them to

eliminate the effects of reactive oxygen species [37] as well as to chelate various transition metals [15]. These results corroborate those of the iron reduction test, where the 95% hydroethanolic extract of leaves and stem barks exhibited the greatest antioxidant activity. There was a positive correlation between the antioxidant activity and the phenol content in the extracts. Many authors have also shown this correlation [16, 38]. These results suggest that the extracts of *Bauhinia rufescens* have an antioxidant activity, due to the phenolic compounds present in these extracts.

5. CONCLUSION

From the results of this work, it can be concluded that the extracts of the plant *Bauhinia rufescens* have antisalmonellal and antioxidant activity *in vitro*. The hydroethanolic extracts of this plant inhibited all isolates and strain of *Salmonella* Typhi (ATCC6539) with Minimum Inhibitory Concentrations ranging from 256 to 1024 µg/ml. The hydroethanolic and aqueous extracts presented the best antiradical activities with IC₅₀ < 20µg/ml. The results demonstrate that the 95% hydroethanolic extract of *Bauhinia rufescens* leaves has both antisalmonellal and antioxidant activity *in vitro*, due to the presence of phytochemical free radical scavenging compounds that could have the ability to inhibit a free radical and consequently could reduce oxidative stress. These results support the claim of the Chadian population to use *Bauhinia rufescens* for the treatment of typhoid fever. However, further studies are needed to determine the *in vivo* antisalmonellal and antioxidant activities, side effects and to define the therapeutic dose that would allow safe use.

Data Availability

No additional data are available.

REFERENCES

1. WHO. Surveillance de la fièvre typhoïde et utilisation des vaccins contre cette maladie, Régions de l'Asie du Sud-Est et du Pacifique occidental, 2009-2013. *Relevé Épidémiologique Hebdomadaire*. 2014 ; 89: 429-440.
2. WHO. Fabrication de comprimés et de solutions orales à base de zinc Directives à l'intention des responsables de programmes et de l'industrie pharmaceutique. Organisation Mondiale de la Santé : Genève, Suisse. 2008 ; 30.
3. Shukla P, Bansode WF, Singh KR. Chloramphenicol toxicity: A Review. *Journal of Medicine and Medical Sciences*. 2011; 2(13): 1313-1316.
4. Murphy EA, Davis JM, Brown AS, Carmichael MD, Carson JA, Van Rooijen N, Ghaffar A, Mayer EP. Benefits of oat-glucan on respiratory infection following

- exercise stress: role of lung macrophages. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. (2007); 10: 1-30.
5. Kodjio N, Atsafack SS, Njateng SSG, Sokoudjou BJ, Kuate RJ, Gatsing D. Antioxidant effect of aqueous extract of *Curcuma longa* Rhizomes (Zingiberaceae) in the typhoid fever induced in Wistar Rats Model. *Journal of Advances in Medical and Pharmaceutical*. 2016 ; 7(3): 1-13.
 6. Rastaldo R, Pagliaro P, Cappello S, Penna C, Mancardi D, Westerhof N, Losano G. Nitricoxide and cardiac function. *Life Sciences*. 2007; 81: 779-793.
 7. Da silva KLD, Filho VC. Plants of the genus Bauhinia: chemical composition and pharmacological potential. *Química Nova*. 2002; 25: 449-454.
 8. Abubakar MS, Musa AM, Ahmed A, Hussaini IM. The perception and practice of traditional medicine in the treatment of cancers and inflammations by the hausa and fulani tribes of northern nigeria. *Journal Ethnopharmacol*. 2007; 111(3): 625-629.
 9. Kamsu GT, Simo Tagne R, Fodouop SPC, Famen NLC, kodjio N, Ekom ES, Gatsing D, “In vitro antisalmonellal and antioxidant activities of leaves extracts of *Tectonagrandis*L. F. (Verbenaceae),” *European Journal of Medicinal Plants*. 2019a; vol. 29, no. 4, pp. 1-13.
 10. Goly C, Soro Y, Kassi B, Dadié A, Soro S, Marcellin DJE. Antifungal activities of the essential oil extracted from the tea of savanna (*Lippia multiflora*) in Côte d’Ivoire. *Intenational Journal Biological and Chemical Science*. 2015; 9(1): 24-34
 11. Mativandlela SPN, Lall N, Meyer JJM. Antimicrobial, antifungal, antitubercular activity of *Pelargonium reniforme* (CURT) and *Pelargonium sidoides* (DC) (Geraniaceae) root extracts. *South Africa Journal of Botanic*. 2006; 72: 232-237.
 12. Mensor LL, Menezes FS, Leitao GG, Reis ASO, Dos Santos TC, Coube CS, Leitao SG. Screening of Brazilian plant extracts for antioxidant activity by the used DPPH free radical method. *Phytotherapy Research*. 2001; 15: 127-130.
 13. Padmaja M, Sravanthi M, Hemalatha KPJ. Evaluation of Antioxidant Activity of Two Indian Medicinal Plants. *Journal of Phytology*. 2011; 3(3): 86-91.
 14. Mohammed AI, Neil AK, Shahidul IM. *In vitro* anti-oxidative activities and gc-ms analysis of various solvent extracts of *cassia singueana* parts. *Drug Research*. 2013; 70(4): 709-719.
 15. Ramde-Tiendrebeogo A, Tibiri A, Hilou A, Lompo M, Millogo-Kone H, Nacoulma OG, Guissou IP. Antioxidative and antibacterial activities of phenolic compounds

- from *Ficus sue Forssk. International Journal of Biological and Chemical Sciences*. 2012; 6(1): 328-336.
16. Luis A, Domingues F, Gil C, Duarte AP. Antioxidant activity of extracts of Portuguese shrubs: *Pterospartum tridentatum*, *Cystisus scoparius* and *Erica spp.* *Journal of Medicinal Plants Research*. 2009; 3(11): 886-893.
 17. Harbone JB. *Phytochemical methods: a guide to modern techniques of plants analysis. Chapman and Hall Ltd: London*. 1973; 50-116.
 18. Mohammedi Z, Atik F. Impact of solvent extraction type on total polyphenols content and biological activity from *Tamarixaphylla(L.) karst.* *International Journal Pharmacology Biology Science*. 2011 ; 2: 609-615.
 19. Bourgou S, Serairi BR, Medini F, Ksouri R. Effet du solvant et de la méthode d'extraction sur la teneur en composés phénoliques et les potentialités antioxydantes d'*Euphorbiahelioscopia*. *Journal of new sciences, Agriculture and Biotechnology*. 2016; 28(12): 1649-1655.
 20. Muhammad A, Sirat MH. Antimicrobial, antityrosinase and brine shrimp lethality test of *Bauhinia rufescens* (Fabaceae). *Journal of Coastal Life Medicine*. 2013; 1(2): 135-140.
 21. Kuete V. Potential of Cameroonian plants and derived products against microbial infections. *Planta Medica*. 2010; 76: 1-13.
 22. Tene M, Tane P, Tamokou JD, Kuate JR, Connolly JD. Antimicrobial diterpenoids and triterpenoids from the stem bark of croton macrostachys. *International Journal of Biological and Chemical Science*. 2009; 3(3): 538-544.
 23. Sokoudjou JB, Atolnani O, Njateng GSS, Khan A, Tagousop NC, Bitombo AN, Kodjio N, Gatsing D. Isolation, characterization and *in vitro* anti-salmonella activity of compounds from stem bark extract of *Canariumschweinfurthii*. *MBC Complementary Medicine and Therapies*. 2020; 20 (316): 1-10.
 24. Gatsing D, Adoga GI. Antisalmonella activity and phytochemical screening of the various parts of *Cassia petersiana Bolle* (Caesalpinaceae). *Research Journal of Microbiology*. 2007 ; 2(11): 876-880.
 25. Dohou N, Yamni K, Badoc A, Douira A. Activité antifongique d'extraits de *Thymelaelythroïdes* sur trois champignons pathogènes du riz. *Société de pharmacie de Bordeaux*. 2004 ; 143: 31-38.
 26. Sokoudjou JB, Njateng GSS, Fodouop SPG, Kodjio N, Atsafack SS, Fowa AB, Djimeli MN, Gatsing D. *In vitro* antisalmonella and antioxidant activities of

- Canariumschweinfurthii* stem bark extracts. *Academia Journal of Medicinal plants*. 2018; 6 (10): 331-341.
27. Frederico SM, Leonardo LB, José RP, Edemilson CC. Impact of different extraction methods on the quality of *Dipteryxalata* extracts. *RevistaBrasileira de Farmacognosia Brazilian Journal of Pharmacognosy*. 2013; 23(3): 521-526.
 28. Abdelfadel MM, Khalaf HH, Sharoba AM, Assous MTM. Effect of extraction methods on antioxydant and activities of some *Spices and Herbs* extracts. *Journal of Food Technology and Nutritional Sciences*. 2016; 1(1): 1-14
 29. Kohanski MA, Dwyer DJ, Collins JJ. How antibiotics kill bacteria: from targets to networks. *Nature Reviews Microbiology*. 2010 ; 8: 423-435.
 30. Kodjio N. Eude des propriétés antisalmonelles, antioxydantes, toxicologiques et formulations galénique des extraits de *Curcuma longa L*. Thèse de Pharmacologie, Département de Biochimie, Université de Dschang-Cameroun. 2017.
 31. Sokoudjou JB. Propriétésantisalmonelles, antioxydantes et étude de la toxicité des extraits d'écorces de *Canariumschweinfurthii*Engl. (Burseraceae) chez le poulet de chair. Thèse de DoctoraleFacultes des Sciences, Ecole Doctorale des Sciences Fondamentales et Technologiques. 2021; 233p.
 32. Huang D, Ou B, Prior RL. The chemistry behind antioxidant capacity assays. *Journal of Agricultural and Food Chemistry*. 2005; 53(6):1841-1856.
 33. Noghogne RL, Gatsing D, Fotso, Kodjio N, Sokoudjou BJ, Kuate RJ. *In vitro* antisalmonellal and antioxidant properties of *Mangifera indica L*. Stem bark crude extracts and fractions. *British Journal of Pharmaceutical Research*. 2015; 5(1): 29-41.
 34. Promprom W, Chatan W. GC-MS analysis and antioxidant activity of *Bauhinia nakhonphanomensis* leaf ethanolic extract. *Pharmacognosie Journal*. 2017; 9(5): 663-667.
 35. Souri E, Amin G, Farsam H, Barazandeh TM. Screening of antioxidant activity and phenolic content of 24 medicinal plants. *Journal of Pharmaceutical Sciences*. 2008; 16(2): 83-87.
 36. Siddhuraju P, Becker K. The antioxidant and free radical scavenging activities of processed cowpea (*Vigna unguiculata (L.) Walp.*) seed extracts. *Food Chemistry*. 2007; 101(1), 10-19.
 37. Ketsawatsakul U, Whiteman M, Halliwell B. A reevaluation of the peroxynitrite scavenging activity of some dietary phenolics. *Biochemical and Biophysical Research Communications*. 2000; 279: 692-699.

38. Tawaha K, Alali F, Gharaibeh M, El- Elimat T. *Antioxidant activity and total phenolic content of selected Jordanian species*. Food Chemistry. 2007; 104: 1372-1378.

UNDER PEER REVIEW