

Original Research Article

Evaluation of Some Haematologic Parameters among Cement Loaders in Port-Harcourt

ABSTRACT

Cement remains a very useful discovery in the modern world and has served many purposes in the circle of construction, the dusts generated in the manufacturing, packaging, transporting, and also usage has impacted negatively on those exposed to the dusts at various levels. This study focuses on assessing the health impact of cement dust on haematologic parameters among cement loaders in Port Harcourt. A cross sectional study design was employed for this study. A total 200 male subjects; 100 healthy cement exposed workers and 100 apparently healthy non exposed male subjects. Subjects were selected randomly based on set inclusion and exclusion criteria. Venous blood samples were collected using venipuncture technique into EDTA bottle for the assay of Erythrocyte Sedimentation Rate (ESR) and Full Blood Count (FBC). ESR was estimated using Westergreen method while FBC was assayed using Haematology analyzer. There was a significant decrease in WBC count, RBC count, haemoglobin (Hb) level and monocyte in cement workers (p -value <0.05) but there was a significant increase in ESR level in cement workers (p -value <0.05). This study has shown that there cement dust exposure results in inflammation and can as well interfere with red blood cell production process.

Keyword: *cement, haematological parameters, exposure*

1.0 INTRODUCTION

Cement is a very fine substance with adhesive and cohesive qualities that acts as glue for the individual elements. There are over ten different varieties of cement used in construction, each with a different composition and made for particular purposes (Dunuweera and Rajapakse, 2018). These include rapid-hardening cement (RHC), quick-setting cement (QSC), low-heat cement (LHC), sulphate-resistant cement (SRC), blastfurnace slag cement (BFSC), high-alumina cement (HAC), white cement (WC), coloured cement (CC), pozzolanic cement (PzC), air-entraining cement (AEC), and hydrophobic cement (HPC). Cements are generally useful in the production of concrete materials, bridges, houses, culverts, and as well as in construction works (Joel and Mbapuan, 2016). Molecules of principal value in cement essentially include 60-67% calcium oxide, 17-25 silicon oxide (SiO_2), 3-5% aluminium (Al) oxide, with some amount of iron oxide, chromium (Cr), potassium, sodium, sulphur and magnesium oxide (Kakooei *et al.*, 2012; Gbadebo and Bankole, 2007). Portland cement is the most used type of cement, and it has a large amount of lime when it comes to rapid-hardening cement (Dunuweera and Rajapakse, 2018). The composition of cement includes; Lime (CaO), Silica (SiO_2), Alumina (Al_2O_3), Calcium Sulphate (CaSO_4), Iron Oxide (Fe_2O_3), Magnesia (MgO), Sulphur (S), and Alkalies. The most extensively used cement for constructions and building is the Ordinary Portland Cement (OPC). It is made through a procedure known as hydration by crushing, grinding, and combining particular proportions of lime or calcium Oxide (CaO : from limestone, chalk, shells, shale or calcareous rock), iron (Fe_2O_3 : from clay, iron ore, scrap iron and fly ash), silica (SiO_2 : from sand, old bottles, clay or argillaceous rock), alumina (Al_2O_3 : from bauxite,

recycled aluminum, clay), and Gypsum, $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$: found together with limestone (Saleh and Rahman, 2018). The chemical interactions between the cement and the water give Portland cement its strength. This is a complicated process that is best understood by first learning about cement's chemical makeup (Saleh and Rahman, 2018). Molecules of principal value in cement essentially include 60-67% calcium oxide, 17-25 silicon oxide (SiO_2), 3-5% aluminium (Al) oxide, with some amount of iron oxide, chromium (Cr), potassium, sodium, sulphur and magnesium oxide (Kakoei *et al.*, 2012; Gbadebo and Bankole, 2007). Portland cement, which is one of the mostly used cements, comprises a mixture of calcium oxide (60–67%), silicon dioxide (17–25%), aluminium trioxide (3–8%) and ferric oxide (0–5%) (Azah *et al.*, 2002). Some of these elements exist freely and abundantly in nature. An example is Aluminium which it has no recognized critical role in biological systems, but has been shown to pose a threat to the health of both humans and animals. Other elements found in cement: iron, chromium, calcium, and silica have all been shown to pose a health hazard to those exposed to them. Different types of cells exist in the human blood, and these cells serve various activities (Koury *et al.*, 2009). These blood cells are produced in the haemopoietic tissue (Gordon, 2005). The most significant cells in the formation of blood cells are stem cells. Gordon (2005) found that soluble factors regulate haemopoiesis, which follows a hierarchical structure in which multipotent stem cells give rise to progenitor cells, which then divide to produce adult blood cells (Gordon, 2005).

Exposure to dust causes alteration in blood cell counts. This can be due to allergic reactions and inflammation which in turn evoked by dust entering the lungs. This has been observed even from dust generated from Rice mill (Patil *et al.*, 2015). A study showed that neutrophil, eosinophil and lymphocyte count among haematological indices were significantly increased in exposed individuals compared with control group. (Patil *et al.*, 2015) thus posit that high level of dust exposure has deleterious effects on blood. Discoveries based on a research by Rahman, (2018) finalized that the exposure to cement components as well as inhalation leads to problems. The haematological indices that have been implicated in cement exposure are ESR, WBC, RBC, Hb levels. These parameters are raised due to exposure to cement particles thereby suggesting, increased activation of allergic response (John and Olubayo, 2011). Also, reports have shown the protective role of haematopoietic products in prevention of oxidative stress related diseases (Feng *et al.*, 2005; Gawad *et al.*, 2009). Specifically, Gawad *et al.* (2009) demonstrated the protective role of erythropoietin against apoptosis induced oxygen radicals. Furthermore, cement dust exposures has been reported to result in significant occupational health problems and long term complications and symptoms. Another study reported that occupational exposure to cement dust may have a negative impact on the haemopoietic function with emphasis on the provision and use of appropriate personal protective equipment.

2.0 MATERIALS AND METHOD

2.1 Study Design

A cross sectional study design was employed for this study. A convenient sampling size of 200 subjects was used. This research work was carried out on 200 male subjects; of which 100 were cement exposed healthy subjects working in one cement depot and eight loading sites and 100 were apparently healthy non exposed male subjects. The exposed subjects were selected randomly. The medical history as well as the bio-data, socio demographics, environmental and lifestyle of the study subjects were obtained with the use of a well-structured questionnaire.

Comment [CC1]: How this sample size was calculated, needs elaboration with reference

2.2 Study Area

The study was carried out in Port Harcourt metropolis, Rivers State, Nigeria. Port Harcourt is the capital and biggest city of Rivers State, Nigeria with its geographic coordinates as latitude: 4°46'38" N, longitude: 7°00'48" E and elevation above sea level: 16 m = 52 ft. It lies along the Bonny Stream and is situated in the Niger Delta. The Port Harcourt urban Harcourt territory has an expected populace of 1,865,000 occupants, up from 1,382,592 starting at 2006 (Weli and Efe, 2015). Despite the change of weather and climate being the influential factors that impact cement consumption, cement still remains a construction raw material, regularly distributed across various parts of Port Harcourt city (Amadi and Dimkpa, 2020). Cement regardless of the economic situation, imparts on the country's gross local product (Mojekwu *et al.*, 2013). The study sites for this research were Dangote cement depot Trans-Amadi and Cement loading points/shops across Port Harcourt Metropolis ranging from Borokiri (latitude: 4.7463, longitude: 7.0364), Choba (latitude 6.50' 00''E; longitude: 007.00' 00E, Eleme junction (latitude: 4°51'21.6"N, longitude: 7°04'00.4"E), Eneka (latitude: 4°53'59.0"N (4.8997100°) and latitude: 7°01'59.9"E (7.0333100°)), Mile 3 Diobu (latitude: 4° 47' 57" North, longitude: 7° 0' 34), Rukpokwu (latitude: 4.902909, latitude: 7.001263), Rumuokoro (latitude: 4° 52' 3" North, longitude: 6° 59' 53" East), and Woji (longitude: 4° 47' 18" North, longitude: 7° 0' 3" East) all in Port Harcourt, Rivers State, Nigeria (Wekpe and Fiberesima, 2020). Dangote cement depot is situated at RIVOC, Trans -Amadi in Port Harcourt, Rivers State.

Comment [CC2]: How did you confirm that The sample selected from this area was exposed to pollutants caused by cement ad not y any other source?
Did you exclude that there was other source of exposure to the said chemicals and the cement was the only soure. Elaborate in detail please

2.3 Study Population

This study involved 100 cement loaders who worked in cement depots and shops, and another 100 control subjects who were not involved in the use of cement occupationally. The biodata of the subjects were obtained using questionnaire. Individuals involved in any occupation that involves the use of cement, were not recruited as control subjects. Informed consent was also obtained from the subjects. All subjects were apparently healthy.

Comment [CC3]: How was it proved that these individuals were exposed to the said pollutants from cement only? There might have been some other sources of exposure too? How did you exclude that?

Comment [CC4]: For example ?

2.4 Eligibility

Inclusion Criteria

The cement loaders involved in the study were those that had been exposed to cement dust for a minimum period of three months, who gave their consent to participate in this study and are adults between the ages of 20 to 60 years of age.

The unexposed control subjects were apparently healthy male subjects who reside outside the vicinity of the cement exposed areas. Subjects with no history or signs suggestive of respiratory, haematologic, bone or liver diseases were considered eligible and selected into both the exposed and unexposed groups.

Comment [CC5]: How was it proved that they did not have any other disease before becoming cement loaders? Did you consult their medical record from some reliable source? If yes kindly provide the evidence
Were they smoker too?

Comment [CC6]: Form where did this figure come, please support it with reference

Comment [CC7]: What about smoking??

Exclusion Criteria

Subjects with previous exposure to any occupational agents other than cement silica etc, were excluded from the study. Also, those with history or diagnosed case of asthma or any respiratory diseases or other diseases like diabetes mellitus, pulmonary tuberculosis, having history of acute or chronic infection or recent case of hospitalization, and those with these chronic illnesses were exempted from the study.

Those who had worked for less than three months as well as those who did not consent were excluded from this study.

With the aid of questionnaire and interview, all participating cement loaders were interviewed by trained interviewers. All participants went through medical assessment to rule out the presence of diseases like asthma, diabetes, hypertension, anemia, cancer, infections or

Comment [CC8]: For example?

Comment [CC9]: What about smokers

Comment [CC10]: Form where did this figure come, please support it with reference

Comment [CC11]: Help

Comment [CC12]: What was this training

those who have recently had blood transfusion, thyroid and heart problems. Participants with diseases, drug therapy and alcohol, antioxidants, exposure to deadly substances or radiation therapy were not included in the study.

Comment [CC13]: What about underweight, obese, those working in some other industry like petroleum, stone work? Addicts?

2.5 Informed Consent and Ethical Clearance

Ethical approval for this research was obtained from the Rivers State Health Research Ethics Committee. Permission was also gotten from the authorities of cement loading sites/shops and Dangote cement depot, RIVOC, Trans-Amadi, Port Harcourt. Informed consent was given by individuals before recruitment into the study.

Comment [CC14]: Taken

Comment [CC15]: And written

2.6 Sample Collection, Transportation, Processing and Preservation

After seeking consent and giving explanations, venous blood samples were drawn from the antecubital fossa of this study subjects using vacutainer sample containers. This is in accordance to the description given by Cheesbrough, (2010).

Venous blood sample of 3ml size was drawn into a vacutainer blood sample container of 0.5 ml of 1.2 mg/ml ethylene diamine tetra-acetic acid (EDTA) and properly mixed for the assessment of ESR, full blood count to ascertain haemoglobin count, white blood cell count and white cell indices, platelet count, platelet indices.

Comment [CC16]: By whom?

Comment [CC17]: How was pseudothrombocytopenia excluded?

Within 24 hours, every venous blood sample drawn into dipotassium ethylene diamine tetra-acetic was completely assayed, to obtain complete blood count.

Comment [CC18]: Why not within 4 hours?

To the point of analysis, all drawn samples were conveyed via cold chain (ice packs/crushed ice in air tight and sealed thermo-container).

Comment [CC19]: Were they frozen? Freezing and thawing may affect the hematological parameters. Kindly clarify

2.7 Sampling Technique

Simple random sampling technique was used for recruitment to give everyone an equal chances of been selected into the study so as to rule out bias.

2.8 Sample Analyses

Determination of Erythrocyte Sedimentation Rate (ESR)

The blood sample for erythrocyte sedimentation rate was estimated using westergreen manual method

Procedure

Comment [CC20]: Who performed this test

0.4 ml (400 ul) of Trisodium citrate solution was added into a westergren bucket, add 1.6 ml (1,600 ul) of blood was also added to the Zero "0" mark. It was stood vertically undisturbed and free from vibrations for 1 hour in the Westergren stand after which the levels of the red cell was read and results were recorded in mm/hr.

Measurement of Full Blood Count

The study samples for complete blood count were all assayed with the use of haematology automated analyzer.

Comment [CC21]: Name and specifications, company?

Procedure

Here, the blood sample to be assayed was mixed up with a vortex mixer then the lid of the sample bottle was turned open and the sample inserted into the Mythic auto-analyser through the machine probe. The results of the analysis were displayed on the machine screen and were printed out.

3.0 RESULT

Demographic Characteristics of Subjects

A total of one hundred (100) apparently healthy cement dust exposed male participants within the age of 20–60 years and one hundred (100) apparently healthy non cement dust exposed male control participants of age between 20 and 60 years were recruited for the study.

In Table 1, the age mean levels of Control and Exposed subjects are 38.69±8.21years and 38.78±1.04years respectively. Marital status was categorized as “married” and “single”. In the Control group, 56subjects were married and 44subjects were single while in the Exposed group, 60subjects were married and 40subjects were single.

Chart 1 :Demographic Characteristics of Subjects

Characteristics	Control subjects (n=100)	Exposed subjects (n=100)
Age(Yrs)	38.69 ± 8.21	38.78 ± 1.04
Marital Status		
Married	56 (28.00%)	60 (30.00%)
Single	44 (22.00%)	40 (20.00%)

Comment [CC22]: Why gender bias?

Comment [CC23]: What is the significance of this parameter? Why the divorced were not considered?

Comment [CC24]: What is the significance of this parameter? Why the divorced were not considered? Give reference

Comment [CC25]: Is shown in table 1

Comparison of Haematological Parameters of Control and Exposed Subjects

The mean ESR levels for the Control and Exposed groups were 10.46±8.74 and 20.56±13.68. There was a significant difference in the ESR values ($p < 0.0001$). The WBC counts for the Control and Exposed groups were $5.97 \pm 2.32 \times 10^9/L$ and $4.44 \pm 3.81 \times 10^9/L$. There was a significant difference in WBC counts ($p = 0.008$). The RBC level for the Control and Exposed groups were $5.47 \pm 0.94 \times 10^{12}/L$ and $5.08 \pm 0.76 \times 10^{12}/L$. There was a significant difference in RBC level ($p = 0.001$). The Hb level for Control and Exposed groups were statistically significant difference 15.03 ± 1.32 g/dL and 4.24 ± 1.28 g/dL. There was a significant difference in Hb ($p < 0.001$) MCV level for Control and Exposed groups were 83.14 ± 5.03 and 83.59 ± 6.38 . There was no significant difference in MCV levels ($p = 0.5843$) MCH level for Control and Exposed groups were 28.71 ± 2.71 pg and 28.59 ± 2.79 pg. There was no significant difference in MCH ($p = 0.7586$). PLT level for Control and Exposed groups were 214.20 ± 70.54 and 218.00 ± 86.45 There was no significant difference ($p = 0.7306$). MPV levels for Control and Exposed groups were 11.13 ± 2.64 0.0699 and 10.61 ± 1.05 0.0699. There was no significant difference in MPV ($p = 0.0699$). Neutrophil levels for Control and Exposed groups were 43.51 ± 11.85 % and 39.32 ± 18.51 %. There was no significant difference in Neutrophil levels ($p = 0.0592$). Lymphocyte levels for Control and Exposed groups were 45.90 ± 12.82 % and 60.97 ± 7.58 %. There was no significant difference in Lymphocyte levels. (P-value=0.0525). Eosinophil levels for Control and Exposed groups were 3.53 ± 2.30 % and 3.07 ± 2.29 %. There was no significant difference in Eosinophil levels ($p = 0.1625$). The Basophil levels for Control and Exposed groups were 0.10 ± 0.09 % and 1.11 ± 0.12 %. There was no significant difference in Basophil levels ($p = 0.4670$). Monocyte levels for Control and Exposed groups were 6.14 ± 3.14 % and 3.44 ± 3.08 %. There was significant difference in Monocyte levels ($p < 0.0001$).

Comment [CC26]: Mere repetition of what is written in table, it is redundant

Table 1 Comparison of Haematological Parameters of Control and Exposed Subjects

Subjects	Parameters													
	ESR (mm/hr)	WBC (x10 ⁹ /L)	RBC (x10 ¹² /L)	Hb (g/dL)	MCV (fl)	MCH (pg)	MCHC (g/dL)	PLT (x10 ⁹ /L)	MPV (fl)	NEUT (%)	LYMP (%)	EOS (%)	BASO (%)	MONO (%)
Control	10.46	5.97	5.47	15.03	83.14	28.71	38.17	214.20	11.13	43.51	45.90	3.53	0.10	6.14
(n=100)	±8.74	±2.32	±0.94	±1.32	±5.03	±2.71	±32.22	±70.54	±2.64	±11.85	±12.82	±2.30	±0.09	±3.14
Exposed	20.56	4.44	5.08	14.24	83.59	28.59	34.18	218.00	10.61	39.32	60.97	3.07	1.11	3.44
(n=100)	±13.68	±3.81	±0.76	±1.28	±6.38	±2.79	±4.68	±86.45	±1.05	±18.51	±7.58	±2.29	±0.12	±3.08
T value	6.151	3.401	3.227	4.324	0.5480	0.3077	1.226	0.3448	1.823	1.897	1.951	1.402	0.7288	6.127
P value	<0.0001	0.0008	0.001	<0.001	0.5843	0.7586	0.2218	0.7306	0.0699	0.0592	0.0525	0.1625	0.4670	<0.0001
Remark	S	S	S	S	NS	NS	NS	NS	NS	NS	NS	NS	NS	S

Key: ESR- erythrocyte sedimentation rate, WBC- white blood cells, RBC- red blood cells, Hb- haemoglobin, MCV- mean corpuscular volume, MCHC—mean corpuscular haemoglobin concentration, MCH- mean corpuscular haemoglobin, PLT- platelets, MPV-mean platelet volume, LYMP-lymphocytes, EOS-eosinophils, BASO- basophils, MONO-monocytes, S-significant at p<0.05 and NS-not significant at p>0.05..

4.0 DISCUSSION

This study evaluated some haematological among individuals exposed to cement dust. In this study, 100 subjects between the ages of 20 and 60 years were selected from the population of interest (cement loaders) and 100 subjects were recruited from non-cement loaders population to form the control group. The subjects were all males since the society considers this occupation as fit for only males. Due to the wide age bracket, it was evident that more married subjects made greater number in the study than the single subjects as 56 (28%) of the exposed subjects and 60 (30%) of the control subjects were married while 44 (22%) and 40 (20%) respectively were single. This choice of age bracket represents the employable and career age range which is ideal for easy subject recruitment especially in a study like this.

Comment [CC27]: Discussion is deficient
Take each parameter turn by turn, compare it with result of other study and mention any difference or similarity giving the possible reasons of any difference if any

Comment [CC28]: Parameters

Comment [CC29]: This is not part of discussion, it may be included in the introduction

The results from this study showed that the exposed subjects had significantly, higher ESR but significantly lower and monocyte count compared to the control subjects. The increase in ESR may be due to inflammation that resulted from the cement dust. According to Bray et al. 2016, ESR was reported to be associated with infections or tissue inflammation. In agreement with this finding, Pollard, (2016) said constituents of cement such as silica-alumina have been reported to cause inflammatory reactions. This finding agrees with the work of Erabhor *et al.*, 2013 and Westerberg *et al.* (2019) reported a notable increase in erythrocyte sedimentation rate in the cement dust exposed subjects compared to the control subjects which is in total consonance with this present study.

This study observed a significantly lower WBC and monocyte count in the exposed subjects, compared to the control subject. The lower WBC level may be due to disruption of white blood cell by the cement dust. According to Emmanuel et al. (2015), cement dust increase the changes of disruption of white blood cell parameters in the body. Farheen et al. (2017) also reported similarly that components in cement dust have effects on haematopoietic system. This finding agrees with the work of Okonkwo *et al.* (2015) who reported decrease in white blood cell count in cement workers. A contrary finding was reported by Erhabor *et al.* (2013). A study conducted by Farheen, (2017); John and Olubayo, (2011) reported significant changes in haematological parameters such that there was significant increase in total white blood cell count (WBC count) which is in contrast to the reduced WBC levels of cement dust exposed subjects compared to the control subjects as recorded in this particular study. The elevated monocyte count among cement exposed subjects in their study is not in consonance with this very study which recorded reduced monocyte counts. According to Patil et al. (2015), generally, exposure to dust causes alteration in blood cell counts. This can be due to allergic reactions and inflammation which in turn is evoked by dust entering the lungs. This has been observed even from dust generated from other dusty materials

Comment [CC30]: How does it prove that this effect was due to cement only? They might have been using some drugs like NSAIDS? How was this excluded

The data from this study showed that the exposed subjects had significantly lower Hb and RBC compared to the control subjects. This may could be that components of cement dust interferes with haemopoietic process. Report by Emeka et al. (2015) suggested that alteration in haematologic parameters could be due to the effect of cement dust components such as aluminum, calcium and silicon on the cellular blood component formation system. This finding agrees with the work of Ahmad and Akhter, (2018). Many ailments have been linked to cement dust exposure and according to them there was significant reduction in the values of haematocrit and haemoglobin level. This study is in consonance with the study carried out by Nwafor *et al.* (2019), which illustrated some haematological parameters with significant decrease as seen in

Comment [CC31]: Poor construction of sentence, kindly rewrite it

the values of red blood cell counts and haemoglobin levels due to exposure to cement dust. Long- term cement dust subjection can lead to the damage of the haemopoietic structure. A study done in Bangladesh, by Ahmad Rahnuma and Akhter Qazi Shamima in (2018) on cement dust related to total count of red blood cell, suggested that cement dust exposure leads to haemoglobin concentration damage and reduction of red blood cell count. This is in agreement with this study. In addition, a study carried out by Jacob *et al.* (2020) on cement workers showed that Hb was decreased in value also.

CONCLUSION

This study has shown that exposure to cement dust can alter haematopoietic processes which could result to anaemia, and affecting total white blood count cells. Also, exposure to cement has shown to cause inflammation.

RECOMMENDATIONS

The cement loaders should be educated on the importance of using adequate PPE and the need for regular medical checkup for the cement workers.

REFERENCES

- Ahmad, R. and Akhter, Q.S. (2018). Effects of exposure to cement dust on haemoglobin concentration and total count of RBC in cement factory workers. *Journal of Bangladesh Society of Physiology*, 13 (7), 68 – 72.
- Amadi, A. I and Dimkpa, K (2020). Towards waste minimization in wet Trades: metrological based model of cement demand in Port Harcourt metropolis. *The Environmental Studies Journal*, 3(1), 61-76.
- Azah, N., Antai, A.B., Peters, E.J. and Osim, E.E. (2002). Effect of exposure to dust generated from crushing of granite rocks on the lung function of south eastern Nigerian children. *Nigerian Journal of Physiology Science*, 17 (1-2), 42-47.
- Bray, C., Bell, L.N., Liang, H., Haykal, R., Kaiksow, F., Mazza, J.J. and Yale, S.H., (2016). Erythrocyte sedimentation rate and c-reactive protein measurements and their relevance in clinical medicine. *World Medical Journal*, 115 (6), 317-322.
- Cheesbrough, M. (2010). District laboratory practice in tropical countries, 2nd edition, Part 2. Cambridge: Cambridge University Press.
- Dunuweera, S.P. and Rajapakse, R. M. G. (2018). Cement Types, Composition, Uses and Advantages of Nanocement, Environmental Impact on Cement Production, and Possible Solutions. *Advances in Materials Science and Engineering*, 2(1), 1-11.

Comment [CC32]: in

Comment [CC33]: Where? How was it proved, did you perform biopsy?

Comment [CC34]: 8 out of 28 references are older than 8 years , need replacement with new ones

Emeka, E. N., Silas, A. U., Peter, U. A. and Victor, C. U. (2015). Cement dust exposure affects haematological parameters in cement loaders in Enugu Metropolis, South-East Nigeria. *Journal of Experimental Research*, 1(4), 1 - 4.

Emmanuel, T.F., Abiam, U.A., Okaka, A.N. and Alabi, O.J. (2015). Effects of cement dust on the haematological parameters in Obajana cement factory workers. *European Scientific Journal*, 11 (27), 256-266.

Erhabor, O., Kebbe, I.Z., Isaac, A., Yakubu, Y., Marafa, A.N., Okwesili, H.A., Buhari, A. and Ezenwa, A.O. (1996). A study of the prevalence of occupational diseases and the factors affecting their identification and reporting in Nigeria. *Nigeria Medical Journal*, 31, 12-14.

Comment [CC35]: Old reference

Farheen, A., Hazanri, M. A. H., Khatoon, F., Fareen, S. and Qudsiya, S. M. (2017). Haematological parameters are acutely effected by cement dust exposure in construction workers. *Annals of Medical Physiology*, 1(1), 31 – 35.

Feng, L., Zhou, L., Gu, D.A., Gell, S. and Mackay, J.P. (2005). Structure of oxidized α -haemoglobin bound to AHSP reveals a protective mechanism for haem. *Nature*, 435, 697-701.

Comment [CC36]: Old reference

Gawad, A.E., Schlichting, O., Strauss, L. and Zeitz, O. (2009). Antiapoptotic properties of erythropoietin: Novel strategies for protection of retinal pigment epithelial cells. *Eye*, 23, 2245-2250.

Gbadebo, A.M. and Bankole, O.D. (2007). Analysis of potentially toxic metals in airborne cement dust around Sagamu, Southwestern Nigeria. *Journal of Applied Science*, 7, 35-40.

Comment [CC37]: Old reference

Gordon, M. (2005). Postgraduate haematology (5th ed.) Oxford: Oxford Blackwell Science.

Comment [CC38]: Old reference

Jacob, R. B., Obianuju, M. C. and Deborah, I. P. (2020). Haematological Alterations among cement loaders in Port Harcourt, Nigeria. *Asian Journal of Medicine and Health*, 18(7), 1 – 8.

John, O. O. and Olubayo, M.A. (2011). Biochemical and haematological profile in Nigerian cement factory workers. *Research Journal of Environmental Toxicology*, 5 (2), 133-140.

Comment [CC39]: Old reference

Kakooei, H., Gholami, A., Ghasemkhani, M., Hosseini, M., Panahi, D. and Pouryaghoub, G. (2012). Dust Exposure and Respiratory Health Effects in Cement Production. *Acta Medica Iranica*, 50(2), 122-126.

Comment [CC40]: Old reference

Koury, M. J., Mahmad, N. and Rhodes, M. M. (2009). Origin and development of blood cells in Greer, J. P., Foerster, J., Rodgers, G. M., Paraskevas, F., Glader, B., Arber, D. & Means, R. T. (Eds). *Clinical Haematology* (pp 79 – 105). Philadelphia: Lippincott Williams.

Comment [CC41]: Old reference

Mojekwu, J. N., Ademola, I. and Sode, O (2013). Analysis of the contribution of imported and locally manufactured cement to the growth of gross domestic production of Nigeria. *African Journal of Business Management*, 7(5), 360-371.

Comment [CC42]: Old reference

Nwafor, P. C., Odukanmi, O. A., Salami, A. T., Owonikoko, M. and Olaleye, S. B. (2017). Evaluation of a cement dust generation and exposure chamber for rodents: Blood Heavy metal status, haematological variables and gastrointestinal motility in Rats. *Africa. Journal of Biomedical Research*, 22, 79 – 87.

Ogunbileje, J. O. and Akinosun, O. M. (2011). Biochemical and haematological profile in nigeria cement factor/workers. *Research Journal of Environmental Toxicology*, 5, 133-144.

Comment [CC43]: Old reference

Okonkwo, C. O. J. Ugwu, C. E., Anakor, A.C.F., Dike, C.C. and Nwobodo, E. (2015). Effects of Cement Dust on Haematological Parameters of Cement Workers in Asaba, Delta State, Nigeria. *Journal of Environmental Science, Toxicology and Food Technology*, 9, 5-8.

Patil, P.S., Aithala, M. and Das, K. K. (2015). Effect of occupational exposure on blood cell counts, electrocardiogram and blood pressure in rice mill workers. *Journal of Clinical and Diagnostic*, 2(1), 221 – 227.

Pollard, K.M., (2016). Silica, Silicosis and Autoimmunity. *Frontiers in Immunology*, 7, 97-107.

Rahmani, A. H., Almatroudi, A., Babiker, A.Y., Khan, A. A. and Alsahly, M.A. (2018). Effect of exposure to cement dust among the workers: An evaluation of health related complications. *Journal of Medical Science*, 6(6), 1159-1162.

Saleh, H. M. & Rahman, R. O. A. (2018). Introductory Chapter: Properties and Applications of Cement- Based Materials. <https://www.intechopen.com/chapters/59180>

Wekpe, V. O. and Fiberesima, D. (2020). Noise mapping around the host communities of the university of port harcourt, Nigeria. *Arts & Humanities Open Access Journal*, 4 (2), 43–48.

Weli, E.V and Efe, I.S. (2015). Climate and epidemiology of malaria in Port Harcourt region, Nigeria. *American Journal of Climate Change*, 4, 40-47.

Westberg, H., Elihn, K., Andersson, E., Persson B., Andersson, L., Bryngelsson, I. L., Karlsson, C. & Sjögren, B. (2016). Inflammatory markers and exposure to airborne particles among workers in a Swedish pulp and paper mill. *International Archives of Occupational and Environmental Health*, 89(5), 813-822.