

Original Research Article

A study on level of Awareness of Farmers about Agri Waste Management practices in Medak district of Telangana

ABSTRACT

Agriculture wastes are the residual from the growing agriculture crops such as cereals, pulses, oilseeds, horticulture and livestock etc. This paper focused on study on level of awareness of the respondents about agri waste management practices and its relationship with the profile of the farmers. The Ex-post facto research design was adopted for the study with a sample of 120, covering Medak district of Telangana state. From the analysis, it was found that majority of the farmers (55.00%) had high level of awareness, followed by medium (27.50%), and low (17.50%) level of awareness about agri waste management practices. From the analysis, it was found that infrastructure facilities and achievement motivation had positive and significant relationship with awareness at one percent level of significance. The variables age, education, farm size, farming experience, cropping pattern, innovativeness, information seeking behavior and training undergone had positive and significant relationship with awareness at five per cent level of significance. The remaining variables, cropping intensity and level of aspiration had positive and non-significant relationship with level of awareness about agri waste management practices.

Keywords : Agriwaste management, awareness

INTRODUCTION

Every year India produce 550 million tonnes of agriculture waste. More than 149 mt of the to residue is un-used, within that 90 mt of residues burned (MNRE, 2018). Telangana generates about 28.89 million tones of agriculture waste. After meeting the requirements surplus residue was 6.96 million tonnes, out of which 2.73 million tonnes were burnt. Ministry of Renewable and Energy Resources, estimates that agri-waste can generate more than 18000 million tones of power every year apart from generating green fertilizers to farms (1).

Agri waste is a serious economic, environmental and social problem and there is a need to analyze different types of waste produced in production, processing and management practices for better utilization of agri waste materials. Proper management of waste from agricultural operations can contribute to significant way of farm operations. By waste management all the crop wastes are placed in the right time and right place for best utilization in order to convert into useful products and control of pollution.

METHODOLOGY

An *Ex-post facto* research design was followed for the study. The State of Telangana was chosen since the researcher was familiar with local language and culture. Medak district of Telangana state was selected. Three mandals in the district and two villages from each mandal were selected using simple random sampling technique. Thus, a total of eight villages were selected. From each village 15 respondents were selected using simple random sampling technique thus constituting 120 respondents for the study.

Comment [WU1]: This paper has potential but needs a lot of improvement in all sections to become dependable academic literature

Comment [WU2]: The abstract does mention key things like sampling techniques used. Also, no implication/conclusion on the findings.
Needs improvement

Comment [WU3]: Residuals

Comment [WU4]: Once you have "and livestock" there is no need for etc

Comment [WU5]: On the level of ...

Comment [WU6]: The study found that, age, education.....

Comment [WU7]: However, cropping.....

Comment [WU8]: This section needs a lot of improvement

1. It is too short and does not give enough background to the issues under consideration. It should give a global outlook on agri waste with key scenarios before narrowing down to Telangana
 2. Not enough references to literature
 3. The problem statement and objectives of the study are not very visible
- A minimum 6 to 10 references should be enough to clearly address the deficiencies in this section.

Comment [WU9]: Provide the justification for the choice of design

The data from the respondents was collected with the help of interview schedule. The data collected was analysed and interpretations were drawn based on results. The statistical techniques frequency, percentage, , exclusive and inclusive class interval were adopted for analyzing data.

Awareness as a stage at which an individual comes to know about the existence of new idea or practice (). In this study awareness operationally defined as consciousness of individual about existence of agri waste management practices and agri waste uses in agriculture.

Comment [WU10]: How was awareness assessed?
Provide a clear description.

RESULTS AND DISCUSSION

The data was collected from the respondents about level of awareness of agri waste management practices analysed, interpreted, and accordingly the following results and conclusion were drawn.

1. LEVEL OF AWARENESS OF THE RESPONDENTS ABOUT AGRICULTURE WASTE MANAGEMENT PRACTICES

The results in the table 1. indicated that majority of the farmers (55.00%) had high level of awareness, followed by medium (27.50%), and low (17.50%) level of awareness about agri waste management practices.

Comment [WU11]: The discussion component of this section is inadequate. It fails to draw on literature to make significant meaning of the findings. A minimum of 15 to 20 references can aid in making very good discussions.
1. How does the current awareness levels impact the current environmental situation?
2. Are the levels good enough and how will it impact future waste practices?
3. Challenges of farmers affecting good sustainable practices and strategies to improve awareness (per the profiles)

Table 1. Distribution of farmers according to their level of awareness about agri waste management practices n=120

S.No.	Category	Frequency	Percentage
1	Low (46-55)	21	17.50
2	Medium (55-64)	28	27.50
3	High (64-75)	66	55.00
Total		120	100.00

Comment [WU12]: indicates

Comment [WU13]: indicate how these categories were obtained. Are they standards or derived?

Comment [WU14]: Check the numbers here because it does not sum up to 120.

The high level of awareness of farmers about agri waste management practices might be due to the fact that majority of the respondents grouped under medium to high farming experience, this might had exposed them to the agri waste management practices and respondents had medium information seeking behavior which imply that the farmers might have accessed agri waste management information sources (3).

Table 2. Distribution of farmers according to their level of awareness about rice waste management practices n=120

Crop	Waste	Practices /uses	Frequency	Percentage
Rice	Straw	Vermi-compost/compost	72	60.00
		Manure preparation	48	40.00
		Straw board and card board	18	15.00
		Animal feed and bedding material	92	76.66
		Mulching material	60	50.00

		Nursery bedding material	76	63.33
		<i>In situ</i> incorporation	80	66.00
	Husk and Bran	Poultry litter	33	27.50
		Energy source for brick klin	26	21.66
		Packing material	39	32.50
		Animal feed	92	76.66

From the table 2 it was observed that among different practices of straw management 76.66 per cent of the farmers were aware about uses of straw as animal feed and bedding material followed by *insitu* incorporation (66.00%), nursery bedding material (63.33%), compost/vermicompost (60.00%) mulching material (50.00%), manure preparation (40.00%) and card board preparation (15.000%).

It's also observed from the table 1 that among different ways of managing rice husk 76.66 per cent of the farmers aware about uses of husk as animal feed, followed by packing material (32.50%), poultry litter (27.50%), and energy source for brick klin (21.66%) respectively (4).

Comment [WU15]: You mean table 2?

Table 3. Distribution of farmers according to their level of awareness about sugarcane waste management practices n=120

Crop	Waste	Practices /uses	Frequency	Percentage
Sugarcane	Trash	Compost	46	38.33
		Animal feed (Tops)	24	20.00
		Mulching material	33	27.50

It could be observed from the table 3 among different ways of managing sugarcane trash, 38.33 per cent of the farmers were aware about use as compost, followed by (27.50%), mulching material and (20.00%) animal feed (5).

Table 4. Distribution of farmers according to their level of awareness about maize waste management practices n=120

Crop	Waste	Practices /uses	Frequency	Percentage
Maize	Stover	Vermi-compost	72	60.00
		Manure preparation	48	40.00
		Mulching material	36	30.00
		<i>In situ</i> incorporation	76	63.33

		Animal feed	85	70.83
	Green leaves(sweet corn)	Silage making	24	20.00
		Animal feed	92	76.66

From the table 4 it was observed that among different ways of managing maize stover, 70.83 per cent of the farmers were aware about use as animal feed, followed by insitu incorporation (63.33%), vermin compost/compost (60.00%), manure preparation (40.00%) and mulching material (30.00%).

It could be observed from the table 4 revealed that among different ways of managing maize green leaves 76.66 percent of the farmers aware about use as animal feed and silage making (20.00%).

Table 5. Distribution of farmers according to their level of awareness about Redgram waste management practices n=120

Crop	Crop waste	Practices /uses	Frequency	Percentage
Redgram	Stubbles and husk	Compost	03	02.50
		Animal shelter	12	10.00
		Animal feed (husk)	18	15.00
		Fuel	30	25.00

From the table 5 it was observed that among different practices of managing redgram stubbles, 25.00 per cent of the farmers were aware about use as fuel, followed by animal feed (15.00%), animal shelter (10.00%), and compost (2.50%).

Table 6. Distribution of farmers according to their level of awareness about cotton waste management practices (n=120)

Crop	Crop waste	Practices /uses	Frequency	Percentage
Cotton	Stalks	Compost	14	11.66
		Fuel	82	68.33
		Mulching material	78	65.00

From the results in the table 6 indicated that among different practices of managing cotton stalks, 68.33 per cent the farmers were aware about use as fuel, followed by mulching material (65.00%) and compost (11.66%).

Table 7. Distribution of farmers according to their level of awareness about vegetable and fruit crops waste management practices (n=120)

Crop	Crop waste	Practices /uses	Frequency	Percentage

Vegetable and fruit crops	Damaged fruits and vegetables and dead plants	Compost	32	26.66
		Animal feed	48	40.00
		Nursery bedding material	18	15.00

From the table 7 it was revealed that among different practices of managing damaged fruits and vegetables, 40.00 per cent of the farmers were aware about use as animal feed, followed by compost (26.66%), and nursery bedding material (15.00%).

Table 8. Distribution of farmers according to their level of awareness about livestock waste management practices (n=120)

Enterprise	waste	Practices /uses	Frequency	Percentage
Live stock	Dung and urine	Biogas	73	60.83
		Vermi composting	79	65.83
		Manure preparation (FYM)	85	70.83
		Bio insecticide	29	24.16
		Dung cake	76	63.33

The results in the table 8 indicated that among different practices of managing dung and urine 70.83 per cent of the farmers were aware about use as manure preparation (FYM), followed by vermi-compost (65.83%), dung cake (63.33%), biogas (60.83%) and bio-insecticide (24.16%).

2. RELATIONSHIP BETWEEN SELECTED PROFILE CHARACTERISTICS OF THE RESPONDENTS WITH THEIR LEVEL OF AWARENESS ABOUT AGRICULTURAL WASTE MANAGEMENT PRACTICES

Null hypothesis : There will be no significant relationship between the level of awareness about agricultural waste management practices and the profile characteristics of the respondents.

Table 9. Relationship between selected profile characteristics of the respondents with their level of awareness (n=120)

S. No.	Profile characteristics	Calculated (r) Value
1.	Age	0.192 *
2.	Education	0.182*
3.	Farm size	0.181*
4.	Farming experience	0.216*
5.	Cropping Intensity	0.117NS
6.	Cropping pattern	0.191*

Comment [WU16]: The Structure of this subsection makes it difficult to follow the arguments

1. Since this section is a combination results and discussion, each profile characteristic should have been thoroughly handled in a paragraph.

2. There should be an introduction to set the tone following discussions by introducing what is to be discussed. This Introduction should also highlight the contents of Table 9 and the various hypothesis

3. Then the discussions follow in well expanded paragraphs.

7.	Infrastructure facilities	0.251**
8.	Level of aspiration	0.172NS
9.	Innovativeness	0.193*
10.	Achievement motivation	0.253**
11.	Information seeking behaviour	0.212*
12.	Training undergone	0.234*

Comment [WU17]: delete

** Significance at 0.01 level

* Significant at 0.05 level

NS Non significant

Comment [WU18]: The focus is on the significant ones

Empirical hypothesis : There will be significant relationship between the level of awareness about agri waste management practices and the selected profile characteristics of the respondents.

Age Vs Awareness

From the findings it is evident that age had positive and significant relationship with awareness about agri waste management. The reason might be majority of the farmers belonged to middle age with medium level of farming experience (6).

Education Vs Awareness

From the findings it was found that there was a positive and significant relationship between education and awareness. The reason might be that most of the farmers were literate and it might have helped them to access information from different sources like magazines, journals, news papers and agriculture officers about agri waste management (7).

Farm size Vs Awareness

From the findings it was found that farm size was positively and significantly correlated with awareness about agri waste management. The reason might be attributed to the fact that diversified farming in the study area increases the scope for having more exposure about different types of agri waste and their management practices (5).

Farming experience Vs Awareness

Farming experience was found to be positively and significantly correlated with awareness about agri waste management. A farming experience increases awareness about various innovative practices of agri waste management and its utilization in agriculture, might have also increased (5).

Comment [WU19]: Discuss these in well expanded paragraphs with sufficient references

Cropping intensity Vs Awareness

Cropping intensity was found to be positively and non significantly correlated with awareness about agri waste management. The reason might be majority of respondents were small farmers and dependent on rainfed farming.

Cropping pattern Vs Awareness

Cropping pattern was found to be positively and significantly correlated with, awareness about agri waste management. The reason might be attributed to the fact that more diversified farming make farmers to access more information about agri waste products and their management.

Infrastructure facilities Vs Awareness

Infrastructure facilities was found to be positively and significantly correlated with awareness about agri waste management. Increased awareness about agri waste management might have encouraged farmers to establish on farm facilities to lead the benefits out of it (8).

Level of aspiration Vs Awareness

Level of aspiration was found to be positively and non significantly correlated with, awareness about agri waste management. The reason might be that majority of the respondents were with poor of infrastructure facilities and other resources required for management of agri waste.

Innovativeness Vs Awareness

Innovativeness was found to be positively and significantly correlated with, awareness about agri waste management. An innovative farmer always tries to adopt new technologies and willing to bear risk involved in adoption. This nature might led them to access more information about technology in order to minimize the risk involved in adoption of the technology (9)

Achievement motivation Vs Awareness

Achievement motivation was found to be positively and significantly correlated with awareness about agri waste management. The reason might be that motivation facilitate the farmers to identify suitability of technologies and access more information about agri waste management(10).

Information seeking behavior Vs Awareness

Information seeking behavior was found to be positively and significantly correlated with, awareness about agri waste management. The result might due to the fact that more exposure and access to the different information sources favours to develop more awareness about agri waste management practices (9).

Training undergone Vs Awareness

From the results it was observed that training received had a significant and positive correlation with awareness about agri waste management. The reason might be the trainings provides knowledge and skills about different agri waste management practices and helps in building up awareness among respondents (11).

Conclusion:

The high level of awareness of farmers about agri waste management practices might be due to the fact that majority of the respondents grouped under medium to high farming experience, this might had exposed them to the agri waste management practices and

Comment [WU20]: Infrastructure

Comment [WU21]: Discuss these in well expanded paragraphs with sufficient references

Comment [WU22]: This is more of summary of findings than conclusion
No recommendations

respondents had medium information seeking behavior which imply that the farmers might have accessed agri waste management information sources.

Correlation analysis of the level awareness about agri waste management practices and profile characteristics of respondents revealed that infrastructure facilities and achievement motivation had positive and significant relationship with awareness at one percent level of significance. The variables age, education, farm size, farming experience, cropping pattern, innovativeness, information seeking behavior and training undergone had positive and significant relationship with awareness at five per cent level of significance. The remaining variables, cropping intensity and level of aspiration had positive and non-significant relationship with level of awareness about agri waste management practices.

References:

1. MNRE, Ministry of New and Renewable Energy sources , Government of India, New delhi, 2018, www.mnre.gov.in/biomass resources.
2. Rogers, 1995. EM. Diffusion of innovations, 5th Ed.. The Free Press, New York; 1995.
3. Sindhu, N. 2012. A study on agriculture waste utilization in diversified agriculture. *M. Sc. (Ag.) Thesis*. Chaudhary Charan Singh Agricultural University, Hissar.
4. Mazhar, H.S. (2016). Analyzing the practices and importance of organic farming. *M.Sc. (Ag.) Thesis*. Bidhan Chandra krishi vishwa vidyalaya, west bengal, India.
5. Anuse, V.R. 2016. Impact of integrated sugarcane trash management technology (ISTM) on sugarcane growers. *M.Sc. (Ag.) Thesis*. Mahatma phule krishi vidyapeeth, Rahuri district , Ahmednagr , Maharashtra, India.
6. Vihariya payal, H. 2017. Awareness and adoption of vegetable growers about hazardous effect caused through pesticide residues in vegetables. *M.Sc. (Ag.) Thesis*. Anand agricultural university. Anandh.
7. Vineetha, A. 2018 Marketing behaviour of groundnut farmers in Anthapur district of Andhrapradesh. *M.Sc (Ag.) Thesis*. Acharya N G Ranga Agricultural University, Guntur , India.
8. Sravan kumar, T. 2012. A study on entrepreneurs of vermicopost technology in Guntur district of Andhra predesh *M.Sc. (Ag.) Thesis*. Acharya N G Ranga Agricultural University, Hyderabad, India.
9. Choudhary, J. G. 2013. Farmer awareness regarding agriculture pollution in Anand district. *M.SC. (Ag.) Thesis*. Anand agriculture university. Anand.

Comment [WU23]: So what does it mean in terms agri waste management
The way forward

Comment [WU24]: Please put together a minimum of 30 references
Ensure that all references conform to that of the journal

Comment [WU25]: Be consistent in the use of brackets

11. Nikunj Kumar, R. P. (2014). Comprehensive awareness among the farmers about the application of bio-fertilizers in Anand district. *M.Sc. (Ag) Thesis*. Anand agricultural university, Anandh.
10. Prashanth, P. 2011. A study on adoption of organic farming in cotton in Karimnagar district of Andhra Pradesh. *M.Sc.(Ag) Thesis*. Acharya N G Ranga Agricultural University, Hyderabad, India.

UNDER PEER REVIEW