

Original Research Article

A Comprehensive Appraisal of the Farming Scenario in Riverine Areas of Lower Assam, India

ABSTRACT

~~The present investigation~~This study was undertaken to ~~study the comprehensive appraisal of~~appraise the farming scenario of Riverine (*Char*) areas of lower Assam ~~based on~~in terms of prevailing climatic parameters, soil-site characteristics, existing cropping pattern and socio-economic conditions of ~~the~~farmers. Three *Chars* of the district, viz., Faujdar *Char*, Simlabari and Bamunpara Part IV ~~was~~were selected for the study. A multistage purposive cum random sampling design was adopted for the study. The average monthly temperature in the chars ranges from 18.9 to 28.6°C, which are usually suitable for most of the crops like *sali* paddy, maize, chilli etc. The analysis of physico-chemical parameters of the soil reveals predominance of sand and sandy loam texture. The available N, P and K contents ranged from 84.2 to 108.5, 38.6 to 42.3, and 127.9 to 152.6 ~~kg~~Kg/ ha, respectively. The average yield efficiency percentage were observed highest (320.21%) in potato. The per cent pest infestation was recorded as 15-25% for almost all the crops studied.

Key words: *Char*, cropping pattern, insect, questionnaire

Comment [U1]: No. The study did not study an appraisal. Simply say to appraise.

Comment [U2]: Not clear. Rephrase.

Comment [U3]: Include something on socio-economic conditions of farmers since it was one of the key factors studied.

INTRODUCTION

The riverbed of the Brahmaputra is dotted with a number of riverine areas (*Chars*) all along its length in the plains of Assam. These *Chars* are formed by the suspended materials brought from the catchment areas of the rivers on account of any obstruction in the course of the rivers. Out of 3.61 lakh ha of *Char* area, an arable area of 2.42 ~~lakh x 10⁵~~ ha is supporting a large population of the farming community (GoA, 2018). The chars account for about 5 percent of the total area of the state spreading across 14 districts, 55 blocks and around 2,300 villages (GoA, 2016). Economic backwardness limits *Char* dwellers' livelihood options. The income opportunities of farmers are severely restricted by recurring floods and erosion (Chakraborty, 2012). As per a-the report (GoA, 2018), the Dhubri district encompasses 480 numbers of villages in the *Char* areas covering an area of 64,767 hectares. The socio-economic backwardness is a matter of serious concern in the *Char* areas. For majority of the *Char* dwellers,

Comment [U4]: Lakh is not so popular in many parts of the world. Since this paper is intended to reach an international audience, I suggest you replace lakh with 10 raised to the power 5, which is the same as 100,000.

Comment [U5]: Indicate the name of the report.

agriculture is the primary source of livelihood (Nath *et al.*, 2021). Identification and quantification of socio-economic factors inhibiting the growth and development of *Char* farmers can pave the way to eradicate the problems of *Char* people.

Most of the *Chars* made up of alluvial deposits, mostly silt and sand. The wide variation in physical and natural features, combined with natural fertility of the alluvial deposits, facilitates the cultivation of a wide ~~varieties~~ variety of *rabi* and summer crops (Talukder *et al.*, 2009). The ease of tillage operations in the *Char* land assist farmers to ~~take up of cultivate~~ various crops (Borkakati *et al.* 1999; Vadivelu *et al.* 2005).

To increase agricultural productivity, considering the immediate and future opportunities of these lands, an evaluation was attempted to study the suitability and performance of various crop. An attempt ~~has also been was also~~ made to prepare an appropriate cropping plan for further enhancement in crop productivity of the study area.

Comment [U6]: In addition to this justification, you need to justify this study based on knowledge/research gap. In other words, you need to formulate a strong problem statement for your paper. What are the gaps in the literature that your paper seeks to address? This has to feature in your introduction.

Comment [U7]: Not clear. Rephrase.

RESEARCH DESIGN AND METHODOLOGY

Selection of the study area:

A preliminary survey cum Participatory Rural Appraisal (PRA) programme was carried out in 2021 in three villages ~~of in~~ Dhubri district, *viz.*, Fauzdar *Char*, Simlabari and Bamunpara Part IV (Fig. 1). A multi-stage ~~random sampling design~~ was adopted for the study.

Comment [U8]: But in the abstract you said 'a multi-stage purposive cum random sampling'. Please reconcile.




Fig. 1: Location of the study areas with GPS coordinates

Period of Study:

Necessary information were collected based on survey using a standard questionnaire during the first quarter of 2021 through multiple visits to the study area.

Secondary data from various sources viz., District Agricultural Office, ~~Deptt~~Department of Economics and Statistics, Assam Agricultural University, Jorhat ~~are~~ were used to corroborate demography, land tenure system, crop activity, land use and cropping pattern, farming system and price of inputs and outputs, maximum and minimum area under certain crops, net benefit and cost benefit analysis. Average yield efficiency percentage (AYEP) was calculated as:

$$\text{AYEP} = \frac{\text{Average crop yield per ha in the given area}}{\text{Mean crop yield per ha in the state}} \times 100$$

Comment [U9]: Why should the average yield/ha in the state be used as the reference? Why don't you use average yield/ha at the national level as the reference instead, so that your figures would be more representative?

Soil samples (0- 15 cm) were collected from the study areas and analysed in the laboratory of Department of Soil Science, SCS College of Agriculture, Assam Agricultural University, Rangamati, Dhubri, Assam. Particle size distribution, soil reaction, organic carbon, and available N, P, K were estimated following standard methods.

Chart 1 : Survey Area:

Sl. No.	Village	GPS Coordinates
1.	Faujdar Char	26.2167430 N, 90.3712150 E, Alt.: 31 m
2.	Bamunpara part IV	26.237328 ⁰ N, 90.258555 ⁰ E, Alt.: 33 m
3.	Simlabari	26.245153 ⁰ N, 90.409363 ⁰ E, Alt.: 36 m

Comment [U10]: This should be captioned as a table, and reference should be made to the table in the text.

RESULTS AND DISCUSSION

Climate:

The Char areas visited received an average annual rainfall of 1,856 mm. About 63.19, 26.38, 6.95 and 3.48 per cent of the annual rainfall occurred in the monsoon, pre-monsoon, post-monsoon and winter seasons, respectively. The rainfall during the period from November to February is sporadic and unreliable (Fig. 2). The occasional rainfall that occurs during this period helps to replenish the moisture deficiency of soils. However, such rainfall depending on its timing and intensity may also cause damage to the standing rabi crops. The average monthly

Comment [U11]: It is better you report your findings in the past tense since the study has already been conducted. Change all the verbs from the present to the past tense.

temperatures in the chars vary between 18.9 and 28.6°C, which are usually suitable for most crops.




Fig. 2. Weather parameters prevalent in Char areas [average of ten years (2012 – 21)]

Rice is a thermal sensitive crop with temperature suitable for growth varying by growth phase of the crop (Wassman *et al.* 2009). The day temperature up to 35°C and night temperature of upto 25°C are suitable for rice growth during *sali* paddy growing period. At tillering the crop requires higher temperature than other stages of its growth. The crop requires 26.50-29.50°C for flowering. At ripening rice requires 20-25°C. Rice grows well in areas receiving average annual rainfall of 1000-1500 mm or more. The prevailing weather conditions at the visited chars were well within the suitable ranges for *sali* paddy.

Chilli grows well in areas with temperatures of 18-27°C during the day and 15-18°C during the night (Doorenbos and Kassam, 1979). A warmer night temperature above this range induces earlier flowering. The weather condition during chilli growing winter months are congenial in the Char areas of Dhubri. Well aerated and well drained soil condition of the visited Char areas also support proper growth, flowering and fruiting of chillis.

A temperature range of 18-20°C is the optimum for germination of maize seeds (Doorenbos and Kassam, 1979). The crop can tolerate hot and dry atmospheric conditions so long as sufficient water is available to the plant. It is also relatively tolerant to water deficits

during the vegetative and ripening stages. Maize grows well in most soils except very sandy soil. The suitable agro-ecological conditions of *Char* areas support excellent crop growth of maize. Jute is a tropical crop growing under wide variation of climatic conditions. The optimum range of temperature required is 18-33°C. The crop is basically self-pollinated and needs long daylight for its growth. It grows well with an annual rainfall of 1,500 mm or more, with the occurrence of at least 250 mm of rainfall during each of the months of March, April and May (Banglapaedia, 2006).

Soil characteristics:

The soils in the study areas have an irregular relief of stratified sandy and silty deposits. The soils form under a diverse set of environmental conditions including vegetation, relief, drainage and age of sedimentation, which together lead to different patterns of soil fertility, crop suitability, land use and agronomic practices. The soil samples were collected during our field visits from the Fauzdar, Bamumparapart IV and Simlabari *chars*. The physio-chemical parameters of the soils at the three *Chars* under study (Table 1) indicated that soils contain sand in a large proportion and the texture varied from sand to loamy sand. Meager quantities of the organic matter (0.38 to 0.51%) in the areas can be attributed to the recent deposit of sand in the top layer. The overall nutrient status of the soils was found to be medium to high (Table 2). The available N, P and K contents ranged from 84.2 to 108.5, 38.6 to 42.3 and 127.9 to 152.6 kg/ ha, respectively. Such soils are usually suitable for many crops. *Char* soil is usually productive as it benefits greatly from seasonal flooding through nitrogen fixation by blue-green algae and bacteria. The flooding also benefits through the addition of organic matter and nutrients from the decomposed flora and fauna. The seasonal fluctuation between aerobic and anaerobic soil conditions also makes phosphorus and possibly potash more readily available to plants (Brammer, 2000).

Table 1: Physical and chemical characteristics of soil.

Location	Colour	% Sand	% Silt	% Clay	Texture	pH	OC (%)
Faujdar <i>Char</i>	10YR7/1	85.8	10.7	3.5	Loamy sand	6.7	0.48
Bamunpara part IV	10YR7/1	78.6	12.5	8.9	Loamy sand	6.8	0.51

Simlabari	10YR6/1	92.4	3.5	4.1	Sand	6.5	0.38
-----------	---------	------	-----	-----	------	-----	------

Table 2: Nutrient status of soil:

Location	Available N (kg/ha)	Available P (kg/ha)	Available K (kg/ha)
Faujdar Char	108.5	38.6	134.4
Bamunpara part IV	122.7	42.3	152.6
Simlabari	84.2	40.5	127.9

Production status of various crops:

The yield performance of summer rice (6600 kg/ha) is superior in the study area compared to *sali* rice (5400 kg/ha) owing to favourable prevalent weather conditions during the summer season. It ~~is~~ was observed that the average yield efficiency of summer and *sali* rice ~~are~~ were 254.53 and 250.00 per cent, respectively compared to the state average (Table 3). Among various pulses grown in Char areas, blackgram exhibits great promise with 175.59 AYEP. In case of oilseed crops, the AYEP of rapeseed, lentil, nigerand sesamum are 194.49, 117.80, 161.29 and 84.51, respectively. Jute is a popular crop among the farmers and perform reasonably well in the Char areas with AYEP of 162.31. The performance of potato (AYEP: 320.21) is very encouraging and suitable soil and climatic conditions might be attributed for such higher yield in the study area. Chilli, though cultivated extensively by the Char dwellers and resulted in better yield compared to state average (AYEP: 146.99), incidence of pests and diseases are of major concern for realizing the potential yield.

Comment [U12]: Report your findings in the past tense.

Table 3: Average yield efficiency percentage of major crops.

Crops	Av. yield in study area(kg/ha)	Av yield in State (kg/ha)	Av yield efficiency percentage (AYEP)
Sali rice	5400	2160	250.00
Summer rice	6600	2593	254.53
Rapeseed	1200	617	194.49
Back gram	1050	598	175.59
Green gram	900	753	119.52
Lentil	900	764	117.80

Comment [U13]: Using the state as the reference is not the best. Use yield at the national level instead so that your findings would be more representative.

Niger	900	558	161.29
Sesamum	600	710	84.51
Jute	3600	2218	162.31
Potato	24000	7495	320.21
Chilli (green)	1439	979	146.99

Recommended land use pattern:

The factors like soil and climate along with food security, market price and local cultivation practice ~~are~~ were the major determinants of crops and cropping system in the *Char* areas. The *Char*-areas under study ~~have~~ had three overlapping cropping seasons, viz., *rabi*, summer and *kharif*. The *rabi* season is usually during a flood-free period, but the later part of the summer and the early part of the *kharif* seasons coincide with the flood period. Chilli, maize, jute, rice, peanut, wheat, millets, oilseeds, onion, garlic, sugarcane, peas and pulses are the crops grown in the *Char* areas. The crops are more diversified during the *rabi* season because of favourable climate during this period.

Rice is a common crop in every *Char*. *Sali* rice is cultivated by the farmers in medium land in spite of great risk of flood damage, as the monsoon ensures water requirement of the crop without irrigation. *Boro* rice (Summer paddy) is also common in all the *chars* under the study. *Ahu* rice (pre-monsoon season) is also found in most of the *chars*. Hybrid maize is also popularly grown in the study area. Jute is the single most dominant crop in the summer season in most of the *chars* of Dhubri district. Sesame is ~~growing~~ grown pre-monsoon (April-June) season in the *char* areas under the study. Different kinds of vegetables are grown, though pumpkin is more popular. Onion, chili, and mustard are traditionally cultivated in *chars* during *rabi* season (dry period). The spices like chili, black cumin and coriander are found to grow extensively. Lathyrus, cowpea, soybean and watermelon are also cultivated in the study area to a limited extent.

The dominant cropping pattern, however, varies from *Char* to *Char*. Several cropping patterns are followed in every *Char*. However, most of the patterns have at least one fallow season, though in a few patterns, two or three crops are also grown, especially in the medium high land. Number of crops in *rabi* season (dry period; November - March) is much higher than that in summer (April-June) or *kharif* season. The cropping patterns suggested for the *Char* areas are as follows:

Comment [U14]: Report in the past tense. Use "varied" instead.

1. Direct seeded early summer rice (February to May) - transplanted late winter rice (August to November)
2. Summer vegetables (February to May) - transplanted late winter rice (August to November) - potato/pea (December to February)
3. Summer vegetables (February to May) - transplanted late winter rice (August to November) - groundnut/pea (December to February)
4. Summer rice/summer pulse (February to May) - fallow – potato/vegetables/toria/wheat/pea (December to February)
5. Groundnut/melons (February to May) – fallow – early pulse/vegetables (December to February)
6. Summer rice (February to June) – fallow – sweet potato (September to February)

Different cropping pattern may be suggested for the *Char* areas under study on the basis of soil suited to crops, socio economic conditions, length of growing period of crops, prevailing pattern of flooding, BC ratio, etc. Crop diversification relying on introduction of new cultivars and adoption of locally adoptable agro-technologies may be suggested to address these issues. It is worth mentioning that, wheat is excluded as one of the major crops of the study areas due to problems associated with threshing, storage, pre-harvest sprouting, etc. Concurrently, late transplanted winter rice may be a viable alternative for such locations. To evade the flood, two crops before and after its occurrence could be suggested. Moreover, choice of the crop is mainly driven by the market demand, soil types, and available resources at the disposal of the farmers. It is expected that implementations of the suggested cropping pattern would enhance the productivity of crops in the areas under study.

Major insect pests:

Farmers, input dealers, extension officials and NGO staffs identified pests and diseases as hazards to crop production and storage across all the chars. The occurrence of pests and diseases in the fields is linked with weather. In the events of cold, fog, cloud, drizzle, hot, and seasonal weather change, the incidences of pests and diseases increase. The quality of seed, its processing/treatment before sowing, the occurrence of flood in the previous monsoon and the soil condition also factor in pest and disease incidences. Chilli is more affected than maize and jute. Plant, stem and root rotting, plant and root turning into yellow color, leaf curl, leaf turning

into white color, formation of root nodule, and cutting of roots and plant bases are the pest- and disease-related hazards for chilli. Cutworm, aphid, thrips, mite, leaf curl virus and anthracnose are among the major pests and diseases for chilli. Maize is affected by different pests and diseases including cutworm, stem borer, aphid and leaf blight at its early growth stage and jute by hairy caterpillar, semi-looper and stem rotting at the mid-stage. Insects and rodents also destroy the stored chilli and jute.

The major insect pests along with percent infestation in all the surveyed *Char* villages of the district are depicted in Table 4. From the investigation it was recorded that in all the crops pest infestation was recorded as 15-25%. The data were collected using simple random sampling in all the crop fields. Majority of the farmers used pesticides for controlling the pests.

Table 4: Major insect pests with per cent infestation.

Crop	Major insect pests	% infestation
Sali rice	Yellow stem borer	20-25
	Leaf folder	20-23
	Case worm	15-18
	Gundhi bug	30-35
	Whorl maggot	15-20
Summer rice	Yellow stem borer	15-20
	Leaf folder	15-22
	Case worm	12-15
	Gundhi bug	25-30
Rapeseed	Aphids	30-35
	Saw fly	15-20
Back gram	Aphids	20-25
	Pod borer	15-20
Green gram	Aphids	20-25
	Pod borer	18-20
Potato	Cut worm	15-20
	Red ant	20-25

Comment [U15]: Use single line spacing to help conserve space.

Formatted: Don't add space between paragraphs of the same style, Line spacing: single

Formatted: Line spacing: single

Formatted: Line spacing: single

Formatted: Line spacing: single

Formatted: Line spacing: single

Formatted: Line spacing: single

	Mole cricket	20-25
Jute	Bihar hairy caterpillar	15-20
Tomato	Fruit borer	25-30
	Aphids	15-20
	White fly	15-20
Okra	Jassids	20-25
	White fly	15-20
	Fruit borer	15-25
Brinjal	Shoot and fruit borer	20-25
Chilli	White fly	15-20
	Fruit borer	20-25
Cole crops	Cut worm	15-20
	Diamond back moth	10-15
	Aphids	15-20
Maize	Cut worm	15-20
	Maize borer	20-25

(Data represents the average of the three *Char* villages)

CONCLUSION

In view of the findings of the present study, some tangible measures may be adopted focusing ~~in~~on the areas of soil testing, IPM, capacity building, infrastructure facilities, etc in the *Char* areas of the district. Multiple cropping and intercropping should be actively encouraged. Mobilization of institutional support for creation of permanent assets, assured and timely availability of critical inputs, market linkage, credit facilities, etc is the need of the hour. It is expected that the socioeconomic standard of the farmers in the study area can be uplifted with the introduction of modern technologies and other amenities in conjunction with the recommendations provided. However, further investigations encompassing a larger study area

Formatted: Line spacing: single

Formatted: Line spacing: single

Formatted: Line spacing: single

Formatted: Line spacing: single

Formatted: Line spacing: single

Formatted: Line spacing: single

Formatted: Line spacing: single

Comment [U16]: The conclusion is missing. What you have stated are all recommendations. Your conclusion should emerge from your problem statement (which is lacking), and should extend the literature.

with higher numbers of parameters concerning socioeconomic status of the community may be advocated.

REFERENCES

- ~~Banglapaedia~~—Banglapaedia—the national encyclopaedia of Bangladesh. Asiatic Society of Bangladesh, Dhaka. 2006.
- Borkakati K., M. C. Talukdar, A. K. Choudhary, G. N. Hazarika, U. C. Bamah and S. Vadivelu Sand deposition by floods in Dhemaji district of Assam, submitted to the Commissioner, Agriculture, Govt. of Assam. 1999.
- Brammer, H Agro-ecological aspects of agricultural research in Bangladesh. The University Press Limited, Dhaka. 2000.
- Chakraborty, G. The 'Ubiquitous' Bangladeshis. *Econ. Pol. Weekly*. 2012; **47**(38):21-23.
- Doorenbos, J. and A.H. Kassam Yield response to water. FAO Irrigation and Drainage Paper 33, Rome. 1979.
- GoA. Assam Human Development Report. Govt. of Assam, Guwahati, Assam. 2016.
- GoA. Economic Survey of Assam. Directorate of Economics and Statistics, Govt. of Assam. 2018.
- Nath R. K., B. Sarma, M. Choudhury, P. Ahmed, G. K. Upamanya, S. M. Khayer, M. Rahman, G. K. Sarma, F. A. Ahmed and R. Sarma Socio Economic Status of Farming Community of Char Area of Dhubri District, Assam. *Asian J. Agril. Ext. Econ. Sociol.* 2021; **39**(9):14-20.
- Talukder, M. C., G. K. Goswami, A. Basumatary and A. K. Das. Crop suitability for Char areas of Nalbari district, Assam. *Agropedology*. 2009; **19** (1):41-46.
- Vadivelu, S., U. Baruah, B. P. Bhaskar, J. Thampi, D. Sarkar and P. S. Butte. Evaluation of soil-site suitability to rice-based cropping system in the river island of Majuli, Assam. *J. Ind. Soc. Soil Sc.* 2005; **53**:35-40.
- Wassman, R., S. V. K. Jagadish, K. Sumfleth, H. Pathak, G. Howell, A. Ismail, R. Serraj, E. Edona, R.K. Singh, and S. Heuer. Regional vulnerability of climate change impacts on Asian rice production and scope for adaptation. *Adv. Agron* 2009; **102**:91-133.