

Original Research Article

An Analysis of the Knowledge of Improved Cultivation Practices of Tomato of Khasi tribes in East Khasi Hills of Meghalaya

Comment [U1]: (*Lycopersicon esculentum*)

Comment [U2]: India

ABSTRACT

The main purpose of the study was to determine the extent of knowledge of Khasi tribe on improved cultivation practices of tomato. The study was conducted in Mawryngkneng Block in East Khasi Hills District, Meghalaya in the year 2022. A descriptive research design was applied for the study. The primary data was collected from 120 respondents by personal interview method using pre-structure interview schedule. Knowledge of the farmers was measured by asking 13 questions in respect to tomato cultivation practices. Pearson's Product Moment Coefficient of Correlation (r) was calculated to determine the relationship between independent variables and knowledge of the respondents. Finding showed that (78.33%) of the respondents had medium level of knowledge, (10%) of the respondents had high level of knowledge and only (3.33%) of the respondents had low level of knowledge about improved cultivation practices of tomato. Co-efficient of correlation (r) analysis indicated that land holding, farming experience, extension contact and sources of information are significantly associated with knowledge of farmers whereas variables like age, education, annual income, family type, size of family, and occupation was found to be non-significant with the knowledge of farmers towards improved cultivation practices of tomato.

Comment [U3]: extent of knowledge of small scale farmers of Khasi Tribeon....

Comment [U4]: How and why did you chose the study area? Purposively or Randomly??

Comment [U5]: pre-structured interview schedule

Comment [U6]: employed

Comment [U7]: colon

Comment [U8]: No recommendation?

Key Words: *Tomato cultivation, Knowledge, correlation co – efficient, Mawryngkneng.*

INTRODUCTION

Tomato (*Lycopersicon esculentum*) is botanically a fruit which is commonly used as a vegetable because of its nutritional content. It is an edible, red berry belonging to the family Solanaceae. Tomato plants are dicotyledon with a number of branching stem and a terminal bud at the tip. Tomato is composed of compound leaves particularly known as regular leaves but some consist of simple leaves known as potato leaves. It originated from western South America and Central America and spread to other countries as it can tolerate different climatic conditions. Tomatoes are widely grown in temperate climate across the world throughout the year. It is usually perennial but is cultivated as annuals. Tomato can be consumed in different ways i.e. raw or cooked, in many dishes and in making sauces, salads and drinks (Wikipedia).

The total cultivated area of tomato in the world was 4.84 and 4.76 million hectares with production 180.94 and 182.25 million tonnes and productivity 37.33 and 38.26 ton/ ha during 2017-18. Major producing countries of tomato in the world include China, India, USA, Turkey, Egypt, Iran, Italy, Spain, Brazil, Uzbekistan in which China was the largest producer of tomato in the world (FAO Statistical Database 2020).

India is the second largest producer of tomato in the world with a total cultivated area of 760,000 ha with production 18,399,000 tonnes and productivity 24,209.2 kg/ha (Anonymous Atlasbig.com 2018-2020). There has been an increasing rate in production of tomato from 18,732,000 during 2016 to 20,708,000 in 2017 but decrease to 19,377,000 during 2018 (FAO Statistical Database 2020). Andhra Pradesh is the leading producer of tomato in India with production 13.90% of tomatoes in the country. Tomato ranks third in priority in India after potato and onion. The major leading states of tomato production in India includes Andhra Pradesh, followed by Madhya Pradesh, Karnataka, Gujarat, Odisha, West Bengal, Telangana, Chattisgarh, Maharashtra, and Bihar (Agri Exchange.APEDA.gov.in). There has been a decreasing rate in the area and production of tomato during 2016-17 to 2017-18 from 797 ha to 789 ha in terms of area and from 20708 MT to 19759 MT in terms of production but there is an increase in 2018 to 814 ha in areas and 20515 MT in production (agricoop.nic.in 2020).

Comment [U9]: Bolden

Comment [U10]: comma

Comment [U11]: stems

Comment [U12]: comma

Comment [U13]: 'regular leaves'

Comment [U14]: 'regular leaves'

Comment [U15]: Cap the W

Comment [U16]: semi-colon

Comment [U17]: Wikipedia is not an authority or individual. Check the Wiki page again, you will see the main author, Use that name and include that you got this information from Wikipedia. Remember to include the year.

Comment [U18]: colon

Comment [U19]: was or is?

Comment [U20]: Anonymous is no person. Goggle this information and you will see the original author, the year and the journal of paper, which the source you are citing copied from.

Comment [U21]: what unit? Kg, Tonnes or what?

Comment [U22]: what unit? Kg, Tonnes or what?

Comment [U23]: semi-colon

Comment [U24]: Find out the unit and include it here

Comment [U25]: remove. Just write: FAO, 2020

Comment [U26]: volume of

Comment [U27]: colon

Comment [U28]: Year

Comment [U29]: remove

Comment [U30]: 20,708 MT to 19,759 MT

Comment [U31]: semi-colon

Comment [U32]: was

Comment [U33]: cultivated

Comment [U34]: 20,515 MT

Meghalaya is an agrarian state and is known for a large scale cultivation of vegetables both tropical and temperate. There has been an increasing rate in the area, production and productivity of vegetable crops. In addition the growth rate of vegetable crops has shown an increasing rate in area and production with respect to tomato. The department of agriculture has taken steps in order to increase production of vegetables in the state by motivating the farmers to cultivate vegetables in polyhouse by providing subsidy on the cost of construction of polyhouse which plays a significant role in agricultural sector as vegetable production from polyhouse increases its productivity by twice. Important tomato varieties grown in Meghalaya include Marglobe, Shillong Selection -1, Vaishali, Rocky, Mangala, Karnataka F1. (Anonymous, 2019. Department of Agriculture Government of Meghalaya, Shillong).

Comment [U35]: its

Comment [U36]: cultivated

Comment [U37]: comma

Comment [U38]: cultivated

Comment [U39]: Department of Agriculture

Comment [U40]: polyhouses,

Comment [U41]: polyhouse,

Comment [U42]: the

Comment [U43]: colon

Comment [U44]: Goggle the main author and replace

Comment [U45]: Knowledge of Improved Cultivation Practices of Tomato, level of awareness and present methods used should come in, plus the extent to which Agricultural Extension has influenced the farmers (respondents) in the study area

Meghalaya is also one of the states in India in which tomato can be grown. According to the Horticultural Statistic 2017 the cultivated area and production under tomato during 2014-15 was 2.20ha and 52 million tonnes, 2015-16 was 2.15ha and 34.02 million tonnes and 2016-17 was 2.12 ha and 33.95 million tonnes (Horticultural Statistics at a Glance, 2017). Horticultural sector in Meghalaya is now moving forward as the horticultural crops have more advantages than the indigenous crops in terms of generating employment, increasing the income and also have the potential to compete at national and international level (Lyngdoh, S. 2014).

Comment [U46]: comma

Comment [U47]: comma

Comment [U48]: average

Comment [U49]: remove

Comment [U50]: remove or you add with

Comment [U51]: National and International levels

Comment [U52]: Remove this S

Comment [U53]: No problem statement?? There could have been a brief reason with your literature, why or what makes you to do this research and what you want to achieve!

Comment [U54]: What is the main goal of your study What are the objectives, which will guide you to achieve these goal?

RESEARCH METHODOLOGY

Descriptive research design was followed for this study as it describes the characteristics or phenomenon that is being studied. The study was conducted among the Khasi tribes of East Khasi Hills District in Meghalaya. Mawryngkneng Block was selected purposively and 6 villages were selected randomly from this block and a total number of 120 farmers were selected proportionately for this study.

Comment [U55]: A brief geographical description of the study area since you are researching on crop production - Tomato

Comment [U56]: Which and which tools are you using to measure the variables? How are you going to measure these variables to achieve your goal?? What of the model?

RESULT AND DISCUSSION

Knowledge level of the respondents toward improved cultivation practices of tomato was studied and presented in Table 1 and Table 2 and figure 1.

Comment [U57]: Since there were no specific objectives, this discussion does not flow well in line with the topic.

Table.1. Distribution of respondents based on the knowledge level of the respondents.

Comment [U58]: You suppose to study the socio-economic characteristics of the respondents as one of the specific objectives

Comment [U59]: S/No.

Sl. No.	Statements	Response					
		Fully correct		Partially correct		Not correct	
		<i>f</i>	%	<i>f</i>	%	<i>f</i>	%
1.	Soil types suitable for tomato crop: Well drained loamy soils	7	5.83	95	79.17	18	15
2.	Climate: 21 - 24°C	41	34.17	71	59.16	8	6.67
3.	Soil testing	0	0	7	5.83	113	94.17
4.	Time of sowing: a) Summer (April – May) b).Autumn (August – September)	99	82.5	21	17.5	0	0
5.	Planting material: seeds	120	100	0	0	0	0
6.	Seed rate: Varieties – 300-350	30	25	80	66.67	10	8.33

Comment [U60]: Most of these variables are not known by educationally-disadvantaged respondents, i. e. small scale farmers who are the major producers of tomatoes.

	g/ha Hybrid – 100- 150g/ha						
7.	Irrigation	0	0	0	0	120	100
8.	Spacing: a). 60 cm X 45 cm b). 45 cm X 30 cm	10	8.33	81	67.5	29	24.17
9.	Varieties: Vaishali, Avinash, Rohit, Arka, Rocky, Cherranjeevi, Marglove, Shillong Selection-1	5	4.17	90	75	25	20.83
10.	FYM : 25 t/ ha	8	6.67	97	80.83	15	12.5
11.	Fertigation	0	0	0	0	120	100
12.	Growth regulators: Spray 1.25 ppm (625 ml in 500 litres of water) Triacantanol at 15 days after transplanting and at full bloom stage	4	3.33	116	96.67	3	2.5
13.	Seed Treatment : Trichoderma viride 4 g or Pseudomonas fluorescens 10 g or Carbendazim 2 g per kg of seeds 24 hours before sowing.	0	0	43	35.83	77	64.17

Comment [U61]: Source, year

Table.2. indicate 79.17 percent of the respondents have partial knowledge about the **soil type** suitable for tomato cultivation i.e. well drain loamy soil. About 59.16 per cent of the respondents have partial knowledge about the **climate** suitable for tomato cultivation. Majority of the respondents i.e. 94.17 per cent has no knowledge about **soil testing** .Majority of the respondents i.e. 82.5 per cent has full knowledge about the **sowing time** .100 per cent of the respondents were fully correct about the **planting material** of tomato. About 66.67 per cent of the respondents has partial knowledge about the **seed rate**|| 100 percent of the respondents do not have knowledge

Comment [U62]: per hectare

Comment [U63]: you do not start a sentence with figures. About 100% of the respondents.....

about irrigation practices since the study area was a rainfed area as they depend on rainfall for the crop. About 67.5 per cent of the respondents have partial knowledge while 24.17 per cent about the spacing required in tomato cultivation. 4.17 per cent of the respondents are correct about the varieties whereas 75 per cent are partially correct and 25 per cent of the respondents are not correct about the varieties for tomato cultivation. 80.83 per cent are partially correct about FYM requirement. 100 percent of the respondents do not have any knowledge about fertigation since the study areas depend on rain for irrigation. Majority (96.67 per cent) of the respondent has partial knowledge about application of Growth Regulator application. 64.17 per cent of the respondents has no knowledge about seed treatment. Similar findings were reported by Hadole *et al.* (2017), Pole (2018), Ansari *et al.* (2021) and Sangavi (2021).

Comment [U64]: a rain-fed

Comment [U65]: About

Table.2. Distribution of the respondents based on the overall level of knowledge (n=120)

S. No.	Knowledge level	Response	
		Frequency	Percentage
1	Low	4	3.33
2	Medium	94	78.33
3	High	12	10

Comment [U66]: Source, year




Fig. 1. Distribution of the respondents based on the overall level of knowledge

Comment [U67]: Source, year

From the above table 2 and figure .1, it is found that the level of knowledge towards tomato production technology is medium with 78.33 per cent followed by and high knowledge level 10 percent respectively and low knowledge level at 3.33 per cent.

Table 3: Association between selected independent variables with knowledge of the respondents.

Sl. No	Variabiles	Correlation Coefficient
1	Age	0.054NS
2	Education	0.063 NS
3	Annual Income	0.102 NS
4	Marital status	0.125NS
5	Family type	0.014 NS
6	Size of family	0.079 NS
7	Occupation	0.199 NS
8	Land holding	0.292 **
9	Farming experience	0.289**
10	Mass media exposure	0.020 NS
11	Extension contact	0.410**
12	Sources of information	0.511**

Comment [U68]: The author did not mention this in the Methodology

* = Significant at p = 0.01% * = Significant at p = 0.05 % NS = Non Significant

The result of correlation analysis in above table revealed the significance and non-significance of the socio-economic profile of the people and their level of knowledge of improved cultivation practices of tomato.

Comment [U70]: Where is the variable in the work?

To ascertain the relationship between profile of the respondents and their knowledge of improved cultivation practices of tomato in the area, the co-efficient correlation was worked out.

Comment [U71]: socio- economic not presented in the work

Table.3. revealed that independent variables like land holding, farming experience, extension contact and sources of information are significantly associated with knowledge of farmers.

Further, the variables age, education, annual income, family type, size of family, and occupation was found to be non-significant with the knowledge of farmers towards improved cultivation practices of tomato.

Comment [U72]: None of these variables were captured, analyzed or presented in the work

Comment [U73]: None of these variables were captured, analyzed or presented in the work

CONCLUSION

It is conclude that majority of the respondents has medium level of knowledge (78.33 per cent) followed by and high knowledge level at 10 per cent and low knowledge level at 3.33 per cent. Co-efficient of correlation (r) analysis indicated that land holding, farming experience, extension contact and sources of information are significantly associated with knowledge of farmers whereas variables like age, education, annual income, family type, size of family, and occupation was found to be non-significant with the knowledge of farmers towards improved cultivation practices of tomato.

Comment [U74]: Conclusion is very scanty. Need to be modified based on the objectives of the study
2. No recommendation

CONSENT

As per international standard or university standard, respondents' written consent has been collected and preserved by the author(s).

REFERENCES

Anonymous. 2015.

- a. Agricultural and Processed Food Products Export Development Authority. New Delhi.
- b. Agri-Exchange. Agricultural and Processed Food Products Export Development Authority. New Delhi.

Anonymous. 2017.

- a) Horticultural Statistics at a Glance. Department of Agriculture, Cooperation & Farmers Welfare Horticulture Statistics Division, Government of India..
- b) Statistical Handbook of Meghalaya. Directorate of Economics and Statistics Government of Meghalaya Shillong.

Anonymous 2018-2020. Atlasbig.com.

Anonymous. 2019. Department of Agriculture Government of Meghalaya, Shillong-Meghalaya.

Comment [U75]: Not proper citing. Goggle and find the real or original authors and acknowledge them appropriately with the period they researched these topics.

Ansari, J.K., Bose, D.K.,Jahanara. (2021) Knowledge And Adoption Behaviour Of Farmers Towards Improved Tomato Production Practices In Sonbhadra District Of Uttar Pradesh DOI: 10.47856/ijaast.2021.v08i9.009.

Hadole,S.M., Pawar, Parmeshwari, Ugale, T.B. (2017) Relationship between selected characteristics of Tomato growers and their adoption level of recommended technology in Nashik District of Maharashtra state. VL - 12 DO - 10.15740/HAS/AU/12.2/210-214

Mulewa Annand (2019) Entrepreneurial Behaviour of Tomato Producers Under National Horticulture Mission (NHM) In Dhar District of Madhya Pradesh "*IOSR Journal of Agriculture and Veterinary Science (IOSR-JAVS)* 12.10 (2019): PP- 47-4

Pole , Vinod Ramrao Kamble, V.B. 2018. Knowledge And Adoption Of Recommended Tomato Production By The Growers Vasantrya Naik Marathwada Krishi Vidyapeeth, Parbhani.

Rashmi N. and Chandrashekar, S. Vaster (2018) A Study On Knowledge, Adoption And Marketing Behaviour Of Tomato Growers In Chickaballapur District Of Karnataka.

Sangavi M,Janaki Rani A. (2021) Knowledge Level on Recommended Cultivation Practices for Tomato Growers in Madurai District. *Journal of Current Crop Science and Technology*.

Comment [U76]: Find out the peculiar referencing format/styles from the Journal you want to publish this work and follow same. E. g. APA, MLA and others