

Biotechnological Production of Carotenoid from Oleaginous Red yeast and its Applications

Abstract

The demand for carotenoids and their derivatives from natural sources is increasing rapidly due to public concern about food safety and health issues, and thus, carotenoid production from microbial fermentation is increasing significantly due to its ability to accumulate higher levels of carotene. Carotenoids, lipid-soluble pigments, are responsible for the vibrant colors in food and microorganisms. Carotenoids have the most important advantages in terms of antioxidant and anticancer activity. These possible applications are used for treating various diseases like xerophthalmia, keratomalacia, skin acne, breast cancer, and tumour formation. They are widely used in the pharmaceutical, cosmetics, and food industries. Due to the overall increase in the cost of carotenoids, carotenoids are produced in the pharmaceutical, food, and cosmetics industries through chemical synthesis or extraction from plants. The oleaginous red yeast, *Rhodotorula minuta*, is well known for producing a high yield of carotenoids with a low production cost. Over the years, these carotenoids have been produced from oleaginous red yeast, using low-cost substrates or agricultural waste for cost-effective purposes. In this paper, we highlighted the production of carotenoids from oleaginous red yeast and its applications.

Keywords: Carotenoids, *Rhodotorula*, Agricultural residue, Antimicrobial activity

1. INTRODUCTION

Color is an important attribute that determines consumers' acceptance of food and health-related drug products. Recently, the global markets for pharmaceuticals, food, and cosmetics industries have been fulfilled via synthetic color pigments. Although the application of chemically synthetic color pigments is restricted because of their toxicity [1]. Carotenoids are natural pigments responsible for the colourant in fruits, flowers, and roots that occur in photosynthetic systems of higher plants, algae, yeast, and bacteria. These carotenoids are hydrophobic in nature and are precursors of vitamin A with beneficial effects, including boosting the immune system and reducing the risk of degenerative diseases such as cardiovascular disease, cancer, skin problems, and eye deficiencies [2]. The vitamin A

precursor helps in the formation of rhodopsin, which is called a photoreceptor pigment in the retina of humans, which is important in the vision process. The deficiency of vitamin A leads to xerophthalmia, keratomalacia, and skin acne. Xerophthalmia is one of the major nutritional problems all over the world that leads to dryness of the conjunctiva, cornea on the surface of the eye, and blindness, particularly in children [3]. Keratomalacia, in which the cornea becomes hazy and develops erosions, automatically leads to major surgery. Some skin problems, i.e., acne, have been diagnosed dermatologically due to a deficiency of vitamin A. Vitamin A is also responsible for maintaining epithelial tissues, lysosome stability, and glycoprotein synthesis. Moreover, carotenoids also show anticancer activity and are important antioxidants for humans that scavenge oxygen and peroxy radicals. In addition, they are also useful in dietary supplements, food coloring agents, cosmetics, and pharmaceutical drugs. These carotenoids are further differentiated according to their cyclic structure into four types, i.e., β -carotene, γ -carotene, torulene and torularhodin.

2. TYPES OF CAROTENOIDS

2.1. β -carotene and γ -carotene

Carotenoids' derivative form, i.e., β -carotene, has an acyclic structure with a long chain that is conjugated with double bonds. They act as an essential dietary supplement for human growth, development, immune system function, and vision. They are organic and give strong red-orange color pigments to plants, fruits, and oleaginous microorganisms [4]. Due to its high antioxidant property and bioactivity, it is also widely used in the pharmaceutical industry [3]. They also act as coloring agents for drugs and ointments. In the food industry, an orange-red pigment (β -carotene) is used in non-alcoholic beverages, edible fats, cheese, pastry, and ice cream. Researchers demonstrated that β -carotene has a dual function in providing color to food while also improving its nutritional value. [The](#) cosmetics industry, it is used in sunscreen, which protects skin problems and exposure to UV radiation [1].

Commercially, these antioxidants also have the ability to reduce the risk of various skin problems and the ageing of humans. Among the numerous functions of β -carotene, the most common advantage is provitamin A supply, which helps in the embryonic development, growth, and vision of a new born child. It is also thought to be a cancer transporter gene inhibitor, which would result in anticancer activity [2].

β -carotene is produced naturally via oleaginous microorganisms by biotechnological processes. According to the studies, *Rhodotorula* species are natural β -carotene producers. For example, *R. glutinis* can produce β -carotene up to 25%–43% and torulene, up to 28%–30%. If grape skin were used as a carbon source, *R. glutinis* could produce 60–70% of the carotenoids. In some cheap carbon sources, such as cheese whey, sugarcane molasses, wheat, rice straw, etc., oleaginous yeast can also accumulate a high amount of carotenoids. Another differentiated carotenoid is γ -carotene. It is also an isomer of β -carotene, which has strong antioxidant properties and is a potential precursor of vitamin A, though less than β -carotene [5]. γ -carotene concentrations are present in orange carrots with high serum concentrations. Its structure is quite equivalent to the molar amount of β -carotene due to the presence of one molecule of biologically active retinol that gets converted after central cleavage. Like other carotenoids, it has anti-carcinogenic properties. Low or high dietary intake of γ -carotene alone may enhance immune function and lower the risk of cardiovascular disease and cancer.

2.2. Torulene and Torularhodin

Torulene is derived from γ -carotene, whereas torularhodin is formed from torulene. Both exhibit the function of vitamin A and antioxidant properties [6]. They have the capacity to neutralise free radicals and also inhibit the growth of prostate cancer and tumour formation. Torulene and torularhodin, in comparison, have higher antioxidant activity than β -carotene [7].

3. SOURCES OF CAROTENOIDS

3.1. Plants

Carotenoids are accumulated in the chloroplasts in a significant amount and are naturally synthesized by flowers, fruits, and vegetables such as lily, marigold, containing β -carotene, γ -carotene, antheraxanthin, and lutein. In fruits, i.e., grapes, mango, melon, papaya, peach, and tomatoes, lutein, β -carotene, γ -carotene, lycopene, and phytopene are present. While green vegetables such as beans, broccoli, cabbage, kale, lettuce, spinach, and yellow-orange vegetables contain β -carotene, lutein, etc., as shown in Table 1 [3]. In addition, xanthophylls, cryptoxanthin and zeinoxanthin, are also present. In purple-colored vegetables, a significant number of non-vitamin A active carotenoids are present. Saffron plants typically contain different apo-carotenoids.

Table 1: Natural Sources of carotenoids

Sources of Carotenoids	Carotenoids
Plants	
Flowers	
<i>Adonis, Chrysanthemum</i> , lily, marigold, <i>Oncidium, Osmanthus</i>	β -carotene, γ -carotene, antheraxanthin, lutein
Fruits	
Capsicum, cashew apple, citrus, grape, Mango, melon, papaya, peach, tomato	β -carotene, lutein, γ -carotene, Phytopene, lycopene
Seeds	
<i>Aradidopsis</i> , canola, maize, pumpkin, red millets, sunflowers, wheat	β -carotene, xanthophyll, lutein
Vegetables	
Green vegetables	
Beans, broccoli, cabbage, kale, lettuce, spinach etc.	β -carotene, lutein
Yellow-orange vegetables	
Carrot, corn, xanthophylls, carotene, ?	xanthophylls, lutein, β -cryptoxanthin, zeinoxanthin
Red-purple vegetables	
Tomato, watermelon, red paprika	Non-vitamin A active, capsanthin, carotenoids, lycopene
Oleaginous micro-organism	
Yeast	
Oleaginous red yeast	Torulene and Torularhodin, β -Carotene
Fungi	β -carotene, Lycopene, Astaxanthin
Micro-algae	Xanthophyll, zeaxanthin, lutein, Antheraxanthin.
Bacteria	β -carotene, salinixanthin, Astaxanthin, Zeaxanthin,
Animals	
Marine animals	β -carotene, fucoxanthin, Peridinin, Diatoxanthin, Alloxanthin, Astaxanthin
Reptiles	lutein, β -carotene

Different types of carotenoids are produced from different sources depending on their genotype, maturation phase, cultivation and climatic conditions, post-harvest and processing practices. Carotenoid-rich sources, mainly fruits and vegetables, have received a lot of attention recently due to the health-related benefits expressed after their regular consumption.

In addition to their key roles in photosynthesis, photo-protection, and response to environmental stress, carotenoids are involved in interactions with pathogens and symbiotic organisms. Moreover, these pigments play important roles in germination, photomorphogenesis, fruit development, and different signaling and regulatory processes. Carotenoids are very abundant in flowers and fruits, conferring yellow, orange, and red colors to attract insects and animals for pollination or seed dispersal [6].

3.2 Oleaginous microorganisms

Natural carotenoids are found in plants and vegetables in the form of carotene and xanthophyll. According to the research, carotenoids are now the most important feedstocks for pharmaceutical, food, and cosmetics industries, but due to limited resources and high costs, they cannot meet the demands economically [8]. Therefore, enormous efforts have been initiated to develop advanced strains for higher carotenoid production for pharmaceutical-based chemicals and food additives [9]. Oleaginous microorganisms such as fungi, bacteria, micro-algae, and yeast have the capability of accumulating higher carotenoids [8]. However, fungi, bacteria, and micro-algae have limited efficiencies in producing carotenoid due to higher costs and time-consuming problems. Oleaginous yeast have a capability of producing carotenoids up to 40% of the total biomass weight, including *Rhodospiridium/Rhodotorula* and *Sporidiobolus/Sporobolomyce*. Interestingly, the advantage of using oleaginous yeast is that it is easily scaled-up, has a higher growth rate, and can be grown on residual substrate from an agricultural source while producing lipids and carotenoids simultaneously [10]. *R. minuta* and other red yeast mainly produce β -carotene, torulene, and torularhodin. *Rhodotorula* yeast strain requires growth parameters such as carbon, nitrogen sources, low-cost substrate, and culture conditions to produce more carotenoid. In the production medium, for example, *Rhodotorula glutinis* and *Rhodotorula graminis* produce 11–15% of the total carotenoid. On the other hand, *Rhodospiridium* species, due to their flexible substrate adaptability, are considered a promising candidate for carotene production. *R. toruloides* yields γ -carotene with of 8.6 ± 0.2 mg/g DW [11]. *R. toruloides* NRRL Y-1091 also produces a high yield of carotene, with the highest carotenoid production of 24.58 ± 1.88 mg/L [12]. *R. toruloides* and *Rhodospiridium diobovatum* red yeast strains also have the capability of producing a high yield of carotenoid and its

derivatives. The composition and yield of different strains are affected by medium composition, environmental conditions, and carbon nitrogen ratio [12].

3.2.1 –*Rhodotorula minuta*

Rhodotorula minuta is a unicellular oleaginous microorganism of the basidiomycetes phylum that has distinctive yellow, orange, and red colonies. These *R. minuta* species are known for synthesizing natural carotenoids under certain cultural conditions, and the main carotenoids produced are torulene, torularhodin, and a minute quantity of β -carotene. They are important in food, cosmetic, and pharmaceutical industries [13]. The oleaginous red yeast has a faster growth rate as compared to similar oleaginous yeasts like *Lipomyces starkeyii* and *R. glutinis*. According to the studies for the production of pigments from red yeast, various dairy waste substrates, agricultural residues and molasses were used [14]. Researchers favored *R. minuta* oleaginous red yeast, one of the most efficient carotenoid producers due to its ability to grow in harsh environmental conditions where other red yeasts cannot. That's why it is a more suitable oleaginous microorganism for a biotechnological application, both in pharmaceutical and food industries on a commercial basis.

3.2.2 –*Rhodospiridium toruloides*

R. toruloides is an aerobic, dimorphic, and non-pathogenic, oleaginous pink yeast that is naturally present in different environmental sources, e.g., soil, wood pulp, conifers, seawater plants, and dry leaves. When excess carbon sources are used during cell culture growth, this oleaginous yeast can accumulate carotenoids and their derivatives to more than 30–40% of the total biomass dry weight [10, 15, 19]. There are various carbon sources such as lignocellulosic biomass, crude glycerol, vegetable and animal waste, sugarcane molasses, cheese whey, etc. *R. toruloides* is able to utilize different types of carbon sources for carotenoid production and shows robustness in terms of resistance against inhibitors produced after pretreatment of lignocellulosic biomass [16]. According to several studies, growing red yeast using cheap carbon and nitrogen sources is the most efficient nutrient for the growth of the cell [17]. They produce carotenoids at a higher level, which makes them promising feedstocks for pharmaceuticals, food colorants, and cosmetic. Therefore, many researchers are working on increasing carotenoid production using engineered strains of oleaginous yeasts [14]. For example, carob pulp syrup and sugarcane molasses were used for carotenoid and lipid production using the *R. toruloides* NCYC 921 strain, which contains a

total sugar concentration of 75 g/L and produces lipid productivity (1.90 g/L) and carotene productivity (9.79 g/L h⁻¹) [18]. In another report, at high salt concentration, *R. toruloides* increases lipid and carotenoid production processes, and it was observed that 27.2% (w/v) increased in carotenoid and 36.2% (w/v) in lipid production under osmotic stress [19]. *R. toruloides* was grown in a variety of food waste extracts to increase lipid and β-carotene production, yielding good results of 1.60 mg/L carotene and 9.6 g/L lipid production [20].

3.2.3 *Rhodotorula glutinis*

R. glutinis is an ~~another~~ important oleaginous red yeast that can accumulate a variety of valuable compounds including microbial lipid, pigmented carotenoid, and enzymes. They are non-pathogenic and mostly found in soil, fruits, dry leaves and plants, animal waste, and milk products. *R. glutinis* is aerobic and mesophilic in nature [21]. Certain carbon sources (glucose, molasses, fructose, glycerol) and nitrogen sources (ammonium sulphate, chloride, yeast extract, or monosodium glutamate wastewater) are used for lipid production and carotene. Additionally, researchers are working on *R. glutinis* to increase microbial lipid and carotene production either by improving the culture medium or by genetically modifying the *R. glutinis* strain. It was demonstrated that *R. glutinis* has the potential capability of producing lipid and carotenoid production on a large scale.

4. CHEMICAL SYNTHESIS OF CAROTENOID

Carotenoids are synthesized on an industrial scale. Among them are β-carotene (-apo-8'-carotenal -apo-8'-carotene), lycopene, xanthophyll (canthaxanthin, astaxanthin, and cytranaxanthin). These chemicals are synthesized from two reactions, i.e., Wittig reactions and Grignard compounds, for producing carotenoids and their derivatives. The Wittig reaction produces alkenes and triphenylphosphine oxides with the chemical reaction of aldehydes or ketones with triphenylphosphine salt. In addition, this salt is called a "witting reagent" in the reaction. The Wittig reaction is most often used to combine di-aldehyde or ketone molecules containing 10 carbon atoms and two phosphonium salt molecules, each containing 15 carbon atoms, as shown in Fig. 1. Then the reaction is subjected to an isomerisation reaction, producing compounds of 40 carbon atoms including β-carotene, astaxanthin or lycopene.




Fig.1: Wittig condensation reaction of carotenoid

The other chemical synthesis of carotenoids is the grignard reaction, where the addition of organomagnesium halide, also known as the grignard reagent, is shown in Fig. 2. In this chemical reaction, one di-ketone molecule and two methanol molecules combine together and,

thereafter, carotenoid is produced with 40 carbon [6]. Among other methods of producing β -carotene, the selective condensation reaction of carbonyl compounds and the homo-dimerisation reaction with dehydration and elimination can also produce the carotenoid. However, the chemical synthesis of carotenoids is harmful for humans as they are toxic and unable to be consumed in food and feed. As a result, researchers are looking for another

method of producing natural carotenoids from oleaginous microorganisms capable of accumulating carotenoid and its derivatives, such as β -carotene, α -carotene, lutein, and so on.




Fig. 2: Grignard reaction of carotenoids

5. BIOSYNTHETIC PATHWAYS OF CAROTENOIDS

Carotenoids are terpenoid compounds with 40 carbon atoms based on their structure. β -carotene (γ -carotene, torulene, torularhodin) and xanthophyll (salinixanthin, Astaxanthin, Zeaxanthin, Canthaxanthin) are the most common and natural carotenoids [1]. Carotenoids are synthesized by prokaryotes and eukaryotic microorganisms, and biosynthesis occurs in two different pathways: via the mevalonate pathway (MVA) and the methylerythritol 4-phosphate pathway (MEP) [6]. Commonly, eukaryotes such as yeast, fungi, and micro-algae synthesize carotenoids by MVA pathways with the condensation of 3 molecules of acetyl-CoA to synthesize mevalonaic acid via acetoacetyl-CoA and get converted into 3-hydroxy-3-methyl-glutaryl (HMG-CoA) with the help of the enzyme hydroxymethylglutaryl-CoA synthase. This HMG-CoA with hydroxymethylglutaryl-CoA reductase enzyme gets converted into Mevalonaic acid and biocatalytic reactions mediated by mevalonate kinase form mevalonate 5 phosphate (MVP), which in turn originates into the mevalonate pyrophosphate (MVPP) mediated by phosphomevalonate kinase, and then the formation of IPP occurs as shown in Fig. 3. Iso-pentenyl pyrophosphate (IPP), isomerized to dimethylallyl diphosphate (DMAPP). After DMAPP, the prenyl transferase enzyme converts it into geranyl pyrophosphate to geranylgeranyl pyrophosphate, which further gives phytoene, which is mediated by phytoene desaturase enzyme to form neurosporene, further classified into lycopene and β -zeacarotene, which gets converted into γ -carotene to β -carotene, torulene, and torularhodin, a carotenoid compound [6]. While in prokaryotes, the MEP pathways create IPP and DMAPP through a primary condensation reaction between pyruvate and glyceraldehyde-3-phosphate. It begins with the glycolysis cycle, where a 6 carbon source (glucose) is converted into 2 pyruvate, which enters the mitochondrial matrix from the cytoplasm of the cell. This acetyl-CoA is further oxidized and moves towards the cytoplasm of the cell for catalyzing the first reaction with glyceraldehyde-3 phosphate in MEP pathways, where 1-deoxy-D-xyl-Phosphate. The chain elongation occurs through the condensation mechanism of IPP to DMAPP. The short-chain-length of DMAPP is mediated by prenyl transferase synthesis, which forms geranyl pyrophosphate and geranylgeranyl pyrophosphate, which are the precursors of carotenoids. Colorless C₄₀ phytoene is formed by the condensation of two molecules of geranylgeranyl pyrophosphate, which is catalyzed by phytoene synthase. Subsequent desaturation of phytoene by phytoene desaturase further produces neurosporene, red-coloured lycopene and β -Zeacarotene. The synthesis of cyclic carotenoids of lycopene and the dehydration of β -Zeacarotene further produces β -carotene,

toluene, and torularhodin production, which in turn can be converted into other types of carotenoids derivatives.




Fig. 3: Biosynthetic pathways of carotenoid production by MVA and MEP pathway

6. METHODS OF EXTRACTION OF CAROTENOIDS

6.1 Organic-solvent extraction

Carotenoids are hydrophobic in nature and are usually extracted from organic solvents. Generally, organic solvents are of two types: polar solvents and non-polar solvents. Polar solvents, i.e., acetone, ethanol, and ethyl acetate, for extracting derivatives of carotenoids, are easy and simple methods for laboratory scale extraction applications. While non-polar solvents like petroleum, hexane, and tetrahydrofuran are excellent choices for extracting precursors of carotene. However, excess use of toxic solvents should usually be avoided because they become toxic for the environment and can cause harm to human health [4].

6.2. Saponification

Mainly, higher production of carotene is found in oleaginous microorganisms after plants, and this carotene exists in free form and has triacylglyceride biosynthesis ability. In order to release carotenoids in ester form, saponification reactions containing alcohol and KOH or NaOH are used. This alcohol saponifies the compound and then removes the ester bond to create functional groups. The saponification reaction requires specific conditions, i.e., temperature, time, and alkali concentration, which are the main factors that affect the yield of carotenoids [1]. In Liu et al.'s (2021) [4] report, it was found that the production of carotenoids in *R. toruloides* by using a saponification method in which alcohol and KOH had a higher recovery rate ($78.7 \pm 2.4\%$) [4].

6.3. Microwave and ultrasound assisted extraction

A variety of combined extraction technologies have been developed in recent years. Microwave and ultrasonic-assisted extraction technologies were used for direct and rapid extraction of carotenoids and their precursors in order to overcome the problems of low yield, toxicity, and high extraction cost of carotenoid production. However, single microwave or ultrasonic extraction can no longer meet the needs of production, so a variety of combined extraction technologies have been developed. For example, it was reported that in *Phaffia rhodozyma*, lactic acid was used to destroy the cell clusters and then carotene derivatives were extracted via ultrasonic extraction, which was then used further in the food and feed industries [22].

6.4. Green solvents as ionic liquids

The organic solvents, for example, methanol, ethanol, acetone, hydrochloric acid, etc., are extremely flammable, volatile, and toxic, and can also cause environmental pollution. Thus, the use of these organic reagents is avoided in the extraction of carotenoids. Ionic liquids (ILs) are a new type of green solvent with many characteristics, including environmentally friendly, non-volatility, non-flammable, thermal stability, control of led miscibility, and the ability to maintain the composition of anionic and cationic salts. Lewis acidic ionic liquid and bronsted acidic ionic liquid are two types of liquids that are commonly used [4].

6.5. Enzyme-assisted extraction

Conventionally, solvent extraction methods were used for extracting the carotenoids, but due to heat production during this extraction process, it affected the stability of carotenoids. In order to obtain a high yield of carotene precursor and improve its stability, enzyme pretreatment was used. They are non-toxic when used in food and feed industries as compared to organic solvents. Since enzymes are environmentally friendly, renewable, and increase production rate and reduce the use of solvents, their disadvantages are that they are expensive and require optimal operating conditions, which makes researchers find an alternative solution [1].

6.6. Supercritical fluid extraction

The new technology, named "supercritical fluid extraction technology," has been developed due to the high demand for carotenoids and their precursors in the last few years. Supercritical fluid is the most preferred extraction method in the pharmaceutical and food industries as it uses non-toxic solvents, namely water and carbon dioxide. This carbon dioxide has an ambient critical temperature range of 31-°C and carotenoids can be extracted from microorganisms under 35-°C. The density of supercritical CO₂ is around 200 bar, which is close to hexane and the properties are similar for extraction. It is also a green alternative to conventional carotenoids extraction methods. More importantly, supercritical fluid extraction technology has higher permeability and diffusibility, so it can obtain carotenoids faster and higher.

7. FACTOR AFFECTING THE PRODUCTION OF CAROTENOIDS

7.1 Low cost feedstock

Earlier, β -carotene was extracted from carrots and fruits of palm trees, but for commercial purposes, it could not meet the higher demand in humans [1, 23]. However, the alternative, i.e., oleaginous microorganisms, is used for carotenoid production, but due to its expensive carbon sources, large area for cultivation and limited growth, it cannot fulfil the demands. Recently, oleaginous red yeasts have been shown to be potentially more beneficial and effective for carotenoid production as compared to other microorganisms. *Rhodospiridium toruloides*, in particular, has the ability to accumulate natural β -carotene, torulene, and torularhodin [5, 24]. *R. toruloides* can assimilate various lignocellulosic substrates for carotenoids production, including corn stover, sugarcane bagasse, rice, wheat straw, banana peels, and food wastes [25]. In another study, camelina meal hydrolysates were used as a lignocellulosic biomass for the cultivation of carotenoids by *Rhodospiridium toruloides* in culture medium, and it was found to produce 13 ± 2.6 mg/L of carotenoids [Breath et al. 2020]. The wheat straw hydrolysate was also used as a carbon source for cultivating *R. toruloides* in culture medium and, after pretreatment, it produced 24.58 ± 1.88 mg/L of carotenoid [26].

7.2. Light

Light intensity is the most important parameter in the synthesis of carotenoids in oleaginous microorganisms (fungi, microalgae, yeast, and bacteria). The biosynthetic pathway and production of carotenoids are affected by light and, more importantly photo-induction, which enhances the carotenoids yields by promoting cell growth density and the activity of enzymes involved in carotenoid biosynthesis. According to the report, in *R. glutinis*, when light facilitated the yield of —torularhodin increases by 18% and the production of β -carotene increases by up to 14%. Carotenoid production is affected by white light, mostly UV-light or LED light. In addition, carotenoid production in the *R. mucilaginosa* strain showed a high yield of carotenoid in the presence of white light [27]. They also stated that the higher production of torularhodin in yeast improves with UV light exposure. Another report showed that β -carotene production results in a concentration of $24.6 \mu\text{g}^{-1}$ under two LED lights, whereas without light intensity, the concentration decreases to $14.69 \mu\text{g}^{-1}$. The synthesis of torularhodin increases when yeast cells are exposed to white light. When *R. glutinis* is

exposed to white light irradiation, it increases the carotenoid production, mostly torularhodin and torularhodin [3]. However, increases in microbial growth and the activities of enzymes are directly related to the increase in carotenoid biosynthesis [27].

7.3. Temperature and pH

Temperature is the most important factor for the biosynthesis of carotenoid pigments. It influences cell growth, enzyme concentrations in cell metabolism [28]. At low temperatures, *R. glutinis* and *R. vulgaris* increase β -carotene content [26]. In another report, the synthesis of torularhodin depends on the corresponding enzyme when it is less active than β -carotene synthesis at low temperatures [4]. The increase or decrease in temperature mostly depends on the growth of oleaginous microorganisms and their synthesis of carotenoid. Temperature increases from 29°C to 30°C increase the overall production of carotenoid and biomass yield in *R. glutinis* [4]. Temperature ranges of 20°C to 30°C are best suited for *Rhodotorula* species growth, according to the report, while higher temperatures, i.e., 32°C or above, reduce the amount of carotenoids synthesis [5]. The carotenoid pigments, i.e., β -carotene and torulene production, decrease at 31°C temperature. Whereas, torularhodin production increases at 31°C.

The pH range is also an important parameter for optimizing the growth of oleaginous yeast for carotenoid production. According to the report, at an initial pH of 5, under various temperature ranges, i.e., (22°C to 28°C), it exhibits the maximum amount of carotenoid pigments in *Rhodotorula* yeast. However, at higher temperatures, i.e., above 30°C, it lowers the yield of carotenoid. It was stated that the biomass growth of oleaginous yeast was influenced when the pH range was between 5 and 7. The highest carotenoid production was at pH 5. At pH 4, however, carotenoid production decreases. The effect of pH on the yield of carotenoids derived from *R. glutinis* was greatest when the pH of the culture medium was 3.0. The carotenoid yield was 115.8 μg^{-1} DW, while at pH 4.0–7.0, the yield of carotenoid was very high (191.7–202.9 μg^{-1} DW) [5].

7.4. Carbon and nitrogen sources (C/N ratio)

Biosynthetic metabolism of *Rhodotorula* species depends on the carbon source and nitrogen sources in the culture medium. The carbon to nitrogen ratio is one of the major factor that assimilate the production of carotenoid pigments. According to the investigation, it was stated that high C/N ratios are advantageous for both biomass and carotenoid yield. Thus, different

experiments are performed to observe the impact of different C/N ratios, i.e., (C/N 70 and 120), glucose concentration, initial pH value, and nitrogen content on the microbial cell growth for a higher yield of carotenoid pigments by *R. glutinis* [28]. Furthermore, the results showed that carotenoid production increases at high C/N ratios when the pH value is greater than (> 2), whereas it decreases at low pH values (< 2). Since the optimum pH levels for carotenoids are around pH 6 [29]. Continuous optimization of the cultural medium in research found that the continuous supplement of organic and inorganic nitrogen sources promoted the production of carotenoids in oleaginous yeast *R. glutinis* (β -carotene, torularhodin, and torulene) by about 25.83%, 11.88%, and 24.50%, respectively.

7.5. Metal ions

Many metal ions such as Ba^{2+} , Fe^{2+} , Mg^{2+} , Ca^{2+} , Zn^{2+} are capable of accumulating carotenoid pigment in *Rhodotorula* species. According to the report, the effect of metal ions in *R. graminis* has been investigated and shows that Al^{3+} and Zn^{2+} metal ions increase the production of both γ -carotene and β -carotene. Metal ions Mn^{2+} and Zn^{2+} , on the other hand, decreases the production of bio-pigments such as torulene and torularhodin. In another study, after adding 0.1 g/L of calcium chloride, magnesium sulphate, or ferric sulphate to the production medium of *Rhodotorula* species, it had a slight influence on cell growth when compared to the control. The carotenoid production with added metal ions showed excellent growth compared to those without metal ions. As per the report, the most suitable metal ion is Fe^{2+} , whose carotenoid concentration was up to 37.6%. Magnesium and calcium ions, increase carotenoid concentration in *R. glutinis* by 21.4% and 9.2%, respectively. Whereas, ferric and magnesium ions increase the total carotenoid concentration by 23.6% and 9.1%, respectively. β -carotene increases after adding metal ions to cell concentration, while torularhodin decreases after involvement of metal ions. The torulene pigment also increases when Fe^{2+} and Mg^{2+} ions are added to the cell, but calcium ions decrease the growth of torulene [4]. The use of metal ions in all microbial growth has a relatively larger impact on increasing carotenoid production. For example, sodium, calcium, and zinc. While potassium and magnesium are the active groups of certain enzymes in carotenoid biosynthetic pathways, they have the capability of regulating cell metabolism and cell membrane permeability in microorganisms [3, 30].

8. APPLICATION OF CAROTENOIDS

8.1 Antimicrobial activity

The secondary metabolites of a few pigmented yeast strains have been observed for their antibacterial and antifungal activities. In general, the antibacterial activity was assessed in a qualitative or quantitative method. The carotenoids were extracted from *R. mucilaginosa*, *R. rubra*, and *R. glutinis*, sp, where the major concentrations of carotenoids were toluene and torularhodin. *Bacillus*, *Salmonella*, *Pseudomonas*, and *Escherichia* are of clinical interest for antimicrobial tests.

The MIC minimum inhibitory concentration was tested against *Salmonella choleraesuis* ATCC 6539, *Staphylococcus aureus* ATCC 13,565, *Pseudomonas aeruginosa* ATCC 25,442, *Listeria monocytogenes* ATCC 19,117, and *Escherichia coli* ATCC 11,229 for carotenoids from *R. mucilaginosa* CCMA 0156 [31]. One of the most relevant applications of purified carotenoids as antimicrobials is their use as antimicrobial coatings in some titanium-medical products, thus improving the effectiveness of torularhodin in implants and protecting humans against bacteria. Soliman et al. (2018) [31] showed a MIC of 1 g/mL of silver nanoparticles added to *Rhodotorula* sp [31]. ATL72 culture it shows antimicrobial activity against *E. coli* and *Bacillus* sp Also, Cunha et al. (2018) [32] showed the lowest MIC reported for the same type of silver nanoparticles added to yeast carotenoids cultures and showed antimicrobial activity against *Candida parapsilosis* [32].

8.2. Anticancer activity

Very promising findings were obtained for inhibiting different types of cancerous cells in humans using carotenoids and their derivatives. α -carotene, β -carotene, lycopene, torulene, torularhodin, and some other derivatives, exhibit anticancer activity for prostate, breast, colon, lung, oral, gastric, and skin cancers, in addition to hepatoma, leukemia, etc. Certain antioxidant response elements are activated within cells to inhibit the androgen receptor activity of cancer cells. These carotenoids are lycopene, phytoene, and phytofluene, and they are further used as therapies for prostate and breast cancer cells. The experiments were performed where ER-negative Hs578T, MDA-MB-231 human breast cancer cells, and oestrogen receptor (ER) positive MCF-7 were treated with carotenoids. Among them, beta-carotene significantly reduced the growth of Hs578T cells and MCF-7, and this lycopene inhibited the growth of MCF-7, MDA-MB-231 cells, and MCF-7 [6]. According to the report, Astaxanthin, a derivative of carotenoids, also inhibits the growth of breast cancer cell

lines, and it is found that oestrogen receptors are the major key factor. The carotene-enriched food was found to be effective against lung and stomach cancer. More recently, tumor metastasis mechanisms in humans are inhibited by lycopene by proliferating diverse cancer cell lines and slowing cell-cycle progression. Some recent reviews discuss the mechanisms of carotenoid cell-cycle arrest, anticancer activity, apoptosis-inducing effect, and anti-metastasis effect. They also act as chemo-protective agents against malignant transformation and cellular mutagenesis. β -carotene, α -carotene, and canthaxanthin can inhibit 3-methylcholanthrene against malignant transformation or X-ray treatment in the fibroblast cell line [6].

8.3. Antioxidant and anti-inflammatory activity

The world is facing severe problems in chronic health complications like cancer, cardiovascular diseases, diabetes, hypertension and other malfunctions. The primary nutrients provided by fruits and vegetables can be the solution to health-related problems as they contain carotenoid pigments. These carotenoids are rich in antioxidants, which is the main factor responsible for their protective effects against chronic diseases. Antioxidants provide additional protection against increased oxidative stress and help in scavenging two of the reactive oxygen species: singlet molecular oxygen (1O_2) and peroxy radical. β -carotene plays a major role in radical transfer chains, as it has a triplet energy level close to that of singlet oxygen molecules and can transfer energy to the cells. In addition, zeaxanthin, cryptoxanthin, and γ -carotene also have an ability to inhibit reactive oxygen species (ROS) in human retinal cells or cancerous cells [6]. These antioxidants have the advantages of lowering the amount of free radicals in our bodies, preventing direct cell damage caused by the chain reaction, and reducing the signs of ageing by preventing oxidation in our skin cells. Carotenoid activity is important for wound healing procedures in the human body and has the capability of inhibiting various carcinogenic diseases and may act as a drug or substance that may activate the immune response in an organism. The immune responsive compounds in the human body, i.e., cytokinins and interleukins, are produced by B lymphocytes, keratinocytes, T lymphocytes, macrophages, and the endocrine system, which release enzymes and antibodies that perform inflammatory action. Interleukin-1 and interleukin-2 are cytokine groups that aid in the regulation of immune and anti-inflammatory responses in human organs in response to infections, as well as acting as mediators to induce healing. They also control the biochemical pathway and prevent the diseases from further spreading [6]. The

study revealed that lycopene, a carotenoid precursor, acts as a mediator that limits the release of excess glucose metabolism in human cells by effectively reducing the oxidation and degradation of free radicals and non-free radicals, i.e., ROS, H₂O₂, O₂, R., etc., that results in negative effects in the mitochondria of human retinal cells. However, these carotenoids block the effects of ROS with the help of antioxidant proteins (superoxide dismutase (SOD), catalase, peroxiredoxin (Prx), glutathione reductase (GR), and thioredoxin reductase (TR)), which then activates the proliferator-activated receptor in bone health, which directly influences anti-inflammatory transcription factors and decreases the risk of osteoporosis. According to the report, the osteoporosis risk in postmenopausal women can be decreased by lycopene. The same report identifies the effects as both lycopene and β-carotene, and β-cryptoxanthin are involved in bone health.

9. CONCLUSIONS

Carotenoids are beneficial color pigments that protect against chronic diseases and enhance the immune system of humans. There is a keen interest in the methods of obtaining carotene and its derivatives through natural sources for the food, cosmetic, and pharmaceutical industries. For many years, they have been widely used as food additives, anti-inflammatory drugs, cosmetics, and natural food colorants. As a matter of concern due to global warming, environmental pollution and health issues, demands for natural carotenoids in the global markets are also increasing. Due to high demand, the main quantity of these compounds are naturally synthesized from fruits and vegetables through physicochemical methods. However, due to high demand, it cannot meet the demands. Currently, the production of carotenoids and their pre-cursors is more expensive than synthetic forms. Therefore, lots of new approaches have recently been developed. Oleaginous microorganisms, especially *Rhodotorula* red yeast (*Rhodotorula minuta*, *Rhodotorula glutinis*, *Rhodospiridium* sp, etc.), are capable of accumulating natural carotenoids. These oleaginous red yeasts produce a high yield of β-carotene, lutein and compete with the composition of carotenoids produced from fruits and vegetables. However, the production of carotenoids from oleaginous yeast requires carbon sources, and because of that, an alternative lignocellulosic biomass has been used. Therefore, researchers are heading towards genetic modification or gene modelling of yeast to gain a high yield of carotenoids.

REFERENCES

1. Saini RK, Keum YS. Microbial platforms to produce commercially vital carotenoids at industrial scale: an updated review of critical issues. *J. Ind. Microbiol. Biotechnol.* (2018); 46:657–674. doi:10.1007/s10295-018-2104-7.
2. Da Silva SRS, Stamford TCM, Albuquerque WWC, Vidal EE, Stamford TLM. (2020). Reutilization of residual glycerin for the produce β -carotene by *Rhodotorula minuta*. *Biotechnol Lett* 2020;42(3):437–443. doi:10.1007/s10529-020-02790-8.
3. Tang W, Wang Y, Zhang J, Cai Y, He Z. Biosynthetic Pathway of Carotenoids in *Rhodotorula* and Strategies for Enhanced Their Production. *J. Microbiol. Biotechnol.* (2019); 29:(4), -507–517 <https://doi.org/10.4014/jmb.1901.01022>.
4. Liu C, Hu B, Cheng Y, Go Y, Yao W, Qian H. *Carotenoids from fungi and microalgae: A review on their recent production, extraction, and developments. Bioresource Technology.* 2021;337, 125398. doi:10.1016/j.biortech.2021.125398.
5. Kot AM, Blazejak S, Gientka I, Kieliszek M, Brys J. (2018). Torulene and torularhodin: “new” fungal carotenoids for industry? *Microb Cell Fact.* 2018;17(1):49. doi:10.1186/s12934-018-0893-z.
6. Igreja WS, de Andrade MF, Lopes AS, Chiste RC. Biotechnological Production of Carotenoids Using Low Cost-Substrates Is Influenced by Cultivation Parameters: A Review, *Int J Mol Sci.* 2021; 22(16):8819. doi: 10.3390/ijms22168819.
7. Du C, Guo Y, Cheng Y, Han M, Zhang W, Qian H. Torulene and torularhodin, protects human prostate stromal cells from hydrogen peroxide-induced oxidative stress damage through the regulation of Bcl-2/Bax mediated apoptosis. *Free. Radic. Res.* 2017; 51(2), 113–123. doi:10.1080/10715762.2017.1285024.
8. Tang W, Wang Y, Zhang J, Cai Y, He Z. Biosynthetic Pathway of Carotenoids in *Rhodotorula* and Strategies for Enhanced Their Production. *J. Microbiol. Biotechnol.* 2019; 29:(4), -507–517 <https://doi.org/10.4014/jmb.1901.01022>.
9. Garay LA, Sitepu IR, Cajka T, Chandra I, Shi S, Lin T et al. Eighteen new oleaginous yeast species. *J. Ind. Microbiol. Biotechnol.* 2016; 43(7): 887–900. doi:10.1007/s10295-016-1765-3.

10. Bertacchi S, Bettiga M, Porro D, Branduardi P. Camelina sativa meal hydrolysate as sustainable biomass for the production of carotenoids by *Rhodospiridium toruloides*. *Biotechnol Biofuels*. 2020; 13(1):47. doi:10.1186/s13068-020-01682-3.
11. Husseiny SM, Abdelhafez AA, Ali AAA, Sand HM. Optimization of β -Carotene Production from *Rhodotorula glutinis* ATCC 4054 Growing on Agro-industrial Substrate Using Plackett–Burman Design. *Proc. Natl. Acad. Sci., India, Sect. B Biol. Sci.* 2017;135: 117-128. doi:10.1007/s40011-017-0908-2.
12. Liu Z, Feist AM, Dragone G, Mussatto SI. Lipid and carotenoid production from wheat straw hydrolysates by different oleaginous yeasts. *J. Clean. Prod.* 2019: 119308. doi:10.1016/j.jclepro.2019.119308.
13. Seveiri RM, Hamidi M, Delattre C, Rahmani B, Darzi S, Pierre G. et al. Characterization of the exopolysaccharides from *Rhodotorula minuta* IBRC-M 30135 and evaluation of their emulsifying, antioxidant and anti proliferative activities. *Med. Sci.* 2019; 23(97): 381–389. <http://eprints.skums.ac.ir/id/eprint/7707>.
14. Lyman, M., Urbin, S., Strout, C., & Rubinfeld, B. The Oleaginous Red Yeast *Rhodotorula/Rhodospiridium*: A Factory for Industrial Bioproducts. *Yeasts Biotechnol.* 2019. doi:10.5772/intechopen.84129
15. Park YK, Nicaud JM, Ledesma-Amaro R. The Engineering Potential of *Rhodospiridium toruloides* as a Workhorse for Biotechnological Applications. *Trends Biotechnol.* 2018; 36(3):304–317. doi:10.1016/j.tibtech.2017.10.013.
16. Singh G, Sinha S, Bandyopadhyay KK, Lawrence M, Paul, D. Triaxial growth of an oleaginous red yeast *Rhodospiridium toruloides* on waste “extract” for enhanced and concomitant lipid and β -carotene production. *Microb Cell Fact.* 2018; 17(1):182. doi:10.1186/s12934-018-1026-4.
17. Fei Q, O’Brien M, Nelson R, Chen X, Lowell A, Dowe N. Enhanced lipid production by *Rhodospiridium toruloides* using different fed-batch feeding strategies with lignocellulosic hydrolysate as the sole carbon source. *Biotechnol Biofuels*. 2016;9:130 doi:10.1186/s13068-016-0542-x.
18. Freitas C, Parreira TM, Roseiro J, Reis A, da Silva TL. Selecting low-cost carbon sources for carotenoid and lipid production by the pink yeast *Rhodospiridium toruloides*

- NCYC 921 using flow cytometry. *Bioresour Technol.* 2014;158, 355–359. doi:10.1016/j.biortech.2014.02.07.
19. Pham KD, Shida Y, Miyata A, Takamizawa T, Suzuki Y, Ara S et al. Effect of light on carotenoid and lipid production in the oleaginous yeast *Rhodospiridium toruloides*. *Biosci. Biotechnol. Biochem.* 2020;1–12. doi:10.1080/09168451.2020.1740581.
 20. Kot, A. M., Błażej, S., Kurcz, A., Gientka, I., & Kieliszek, M. *Rhodotorula glutinis*—potential source of lipids, carotenoids, and enzymes for use in industries. *Applied Microbiology and Biotechnology.* 2016;100(14):6103–6117. doi:10.1007/s00253-016-7611-8.
 21. Karamerou EE, Theodoropoulos C, Webb C. A biorefinery approach to microbial oil production from glycerol by *Rhodotorula glutinis*. *Biomass and Bioenergy.* 2016; 89:113–122. doi:10.1016/j.biombioe.2016.01.007.
 22. Cheng XY, Xiong YJ, Yang MM, Zhu MJ. *Preparation of astaxanthin mask from Phaffia rhodozyma and its evaluation.* *Process Biochem.* 2018;79:195-202. doi:10.1016/j.procbio.2018.12.027.
 23. Park YK, Nicaud JM, Ledesma-Amaro R. The Engineering Potential of *Rhodospiridium toruloides* as a Workhorse for Biotechnological Applications. *Trends Biotechnol.* 2018; 36(3):304–317. doi:10.1016/j.tibtech.2017.10.013.
 24. Uprety BK, Dalli SS, Rakshit SK. Bioconversion of crude glycerol to microbial lipid using a robust oleaginous yeast *Rhodospiridium toruloides* ATCC 10788 capable of growing in the presence of impurities. *Energy Conversion and Management.* 2017;135, 117–128. doi:10.1016/j.enconman.2016.12.071 .
 25. Bertacchi S, Bettiga M, Porro D, Branduardi P. Camelina sativa meal hydrolysate as sustainable biomass for the production of carotenoids by *Rhodospiridium toruloides*. *Biotechnol Biofuels.* 2020; 13(1):47. doi:10.1186/s13068-020-01682-3.
 26. Mata-Gomez LC, Montanez JC, Mendez-Zavala A, Aguilar CN. Biotechnological production of carotenoids by yeasts: An overview. *Microb. Cell Fact.* 2014; 13, 1–11. <https://doi.org/10.1186/1475-2859-13-1>.
 27. Zhang Z, Zhang X, Tan T. Lipid and carotenoid production by *Rhodotorula glutinis* under irradiation/high-temperature and dark/low-temperature cultivation. *Bioresour. Technol.* 2014; 157:149–153. doi:10.1016/j.biortech.2014.01.039.

28. Kot AM, Błazejak S, Kieliszek M, Gientka I, Brys J et al. Effect of initial pH of medium with potato wastewater and glycerol on protein, lipid and carotenoid biosynthesis by *Rhodotorula glutinis*. *Electronic Journal of Biotechnology*. 2017; 27: 25–31. doi:10.1016/j.ejbt.2017.01.007.
29. Yolmeh M, Khomeiri M. Using physical and chemical mutagens for enhanced carotenoid production from *Rhodotorula glutinis* (PTCC 5256). *Biocatal. Agric. Biotechnol.* 2016; 8:158–166. doi:10.1016/j.bcab.2016.09.004
30. Rapoport A, Guzhova I, Bernetti L, Buzzini P, Kieliszek M, et al. *Carotenoids and Some Other Pigments from Fungi and Yeasts. Metabolites.* 2021;11(2), 92. doi:10.3390/metabo11020092.
31. Soliman H, Elsayed A, Dyaa A. Antimicrobial activity of silver nanoparticles biosynthesised by *Rhodotorula* sp. strain ATL72. *Egypt. j. basic appl. sc.* 2018; 5(3): 228-233. doi:10.1016/j.ejbas.2018.05.005
32. da Frota SM, Cunha FA, Cunha MDCDSO, Martins RT, Menezes EA, Fachine PBA. Synergistic Effect of Polyene Antifungals and Silver Nanoparticles Against *Candida Parapsilosis*. 2018. *J. Antibiot. Res.* 2018;2(1).
33. Shin J, Song MH, Oh JW, Keum YS, Saini RK. Pro-Oxidant Actions of Carotenoids in Triggering Apoptosis of Cancer Cells: A Review of Emerging Evidence. *Antioxidants.* 2020; 9(6), 532. doi:10.3390/antiox9060532

UNDER PEER REVIEW